
On the Generalization Power of the Overfitted Three-Layer Neural Tangent Kernel Model

Peizhong Ju
Department of ECE
The Ohio State University
Columbus, OH 43210
ju.171@osu.edu

Xiaojun Lin
School of ECE
Purdue University
West Lafayette, IN 47906
linx@purdue.edu

Ness B. Shroff
Department of ECE
The Ohio State University
Columbus, OH 43210
shroff.11@osu.edu

Abstract

In this paper, we study the generalization performance of overparameterized 3-layer NTK models. We show that, for a specific set of ground-truth functions (which we refer to as the “learnable set”), the test error of the overfitted 3-layer NTK is upper bounded by an expression that decreases with the number of neurons of the two hidden layers. Different from 2-layer NTK where there exists only one hidden-layer, the 3-layer NTK involves interactions between two hidden-layers. Our upper bound reveals that, between the two hidden-layers, the test error descends faster with respect to the number of neurons in the second hidden-layer (the one closer to the output) than with respect to that in the first hidden-layer (the one closer to the input). We also show that the learnable set of 3-layer NTK without bias is no smaller than that of 2-layer NTK models with various choices of bias in the neurons. However, in terms of the actual generalization performance, our results suggest that 3-layer NTK is much less sensitive to the choices of bias than 2-layer NTK, especially when the input dimension is large.

1 Introduction

Neural tangent kernel (NTK) models (Jacot et al., 2018) have been recently studied as an important intermediate step to understanding the exceptional generalization power of overparameterized deep neural networks (DNNs). Deep neural networks (DNNs) usually have so many parameters that they can perfectly fit all train data, yet they still have good generalization performance (Zhang et al., 2017; Advani et al., 2020). This seems contradicting to the classical wisdom of “bias-variance-tradeoff” in the statistical machine learning methods (Bishop, 2006; Hastie et al., 2009; Stein, 1956; James & Stein, 1992; LeCun et al., 1991; Tikhonov, 1943). To understand this distinct behavior of DNNs, a recent line of work studies the so-called “double-descent” phenomenon, beginning with overfitted linear models. These results on linear models suggest that the test error indeed decreases again in the overparameterized region, as the model complexity increases beyond the number of samples (Belkin et al., 2018, 2019; Bartlett et al., 2020; Hastie et al., 2019; Muthukumar et al., 2019; Ju et al., 2020; Mei & Montanari, 2019). However, these studies use linear models with simple features such as Gaussian or Fourier features, and hence they fail to capture the non-linearity in neural networks. In contrast, NTK models adopt features generated by non-linear activation functions (i.e., neurons of DNNs), and thus they can be viewed as an intermediate step between simple linear models and DNNs. Along this line, the work in Ju et al. (2021) studies 2-layer NTK models, and shows that the 2-layer NTK model indeed exhibits better and different descent behavior in the overparameterized region, which might be closer to that of an actual neural network.

Motivated by Ju et al. (2021), it is of great interest to understand whether similar insights extend to deeper NTK models. In particular, in this paper we study NTK models with 3 layers. Although

both 2-layer and 3-layer NTK models share similar assumptions (e.g., trained weights do not change much from initialization, and features are linearized around the initial state), their difference in structure leads to completely different feature formation. Compared with 2-layer NTK models that only contain one hidden-layer of neurons, 3-layer NTK models have two hidden-layers, which interact in more complex ways not observed in 2-layer NTK models. Specifically, let p_1 and p_2 denote the number of neurons in the two hidden layers. Then, the ultimate features of the 3-layer NTK models depend on both p_1 and p_2 . This dependency leads to the following questions. First, the width of which layer is more important in governing the descent behavior, p_1 or p_2 ? Further, to get better descent behaviors, should p_1 and p_2 grow at the same speed, or should one of them grow faster than the other? Second, do 3-layer NTK models have any performance advantage over 2-layer NTK models?

To answer these questions, in this paper we study the generalization performance of overfitted min- ℓ_2 -norm solutions for 3-layer NTK models where the middle layer is trained. For a set of learnable functions (which we refer to as the “learnable set”), we provide an upper bound on the test error for finite values of p_1 and p_2 . To the best of our knowledge, this upper bound is the first result that can reveal the dependency of the descent behavior on p_1 and p_2 separately. We then compare 3-layer NTK with 2-layer NTK with respect to the corresponding learnable set and the actual generalization performance. Our comparison reveals several important differences between 3-layer NTK and 2-layer NTK, in terms of the descent behavior, the size of the learnable set, and the sensitivity of the generalization performance to the choice of bias of the neurons.

Analyzing the Generalization Error: First, we show that the generalization error (denoted by the absolute value of the difference between the model output and the ground-truth for a test input) is upper bounded by the sum of several terms on the order of $O(1/\sqrt{n})$ (n denotes the number of training data), $O(1/p_2)$ (p_2 denotes the number of neurons in the second hidden-layer), $O(\sqrt[4]{\log p_1/p_1})$ (p_1 denotes the number of neurons in the first hidden-layer), plus another term related to the magnitude of noise. Similar to 2-layer NTK (Arora et al., 2019; Ju et al., 2021; Satpathi & Srikant, 2021), our upper bound suggests that when there are infinitely many neurons, the generalization error decreases with the number of samples n at the speed of \sqrt{n} and will approach zero when $n \rightarrow \infty$ in the noiseless situation. Further, the noise term will not explode when the number of neurons goes to infinity, which is also similar to that for 2-layer NTK. However, our upper bound also reveals new insights that are different from the results for 2-layer NTK. Specifically, our upper bound decreases slower with respect to the number of neurons in the first hidden-layer p_1 at the speed of $\sqrt[4]{(\log p_1)/p_1}$, and decreases faster with respect to the number of neurons in the second¹ hidden-layer p_2 at the speed of $1/\sqrt{p_2}$. Further, our upper bounds hold regardless of how fast p_1 and p_2 increase relative to each other (e.g., they could increase at the same speed, or one could increase faster than the other).

Characterizing the Learnable Set: We then show that, even if we only train the middle-layer weights, the learnable set (i.e., the set of ground-truth functions for which the above upper bound holds) of the 3-layer NTK without bias contains all finite degree polynomials, which is strictly larger than that of the 2-layer NTK without bias and is at least as large as the 2-layer NTK with bias. Recently, Geifman et al. (2020); Chen & Xu (2020) show that when all layers are trained, 3-layer NTK leads to exactly the same reproducing kernel Hilbert space (RKHS) as 2-layer NTK with biased ReLU (although they assumed an infinite number of neurons, and did not characterize the descent behavior of the generalization error). Combining with their results, we can draw the conclusion that training only the middle-layer weights is at least as effective as training all layers in 3-layer NTK, in terms of the size of the learnable set.

Sensitivity to the Choices of Bias: Even though a similar learnable set can be attained by 3-layer NTK (with or without bias) and 2-layer NTK (with bias), our results suggest that the actual generalization performance can still differ significantly in terms of the sensitivity to the choice of bias, especially when the input dimension d is large. One type of bias setting commonly used in literature (Ghorbani et al., 2021a; Satpathi & Srikant, 2021) is that the bias has a similar magnitude as each element of the input vector, which we refer to as “normal bias”. However, we show that such a normal bias setting has a negative impact on the generalization error for overfitted 2-layer NTK when d is large. To avoid this negative impact, it is important to use another type of bias setting where the bias has a similar magnitude as the norm of the whole input vector, which we refer to as

¹In this paper, the first hidden-layer denotes the one closer to the input layer, while the second hidden-layer denotes the one closer to the output layer.




Figure 1: A fully-connected three-layer neural network where input dimension $d = 2$, the number of neurons of the first hidden-layer $p_1 = 3$, and the number of neurons of the second hidden-layer $p_2 = 4$.

“balanced bias”. In contrast, for 3-layer NTK, different bias settings do not have an obvious effect on the generalization performance. In summary, compared with 2-layer NTK, the use of an extra non-linear layer in 3-layer NTK appears to significantly reduce the impact due to the choice of bias, and therefore makes the learning more robust.

Our work is related to the growing literature on the generalization performance of such shallow and fully-connected neural network. However, most of these studies focus on 2-layer neural networks. Among them, they differ in which layer to train. For example, [Mei & Montanari \(2019\)](#); [d’Ascoli et al. \(2020\)](#); [Mei et al. \(2022\)](#) consider the “random feature” (RF) model that only trains the top-layer weights and fixes the bottom-layer weights, while 2-layer NTK trains the bottom-layer weights. In contrast, our work on 3-layer NTK neither trains the bottom-layer or top-layer weights. Instead, we train the middle-layer weights, since the middle-layer of a 3-layer model involves the interaction between two hidden-layers, which does not exist in 2-layer models. The above studies of 2-layer network also differ in how the number of neurons/features p , the number of training samples n , and the input dimension d grow. [Mei & Montanari \(2019\)](#); [Mei et al. \(2022\)](#) study the generalization performance of the RF model where the number of neurons p , the number of training data n , and the input dimension grow proportionally to infinity. While [Ghorbani et al. \(2021b\)](#) focuses on the approximation error (i.e., expressiveness) of both RF and NTK models, their analysis on generalization error is only on the limit n or $p \rightarrow \infty$. All of these studies are quite different from ours with fixed n and finite p . Other works such as [Arora et al. \(2019\)](#); [Satpathi & Srikant \(2021\)](#); [Fiat et al. \(2019\)](#) study the situation where the number of training samples n is given and the number of neurons p is larger than a threshold, which is closer to our setup. However, these studies usually do not quantify how the generalization performance depends on the number of neurons p . Specifically, they usually provide an upper bound on the generalization error when the number of neurons p is greater than a threshold, while the upper bound itself does not depend on p . Thus, such an upper bound cannot explain the descent behavior of NTK models. The work in [Ju et al. \(2021\)](#) does study the descent behavior with respect to p , and is therefore the closest to our work. However, as we have explained earlier, there are crucial differences between 2 and 3 layers in both the descent behavior and the learnable set of ground-truth functions. In addition to the above references, our work is also related to [Allen-Zhu et al. \(2019\)](#) (which studies NTK without overfitting) and [Ji & Telgarsky \(2019\)](#) (which studies classification by NTK). Their settings are however different from ours in that we consider overfitted solutions for regression. In summary, our paper is the first to provide a high-probability upper bound on the generalization error of the overfitted 3-layer NTK (where only its middle layer weights are trained), and to characterize how the generalization error decreases with the number of neurons p_1 and p_2 .

2 System Model

Let $f : \mathbb{R}^d \mapsto \mathbb{R}$ denote the ground-truth function. Let $(\mathbf{X}_i, f(\mathbf{X}_i) + \epsilon_i), i = 1, 2, \dots, n$ denote n pieces of training data, where $\mathbf{X} \in \mathbb{R}^{n \times d}$ is the matrix, each column of which is the input of one training sample, $\epsilon \in \mathbb{R}^{n \times 1}$ denotes the noise in the output of training data. We define the training output vector generated by the ground-truth function as $\mathbf{F}(\mathbf{X}) := [f(\mathbf{X}_1) f(\mathbf{X}_2) \dots f(\mathbf{X}_n)]^T \in \mathbb{R}^n$.

We use $\mathbf{a}[j]$ to denote the j -th part (sub-vector) of the vector \mathbf{a} . The part size depends on \mathbf{a} . Specifically, If \mathbf{a} has p_1 elements, then each part has 1 elements. If \mathbf{a} has dp_1 elements, then each part has d elements. If \mathbf{a} has p_1p_2 elements, then each part has p_1 elements.

We consider a fully-connected 3-layer neural network as illustrated in Fig. 1, which consists of normalized d -dimensional input $\mathbf{x} \in \mathcal{S}^{d-1}$ (a unit hyper-sphere), p_1 ReLUs (rectifier linear units $\max(\cdot, 0)$) at the first hidden-layer, p_2 ReLUs at the second hidden-layer, bottom-layer weights (between input and 1st hidden-layer) $\mathbf{V} \in \mathbb{R}^{(p_1d) \times 1}$, middle-layer weights (between 1st hidden-layer and 2nd hidden-layer) $\mathbf{W}_0 \in \mathbb{R}^{(p_1p_2) \times 1}$, and top-layer weights $w \in \mathbb{R}^{p_2 \times 1}$ (between 2nd hidden-layer and output).

2.1 Overfitted NTK solution

In this subsection, we will derive the overfitted solution for this 3-layer neural network using the NTK approximation. Let $\mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}} \in \mathbb{R}^{p_1 \times 1}$ denote the output of the first hidden-layer. We then have

$$\mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}}[j] := (\mathbf{x}^T \mathbf{V}[j]) \mathbf{1}_{\{\mathbf{x}^T \mathbf{V}[j] > 0\}}, j = 1, 2, \dots, p_1. \quad (1)$$

(We use the superscript ‘‘RF’’ because $\mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}}$ is indeed the feature vector of a random feature model (Mei & Montanari, 2019).) After training the middle-layer weights, \mathbf{W}_0 changes to $\mathbf{W}_1 := \mathbf{W}_0 + \overline{\Delta \mathbf{W}}$. Then, the change of the output is

$$\sum_{k=1}^{p_2} w_k \mathbf{1}_{\{\mathbf{W}_1[k]^T \mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}} > 0\}} \mathbf{W}_1[k]^T \mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}} - \sum_{k=1}^{p_2} w_k \mathbf{1}_{\{\mathbf{W}_0[k]^T \mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}} > 0\}} \mathbf{W}_0[k]^T \mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}}.$$

The NTK model (Jacot et al., 2018) assumes that $\overline{\Delta \mathbf{W}}$ is very small and thus the activation pattern does not change much. In other words, we can approximate $\mathbf{1}_{\{\mathbf{W}_1[k]^T \mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}} > 0\}}$ by $\mathbf{1}_{\{\mathbf{W}_0[k]^T \mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}} > 0\}}$. Define $\Delta \mathbf{W} \in \mathbb{R}^{(p_1p_2) \times 1}$ as $\Delta \mathbf{W}[k] := w_k \cdot \overline{\Delta \mathbf{W}}[k]$, $k = 1, 2, \dots, p_2$. Define $\mathbf{h}_{\mathbf{V},\mathbf{W}_0,\mathbf{x}}^{\text{Three}} \in \mathbb{R}^{1 \times (p_1p_2)}$ such that

$$\mathbf{h}_{\mathbf{V},\mathbf{W}_0,\mathbf{x}}^{\text{Three}}[k] := (\mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}})^T \cdot \mathbf{1}_{\{(\mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}})^T \mathbf{W}_0[k] > 0\}}, \quad (2)$$

where $k = 1, 2, \dots, p_2$. Therefore, the change of the output can be approximated by

$$\sum_{k=1}^{p_2} w_k \mathbf{1}_{\{\mathbf{W}_0[k]^T \mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}} > 0\}} \overline{\Delta \mathbf{W}}[k]^T \mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}} = \mathbf{h}_{\mathbf{V},\mathbf{W}_0,\mathbf{x}}^{\text{Three}} \Delta \mathbf{W}.$$

We thus obtain a linear model in $\Delta \mathbf{W}$. We provide an illustration of the formation and structure of these vectors in Fig. 4, Appendix A.1 in Supplementary Material. Define the design matrix $\mathbf{H} \in \mathbb{R}^{n \times (p_1p_2)}$ such that its i -th row is $\mathbf{H}_i = \mathbf{h}_{\mathbf{V},\mathbf{W}_0,\mathbf{x}_i}^{\text{Three}}$. Notice that overfitted gradient descent on a linear model converges to the min ℓ_2 -norm solution², which is denoted by

$$\Delta \mathbf{W}^{\ell_2} := \arg \min_{\mathbf{w} \in \mathbb{R}^{(p_1p_2) \times 1}} \|\mathbf{w}\|_2 \text{ subject to } \mathbf{H}\mathbf{w} = \mathbf{F}(\mathbf{X}) + \epsilon.$$

When \mathbf{H} is full row-rank (which holds with high probability under certain conditions), the trained model is then

$$\hat{f}^{\ell_2}(\mathbf{x}) = \mathbf{h}_{\mathbf{V},\mathbf{W}_0,\mathbf{x}}^{\text{Three}} \Delta \mathbf{W}^{\ell_2} = \mathbf{h}_{\mathbf{V},\mathbf{W}_0,\mathbf{x}}^{\text{Three}} \mathbf{H}^T (\mathbf{H}\mathbf{H}^T)^{-1} (\mathbf{F}(\mathbf{X}) + \epsilon). \quad (3)$$

Notice that the trained model is determined by multiple random variables.

In order to analyze the generalization performance of the trained model, we have to make assumptions on the distribution of those random variables. Let $\mu(\cdot)$, $\lambda(\cdot)$, and $\gamma(\cdot)$ denote the probability density function of \mathbf{x} , $\mathbf{V}[j]$, and $\mathbf{W}_0[k]$, respectively. For simplicity, we make the following assumption that all random variables follow uniform distribution.

²As suggested by other prior works Satpathi & Srikant (2021); Hastie et al. (2019); Ju et al. (2021), if we perform gradient descent training on a linear model from zero initial point until the training error is zero (i.e., overfitting), then the solution will be exactly the min ℓ_2 -norm solution. Note that we do not need to be concerned about the training dynamics here, because the min ℓ_2 -norm overfitted solution can be written down exactly as in Eq. (3).

Assumption 1. *The input \mathbf{x} and the bottom-layer initial weights $\mathbf{V}[j]$'s ($j = 1, 2, \dots, p_1$) are i.i.d. and uniformly distributed in \mathcal{S}^{d-1} . In other words, $\mu(\cdot)$ and $\lambda(\cdot)$ are both $\text{unif}(\mathcal{S}^{d-1})$. The middle-layer initial weights $\mathbf{W}_0[k]$'s ($k = 1, 2, \dots, p_2$) are i.i.d. and uniformly distributed in \mathcal{S}^{p_1-1} . In other words, $\gamma(\cdot)$ is $\text{unif}(\mathcal{S}^{p_1-1})$. The top-layer weights w are all non-zero³.*

Remark 1. Readers may be curious why we only train the middle-layer weights. Part of the reason is technicality: if the bottom layer is also trained, the aggregate output of the first hidden-layer may have changed so much that the second hidden-layer's inputs and ReLU activation patterns change significantly from initialization, which may violate the NTK assumption. The work in Geifman et al. (2020); Chen & Xu (2020) is not concerned about this difficulty, since they are mostly interested in the expressive power of the RKHS, assuming an infinite number of neurons. In contrast, we wish to capture the effect of finite width, and thus train only the middle layer to avoid this difficulty. More importantly, the middle-layer weights interact with both the first hidden layer and the second hidden-layer, and are the major structural distinction compared with 2-layer NTK. This setting thus helps us to answer the following interesting question: will training the middle layer alone already achieve the same (potential) benefit as training all layers (especially given that the latter encounters more technical difficulty)?

3 Generalization Performance

In this section, we will show our main results about the generalization performance of the aforementioned 3-layer NTK model for a specific set of functions. We first introduce a set of ground-truth functions that may be learnable and then provide a high-probability upper bound on the test error. We then discuss some useful implications of our upper bound.

3.1 A set of ground-truth functions that may be learnable

We define kernel functions K^{RF} , K^{Two} , and $K^{\text{Three}} : [-1, 1] \mapsto \mathbb{R}$ as follows (whose meanings will be explained soon):

$$K^{\text{RF}}(a) := \frac{\sqrt{1-a^2} + a \cdot (\pi - \arccos(a))}{2d\pi}, \quad (4)$$

$$K^{\text{Two}}(a) := a \cdot \frac{\pi - \arccos(a)}{2\pi}, \quad (5)$$

$$K^{\text{Three}}(a) := \frac{K^{\text{Two}}(2d \cdot K^{\text{RF}}(a))}{2d}. \quad (6)$$

(Notice that $2d \cdot K^{\text{RF}}(a) \in [0, 1]$ for all $a \in [-1, 1]$ by Lemma 43 in Supplementary Material, Appendix I, and hence $K^{\text{Three}}(\cdot)$ is well defined.) We define a set $\mathcal{F}_{(3)}^{\ell_2}$ of ground-truth functions based on those kernels:

Definition 1 (learnable set of 3-layer NTK).

$$\mathcal{F}_{(3)}^{\ell_2} := \left\{ f_g : \mathcal{S}^{d-1} \mapsto \mathbb{R} \mid f_g(\mathbf{x}) = \int_{\mathcal{S}^{d-1}} K^{\text{Three}}(\mathbf{x}^T \mathbf{z}) g(\mathbf{z}) d\mu(\mathbf{z}), \|g\|_\infty < \infty \right\}, \quad (7)$$

where $\|g\|_\infty := \sup_{\mathbf{z} \in \mathcal{S}^{d-1}} |g(\mathbf{z})|$.

To see why functions in $\mathcal{F}_{(3)}^{\ell_2}$ may be learnable, we can check what the learned result \hat{f}^{ℓ_2} in Eq. (3) should look like. When there are infinite number of neurons and there is no noise (i.e., $\epsilon = \mathbf{0}$), what remains on the right-hand-side of Eq. (3) can be viewed as the product of two terms, $\mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}} \mathbf{H}^T$ and $(\mathbf{H}\mathbf{H}^T)^{-1} \mathbf{F}(\mathbf{X})$. For the first term $\mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}} \mathbf{H}^T$, note that each row of \mathbf{H} is given by $\mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}_i}^{\text{Three}}$ for $i = 1, 2, \dots, n$. Thus, when $p_1, p_2 \rightarrow \infty$, the i -th element of $\mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}} \mathbf{H}^T$, which is the inner product between $\mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}}$ and $\mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}_i}^{\text{Three}}$, converges in probability to $K^{\text{Three}}(\mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}} (\mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}_i}^{\text{Three}})^T)$, which is exactly the kernel function of 3-layer NTK. By representing the second term $(\mathbf{H}\mathbf{H}^T)^{-1} \mathbf{F}(\mathbf{X})$ with a certain $g(\cdot)$, \hat{f}^{ℓ_2} must then approach the form

³We do not need to specify the distribution of w , since w is absorbed into the regressor $\Delta \mathbf{W}$ by definition $\Delta \mathbf{W}[k] := w_k \cdot \overline{\Delta \mathbf{W}}[k]$, $k = 1, 2, \dots, p_2$.

in Eq. (7). (See Supplementary Material, Appendix B for details.) Intuitively, K^{Three} can be thought of as the composition of the kernels of each of the two layers, which are K^{Two} and K^{RF} given in Eq. (5) and Eq. (4). Specifically, suppose that we fix the output of the first hidden layer (i.e., $\mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}}$) and regard it as the input of a 2-layer NTK formed by the top two layers of the 3-layer neural network. By letting $p_2 \rightarrow \infty$, we can show that the inner product between $\mathbf{h}_{\mathbf{V},\mathbf{W}_0,\mathbf{x}}^{\text{Three}}$ and $\mathbf{h}_{\mathbf{V},\mathbf{W}_0,\mathbf{X}_i}^{\text{Three}}$ approaches $K^{\text{Two}}((\mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V},\mathbf{X}_i}^{\text{RF}})$ (with necessary normalization of $\mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}}$ and $\mathbf{h}_{\mathbf{V},\mathbf{X}_i}^{\text{RF}}$), where K^{Two} is exactly the kernel of 2-layer NTK in Ju et al. (2021). Second, when $p_1 \rightarrow \infty$, we can show that $(\mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V},\mathbf{X}_i}^{\text{RF}}$ approaches $K^{\text{RF}}(\mathbf{x}^T \mathbf{X}_i)$, where K^{RF} is exactly the kernel of the random-feature model (Mei & Montanari, 2019). In summary, we expect that functions in $\mathcal{F}_{(3)}^{\ell_2}$ can be approximated by $\hat{f}^{\ell_2}(\cdot)$. However, we note that the above deviation is only about the expressiveness of 3-layer NTK and it does not precisely reveal its generalization performance.

3.2 An upper bound on the generalization error

We now present the first main result of this paper, which is an upper bound that quantifies the relationship between the generalization performance and system parameters.

Theorem 1. *For any ground-truth function $f(\mathbf{x}) = f_g(\mathbf{x}) \in \mathcal{F}_{(3)}^{\ell_2}$, when d is fixed and p_1, p_2 are much larger than n , (with high probability) we have*

$$|\hat{f}^{\ell_2}(\mathbf{x}) - f(\mathbf{x})| = \underbrace{O\left(\frac{\|g\|_{\infty}}{\sqrt{n}}\right)}_{\text{Term A}} + \underbrace{\left(O\left(\frac{\|g\|_1}{\sqrt{p_2}}\right)\right)}_{\text{Term B}} + \underbrace{\left(O\left(\|g\|_1 \sqrt[4]{\frac{\log p_1}{p_1}}\right) + \frac{\|\epsilon\|_2}{\sqrt{n}}\right)}_{\text{Term C}} \cdot \underbrace{O\left(n^{\frac{2}{d-1} + \frac{1}{2}} \cdot \sqrt{\log n}\right)}_{\text{Term E}}. \quad (8)$$

A more precise version of the upper bound and the condition of Theorem 1 as well as its derivation can be found in Supplementary Material, Appendix C.

As we can see, Eq. (8) captures how the test error depends on finite values of parameters $n, p_1, p_2, \|\epsilon\|_2$, and g . Later in this section we will examine more closely how n, p_1 , and p_2 affect the value of the upper bound. Regarding the dependency on g , Eq. (8) works as long as $\|g\|_1$ and $\|g\|_{\infty}$ are finite⁴. Intuitively, the norm of g represents the complexity of the ground-truth function in $\mathcal{F}_{(3)}^{\ell_2}$. When the norm of g is larger, then the right-hand side of Eq. (8) becomes larger, which indicates that such ground-truth function is harder to learn. A simple example is that if we enlarge a ground truth function $f_g \in \mathcal{F}_{(3)}^{\ell_2}$ by 2 times (which means g is 2 times larger), then since the model is linear, the test error $|\hat{f}^{\ell_2}(\mathbf{x}) - f(\mathbf{x})|$ will become 2 times larger. We will discuss more about which types of functions satisfy the condition of finite norm of g in Section 4.

Next, we will discuss some implications of this upper bound of 3-layer NTK. While some of them are similar to 2-layer NTK, others are significantly different, revealing the complexity due to having more layers.

3.3 Interpretations similar to 2-layer NTK

Based on the upper bound in Theorem 3, we have the following insights for 3-layer NTK, which are similar to those for 2-layer NTK shown in Ju et al. (2021). These similarities may reveal some intrinsic properties of the NTK models regardless of the number of layers.

Zero test error with $n \rightarrow \infty$ in the ideal situation: In the ideal situation where there are infinitely many neurons and no noise, the only remaining term in Eq. (8) is Term A. Notice that Term A decreases to zero as $n \rightarrow \infty$, which indicates that the generalization error decreases to zero when

⁴Indeed, as long as $\|g\|_{\infty} < \infty$, then $\|g\|_1 < \infty$. That is why we only include the condition $\|g\|_{\infty} < \infty$ in Eq. (7). Notice that the assumption $\|g\|_{\infty} < \infty$ can be relaxed to $\|g\|_1 < \infty$ by similar methods showing in Ju et al. (2021). However, as shown in Ju et al. (2021), such relaxation leads to a different upper bound with slower descent speed with respect to n .




Figure 2: Curves of MSE for 3-layer NTK (no-bias) with respect to p_1 or p_2 when there exists Gaussian noise whose mean is zero and the variance is σ^2 . The ground-truth function is $f(\mathbf{x}) = (\mathbf{x}^T \mathbf{e}_1)^2 + (\mathbf{x}^T \mathbf{e}_1)^3$ where $d = 3$. Sample size is $n = 200$. Every curve is the median of 20 random simulations.

more training data are provided in the ideal situation. Term A suggests that such decreasing speed is at least $1/\sqrt{n}$. Such result is consistent with that of 2-layer NTK, e.g., in [Arora et al. \(2019\)](#).

3.4 Insights that are distinct compared with 2-layer NTK

Compared with 2-layer NTK, an important difference for 3-layer NTK is that there are more than one hidden-layers. Therefore, the speed of the descent of 3-layer NTK involves the interaction between two hidden-layers.

Descent with respect to the number of neurons: In Eq. (8), Term B and Term C contain p_1 and p_2 , respectively. For any given n and noise level $\|\epsilon\|_2$, Terms A and D do not change, and Term E decreases with p_1 and p_2 . (More discussion about Term E can be found in Supplementary Material, Appendix D, where we discuss the noise effect.) Therefore, by increasing p_1 and p_2 , Term B and Term C keep decreasing. In summary, right-hand-side of Eq. (8) decreases as the number of neurons p_1 and p_2 increases, which validates the descent in the overparameterized region of 3-layer NTK.

Different descent speed: As shown in Eq. (8), p_1 and p_2 play different roles in the descent of the generalization error. Comparing Term B and Term C of Eq. (8), we can see that the upper bound of the test error $|\hat{f}^{\ell_2}(\mathbf{x}) - f(\mathbf{x})|$ decreases faster with respect to p_2 (at the speed of $\sqrt{p_2}$) and slower with respect to p_1 (at the speed of $\sqrt[4]{p_1/\log p_1}$). We emphasize that this difference is not due to the number of weights/parameters contributed by the number of neurons in each hidden-layer of p_1 and p_2 ⁵. Instead, we conjecture that such difference in the speed of descent may be due to the different positions in this 3-layer neural network structure, where the second hidden-layer takes the trained middle-layer weights as its input (and thus utilizes the trained weights better than the first hidden-layer).

We use numerical results to illustrate the different roles of p_1 and p_2 in reducing the generalization error. We fix $p_2 = 200$ and plot the MSE with respect to p_1 in Fig. 2(b). Although the test error decreases when p_1 increases, the decreasing speed is slow, especially for the noisy situation. Such a slow decreasing speed with p_1 remains even when p_2 is fixed to a much higher value. For example, in Fig. 2(b), we fix $p_2 = \infty$, we still observe the similarly slow decreasing speed with p_1 as shown by Fig. 2(c). In contrast, the descent with respect to p_2 should be easier to observe and can reach a lower test MSE. In Fig. 2(d), we fix $p_1 = 200$ and increase p_2 (i.e., we exchange the values of p_1 and p_2 in Fig. 2(c)(d)). As we can see, all three curves in Fig. 2(d) have a more obvious descent and decrease to lower MSE compared with those in Fig. 2(c), which validates our conjecture that the descent speed with respect to the number of neurons of the second hidden-layer is faster.

Notice that our upper bound Eq. (8) also suggests a descent when both p_1 and p_2 increase simultaneously. We use simulation result by Fig. 2(a) to support this point. We fixed $n = 200$ and let $p_1 = p_2$ increase simultaneously. The ground-truth model in this figure is $f(\mathbf{x}) = (\mathbf{x}^T \mathbf{e})^2 + (\mathbf{x}^T \mathbf{e})^3$ where $d = 3$. The green, orange, and blue curves denote the situations of $\sigma^2 = 0$ (no noise), $\sigma^2 = 0.01$, and $\sigma^2 = 0.04$, respectively. Every point in this figure is the median of 20 simulation runs. We also

⁵Specifically, the number of weights that get trained equals to $p_1 p_2$ and the total number of weights for bottom, middle, and top layers equals to $d p_1 + p_1 p_2 + p_2$. In other words, the number of weights (either for the middle layer or for all layers) does not increase faster by increasing p_2 instead of p_1 .

provide the box plot⁶ of the situation of $\sigma^2 = 0.01$ (correspond to the orange curve). It is obvious that all three curves descend, which verifies that the generalization error of the overfitted 3-layer NTK model decreases when p_1 and p_2 increase simultaneously at the same speed. By observing the box plot for the situation $\sigma^2 = 0.01$ (the orange curve), we also notice that when $p_1 = p_2$ becomes large, the variance becomes small. This is because all initial weights are *i.i.d.* random and a large number of weights may reduce the variance of the model due to the law of large numbers. Our upper bound in Theorem 3 also suggests such reduced variance as the probability in Theorem 3 increases as p_1 increases.

4 Types of Ground-Truth Functions

Are 3-layer (i.e., deeper) networks better than 2-layer networks in any way for generalization performance? In the last section, we have seen that both 3-layer NTK and 2-layer NTK can achieve zero test error when $n \rightarrow \infty$ in the ideal noiseless situation, when the ground-truth functions are in their respective learnable set⁷. A natural question is then to compare the learnable sets between these two models, and to compare the generalization performance when the ground-truth function belongs to both learnable sets. In this section, we provide some answers by studying various types of ground-truth functions and their effects on the generalization performance.

4.1 Size of the learnable set

For a 2-layer NTK, as shown in Ju et al. (2021), when no bias is used in ReLU, the corresponding learnable set $\mathcal{F}_{(2)}^{\ell_2}$ contains all even polynomials and linear functions, but does not contain other odd polynomials. In order to learn both even and odd polynomials, it is critical that bias is added to ReLU (Satpathi & Srikant, 2021; Ju et al., 2021). In contrast, we prove the following result:

Proposition 2. $\mathcal{F}_{(3)}^{\ell_2}$ (with unbiased ReLU, middle layer being trained) already contains all polynomials with finite degree (i.e., including both even and odd polynomials). Further, the learnable set $\mathcal{F}_{(3)}^{\ell_2}$ of 3-layer NTK is strictly larger than that of the 2-layer NTK with unbiased ReLU, and is at least as large as that of the 2-layer NTK with biased ReLU.

This independence to bias shown by Proposition 2 can be seen as one performance advantage of 3-layer NTK compared to 2-layer NTK. Details (including more precise statement) about this result is in Supplementary Material, Appendix J. Notice that Geifman et al. (2020); Chen & Xu (2020) show that when training all layers, 3-layer NTK leads to the same RKHS as 2-layer NTK with biased ReLU. However, it is unclear whether training one layer is already sufficient for achieving the same RKHS as training all layers. Our result in Proposition 2 answers this question positively, i.e., only training the middle layer has already achieved all benefits of training all layers in terms of the size of the learnable set. (In other words, training all three layers will not expand the learnable set over training only the middle layer.)

4.2 Different bias settings with high input dimension

Even when a ground-truth function belongs to both $\mathcal{F}_{(2),b}^{\ell_2}$ and $\mathcal{F}_{(3)}^{\ell_2}$, their generalization performance may still exhibit some differences. In this subsection, we will show that when the input dimension d is high, some specific choice of bias of the 2-layer NTK has better generalization performance than others. In contrast, the 3-layer NTK is less sensitive to different bias settings.

Notice that adding bias to each ReLU in 2-layer NTK is equivalent to appending a constant to \mathbf{x} while still using ReLU without bias. Specifically, the input vector for biased 2-layer NTK is

$$\mathbf{x}_b := \left[\sqrt{1-b^2} \cdot \mathbf{x} \right] \in \mathbb{R}^{d+1}, \quad (9)$$

⁶From bottom to top, the five horizontal lines of each marker of a box plot represent the minimum (excluding outliers), first quartile (25%), median (50%), third quartile (75%), and maximum (excluding outliers), respectively. See (McGill et al., 1978) for more details.

⁷We illustrate the generalization performance of ground-truth functions outside the learnable set in Supplementary Material, Appendix J.3.

Model	Learnable functions set	Category
3-layer NTK, no-bias	$\mathcal{F}_{(3)}^{\ell_2} = \left\{ \int_{\mathcal{S}^{d-1}} K^{\text{Three}}(\mathbf{x}^T \mathbf{z}) g(\mathbf{z}) d\mu(\mathbf{z}) \right\}$	(i)
2-layer NTK, no-bias	$\mathcal{F}_{(2)}^{\ell_2} = \left\{ \int_{\mathcal{S}^{d-1}} K^{\text{Two}}(\mathbf{x}^T \mathbf{z}) g(\mathbf{z}) d\mu(\mathbf{z}) \right\}$	(ii)
2-layer NTK, normal-bias	$\mathcal{F}_{(2),\text{NLB}}^{\ell_2} = \left\{ \int_{\mathcal{S}^{d-1}} K^{\text{Two}}\left(\frac{d}{d+1} \mathbf{x}^T \mathbf{z} + \frac{1}{d+1}\right) g(\mathbf{z}) d\mu(\mathbf{z}) \right\}$	(i)
2-layer NTK, balanced-bias	$\mathcal{F}_{(2),\text{BB}}^{\ell_2} = \left\{ \int_{\mathcal{S}^{d-1}} K^{\text{Two}}\left(\frac{1}{2} \mathbf{x}^T \mathbf{z} + \frac{1}{2}\right) g(\mathbf{z}) d\mu(\mathbf{z}) \right\}$	(i)

Table 1: Learnable functions for different NTK models. Category: (i) can learn both even- and odd-power polynomials; (ii) cannot learn other odd-power polynomials except linear functions. (We omit the condition $\|g\|_\infty < \infty$ in the expression of learnable sets to save space.)




Figure 3: Comparison of test MSE with respect to n between different NTK models when the number of neurons is infinite and without noise. The ground-truth function is $f(\mathbf{x}) = \frac{d+2}{3} (\mathbf{x}^T \mathbf{e}_1)^3 - \mathbf{x}^T \mathbf{e}_1$. Every curve is the average of 10 random simulations.

where $b \in (0, 1)$ denotes the initial bias. We also normalize the first d elements of \mathbf{x}_b by $\sqrt{1-b^2}$ in Eq. (9) to make sure that $\|\mathbf{x}_b\|_2 = 1$. Under this biased setting, the 2-layer NTK model has the learnable set $\mathcal{F}_{(2),b}^{\ell_2} := \left\{ \int_{\mathcal{S}^{d-1}} K^{\text{Two}}((1-b^2)\mathbf{x}^T \mathbf{z} + b^2) g(\mathbf{z}) d\mu(\mathbf{z}), \|g\|_\infty < \infty \right\}$.

A common setup for the initial magnitude of the bias of each ReLU is to use a value that is close or equal to the average magnitude of each element of input \mathbf{x} , e.g., [Satpathi & Srikant \(2021\)](#); [Ghorbani et al. \(2021a\)](#). Specifically, we let $b = \frac{1}{\sqrt{d+1}}$ in Eq. (9), and denote the corresponding learnable set by $\mathcal{F}_{(2),\text{NLB}}^{\ell_2}$. We refer to this setting as the “normal-bias” setting. Alternatively, the initial magnitude of the bias can be chosen to be close or equal to $\|\mathbf{x}\|_2$. Specifically, we let $b = \frac{1}{\sqrt{2}}$ in Eq. (9) and denote the corresponding learnable set by $\mathcal{F}_{(2),\text{BB}}^{\ell_2}$. We refer to this second setting as “balanced-bias”. The specific expression of $\mathcal{F}_{(2)}^{\ell_2}$, $\mathcal{F}_{(2),\text{NLB}}^{\ell_2}$, and $\mathcal{F}_{(2),\text{BB}}^{\ell_2}$ can be derived by using similar methods shown in Section 3.1 (results are listed in Table 1).

We now discuss how the two different bias settings could affect the generalization performance when d is large. For 2-layer NTK under the normal-bias setting, the kernel is $K^{\text{Two}}\left(\frac{d}{d+1} \mathbf{x}^T \mathbf{z} + \frac{1}{d+1}\right)$. Although it contains both even and odd power polynomials, we notice that when d increases, K^{Two} approaches its no-bias counterpart $K^{\text{Two}}(\mathbf{x}^T \mathbf{z})$, which only contains even power polynomials and linear term. Thus, we conjecture that, by increasing d , the generalization performance of 2-layer NTK with normal-bias will deteriorate for those ground-truth functions inside $\mathcal{F}_{(2),\text{NLB}}^{\ell_2}$ but far away from $\mathcal{F}_{(2)}^{\ell_2}$ (e.g., odd-degree non-linear polynomials). In contrast, for 2-layer NTK under the balanced-bias setting, the kernel is $K^{\text{Two}}\left(\frac{1}{2} \mathbf{x}^T \mathbf{z} + \frac{1}{2}\right)$, which does not change with d . Therefore, we expect that such deterioration should not happen. Note that in 3-layer NTK, although normal-bias setting still approaches no-bias setting when d increases, there does not exist such performance deterioration, because $\mathcal{F}_{(3)}^{\ell_2}$ (the learnable set of 3-layer NTK without bias) already contains both even and odd power polynomials. These insights will be verified by the numerical results below.

We now use simulation results in Fig. 3 to validate the conjecture that 3-layer NTK models are less sensitive to different bias settings than 2-layer NTK models. We let the ground-truth function be $f(\mathbf{x}) = \frac{d+2}{3} (\mathbf{x}^T \mathbf{e}_1)^3 - \mathbf{x}^T \mathbf{e}_1$, which is orthogonal to $\mathcal{F}_{(2)}^{\ell_2}$. In Fig. 3(a) when $d = 2$, all settings have similar performance except 2-layer NTK without bias, whose test error is always above the null risk. In Fig. 3(b) when $d = 15$, the purple curve of 2-layer NTK with normal bias gets closer to the

red curve of 2-layer NTK without bias (and thus the generalization performance becomes worse), while other curves are still close to each other. This validates our conjecture that 3-layer NTK models are less sensitive to different bias settings than 2-layer NTK models. Further simulations can be found in Appendix A.2.

5 Conclusion

In this paper, we studied the generalization performance of overfitted 3-layer NTK models. Compared with 2-layer NTK models, 3-layer NTK is less sensitive to different bias settings. Further, training only the middle layer can get most of the performance advantage of 3-layer NTK, in terms of the learnable set. Possible future directions include: (i) studying whether training other layers will get the same benefit as training the middle layer; (ii) approximating the actual neural network where the learned result is far away from the initial state; (iii) investigating deeper network as well as other structures such as convolutional neural network (CNN) and recursive neural network (RNN).

Acknowledgments and Disclosure of Funding

This work has been supported in part by NSF grants: 2112471 (also partly funded by DHS), CNS-2106933, CNS-1901057, CNS-2113893, and the Bilisland Dissertation Fellowship at Purdue University, and a grant from the Army Research Office: W911NF-21-1-0244.

References

- Advani, M. S., Saxe, A. M., and Sompolinsky, H. High-dimensional dynamics of generalization error in neural networks. *Neural Networks*, 132:428–446, 2020.
- Allen-Zhu, Z., Li, Y., and Liang, Y. Learning and generalization in overparameterized neural networks, going beyond two layers. In *Advances in neural information processing systems*, pp. 6158–6169, 2019.
- Arora, S., Du, S., Hu, W., Li, Z., and Wang, R. Fine-grained analysis of optimization and generalization for overparameterized two-layer neural networks. In *International Conference on Machine Learning*, pp. 322–332, 2019.
- Bartlett, P. L., Long, P. M., Lugosi, G., and Tsigler, A. Benign overfitting in linear regression. *Proceedings of the National Academy of Sciences*, 2020.
- Belkin, M., Ma, S., and Mandal, S. To understand deep learning we need to understand kernel learning. In *International Conference on Machine Learning*, pp. 541–549, 2018.
- Belkin, M., Hsu, D., and Xu, J. Two models of double descent for weak features. *arXiv preprint arXiv:1903.07571*, 2019.
- Bell, H. E. Gershgorin’s theorem and the zeros of polynomials. *The American Mathematical Monthly*, 72(3):292–295, 1965.
- Bishop, C. M. *Pattern recognition and machine learning*. Springer, 2006.
- Chaudhry, M. A., Qadir, A., Rafique, M., and Zubair, S. Extension of euler’s beta function. *Journal of computational and applied mathematics*, 78(1):19–32, 1997.
- Chen, C.-P. and Qi, F. The best bounds in wallis’ inequality. *Proceedings of the American Mathematical Society*, pp. 397–401, 2005.
- Chen, L. and Xu, S. Deep neural tangent kernel and laplace kernel have the same rkhs. *arXiv preprint arXiv:2009.10683*, 2020.
- Dokmanic, I. and Petrinovic, D. Convolution on the n -sphere with application to pdf modeling. *IEEE transactions on signal processing*, 58(3):1157–1170, 2009.
- Dutka, J. The incomplete beta function—a historical profile. *Archive for history of exact sciences*, pp. 11–29, 1981.

- d'Ascoli, S., Refinetti, M., Biroli, G., and Krzakala, F. Double trouble in double descent: Bias and variance (s) in the lazy regime. In *International Conference on Machine Learning*, pp. 2280–2290. PMLR, 2020.
- Fiat, J., Malach, E., and Shalev-Shwartz, S. Decoupling gating from linearity. *arXiv preprint arXiv:1906.05032*, 2019.
- Geifman, A., Yadav, A., Kasten, Y., Galun, M., Jacobs, D., and Ronen, B. On the similarity between the laplace and neural tangent kernels. *Advances in Neural Information Processing Systems*, 33: 1451–1461, 2020.
- Ghorbani, B., Mei, S., Misiakiewicz, T., and Montanari, A. Linearized two-layers neural networks in high dimension. *The Annals of Statistics*, 49(2):1029–1054, 2021a.
- Ghorbani, B., Mei, S., Misiakiewicz, T., and Montanari, A. Linearized two-layers neural networks in high dimension. *The Annals of Statistics*, 49(2):1029–1054, 2021b.
- Hastie, T., Tibshirani, R., and Friedman, J. *The elements of statistical learning: data mining, inference, and prediction*. Springer Science & Business Media, 2009.
- Hastie, T., Montanari, A., Rosset, S., and Tibshirani, R. J. Surprises in high-dimensional ridgeless least squares interpolation. *arXiv preprint arXiv:1903.08560*, 2019.
- Jacot, A., Gabriel, F., and Hongler, C. Neural tangent kernel: Convergence and generalization in neural networks. In *Advances in neural information processing systems*, pp. 8571–8580, 2018.
- James, W. and Stein, C. Estimation with quadratic loss. In *Breakthroughs in Statistics*, pp. 443–460. Springer, 1992.
- Ji, Z. and Telgarsky, M. Polylogarithmic width suffices for gradient descent to achieve arbitrarily small test error with shallow relu networks. *arXiv preprint arXiv:1909.12292*, 2019.
- Ju, P., Lin, X., and Liu, J. Overfitting can be harmless for basis pursuit, but only to a degree. *Advances in Neural Information Processing Systems*, 33, 2020.
- Ju, P., Lin, X., and Shroff, N. On the generalization power of overfitted two-layer neural tangent kernel models. In *International Conference on Machine Learning*, pp. 5137–5147. PMLR, 2021.
- Laha, R. and Rohatgi, V. *Probability Theory*. Wiley Series in Probability and Statistics. Wiley, 1979. ISBN 9780471032625. URL <https://books.google.com/books?id=HBHvAAAAMAAJ>.
- LeCun, Y., Kanter, I., and Solla, S. A. Second order properties of error surfaces: Learning time and generalization. In *Advances in Neural Information Processing Systems*, pp. 918–924, 1991.
- Li, S. Concise formulas for the area and volume of a hyperspherical cap. *Asian Journal of Mathematics and Statistics*, 4(1):66–70, 2011.
- Li, Y. and Wong, R. Integral and series representations of the dirac delta function. *arXiv preprint arXiv:1303.1943*, 2013.
- McGill, R., Tukey, J. W., and Larsen, W. A. Variations of box plots. *The american statistician*, 32(1):12–16, 1978.
- Mei, S. and Montanari, A. The generalization error of random features regression: Precise asymptotics and double descent curve. *arXiv preprint arXiv:1908.05355*, 2019.
- Mei, S., Misiakiewicz, T., and Montanari, A. Generalization error of random feature and kernel methods: Hypercontractivity and kernel matrix concentration. *Applied and Computational Harmonic Analysis*, 59:3–84, 2022.
- Muthukumar, V., Vodrahalli, K., and Sahai, A. Harmless interpolation of noisy data in regression. In *2019 IEEE International Symposium on Information Theory (ISIT)*, pp. 2299–2303. IEEE, 2019.
- Satpathi, S. and Srikant, R. The dynamics of gradient descent for overparametrized neural networks. In *Learning for Dynamics and Control*, pp. 373–384. PMLR, 2021.

- Stein, C. Inadmissibility of the usual estimator for the mean of a multivariate normal distribution. Technical report, Stanford University Stanford United States, 1956.
- Tikhonov, A. N. On the stability of inverse problems. In *Dokl. Akad. Nauk SSSR*, volume 39, pp. 195–198, 1943.
- Vilenkin, N. Y. Special functions and the theory of group representations. providence: American mathematical society. *sftp*, 1968.
- Wainwright, M. Uniform laws of large numbers, 2015. https://www.stat.berkeley.edu/~mjwain/stat210b/Chap4_Uniform_Feb4_2015.pdf, Accessed: Feb. 7, 2021.
- Zhang, C., Bengio, S., Hardt, M., Recht, B., and Vinyals, O. Understanding deep learning requires rethinking generalization. In *5th International Conference on Learning Representations, ICLR 2017*, 2017.

Checklist

1. For all authors...
 - (a) Do the main claims made in the abstract and introduction accurately reflect the paper’s contributions and scope? [Yes]
 - (b) Did you describe the limitations of your work? [Yes] E.g., see Remark 1.
 - (c) Did you discuss any potential negative societal impacts of your work? [No] This work is on the learning theory and should have no potential negative societal impacts.
 - (d) Have you read the ethics review guidelines and ensured that your paper conforms to them? [Yes]
2. If you are including theoretical results...
 - (a) Did you state the full set of assumptions of all theoretical results? [Yes]
 - (b) Did you include complete proofs of all theoretical results? [Yes]
3. If you ran experiments...
 - (a) Did you include the code, data, and instructions needed to reproduce the main experimental results (either in the supplemental material or as a URL)? [No] As this is mainly a theoretical paper.
 - (b) Did you specify all the training details (e.g., data splits, hyperparameters, how they were chosen)? [Yes]
 - (c) Did you report error bars (e.g., with respect to the random seed after running experiments multiple times)? [Yes] See Fig. 2.
 - (d) Did you include the total amount of compute and the type of resources used (e.g., type of GPUs, internal cluster, or cloud provider)? [No] As this is mainly a theoretical paper.
4. If you are using existing assets (e.g., code, data, models) or curating/releasing new assets...
 - (a) If your work uses existing assets, did you cite the creators? [N/A]
 - (b) Did you mention the license of the assets? [N/A]
 - (c) Did you include any new assets either in the supplemental material or as a URL? [N/A]
 - (d) Did you discuss whether and how consent was obtained from people whose data you’re using/curating? [N/A]
 - (e) Did you discuss whether the data you are using/curating contains personally identifiable information or offensive content? [N/A]
5. If you used crowdsourcing or conducted research with human subjects...
 - (a) Did you include the full text of instructions given to participants and screenshots, if applicable? [N/A]
 - (b) Did you describe any potential participant risks, with links to Institutional Review Board (IRB) approvals, if applicable? [N/A]
 - (c) Did you include the estimated hourly wage paid to participants and the total amount spent on participant compensation? [N/A]




Figure 4: Formation and structure of $h_{\mathbf{V}_0, \mathbf{W}_0, \mathbf{x}}^{\text{Three}}$ and $h_{\mathbf{V}_0, \mathbf{x}}^{\text{RF}}$.

A Additional Figures

A.1 Formation of features

In Fig. 4, we illustrate the formation and the structure of the features shown in Section 2.

A.2 About the conjecture in Section 4.2

We provide some additional simulation results (in addition to Fig. 3) to validate our conjecture in Section 4.2 that 2-layer NTK is more sensitive to different bias settings, especially when d is large. Note that in Fig. 3, we only consider one type of ground-truth functions that contains only odd-power polynomials. Here, we also examine other types of ground-truth functions.

Similar to Fig. 3, in Fig. 5, we consider the ideal case where there are infinite number of neurons. We plot curves of MSE with respect to n when $p, p_1, p_2 \rightarrow \infty$. The simulation setup is similar to Fig. 3, but here we consider more types of ground-truth functions (whose exact forms are given in the caption of Fig. 5). In sub-figures (a)(b), Type A function corresponds to even-power polynomials. We can see that all curves are close to each other in both low-dimensional case ($d = 2$) and high-dimensional case ($d = 15$). This is because the 2-layer NTK without bias can learn even-power polynomials. In other words, in high-dimensional cases, although the performance of the normal-bias setting approaches that of the no-bias setting, it does not hurt the generalization performance because the no-bias setting can already learn the Type A function. Sub-figures (c)(d) are exactly the same as Fig. 3, which uses the Type B ground-truth function corresponds to odd-power polynomials. Sub-figures (e)(f) adopt the Type C ground-truth function that contains both odd-power and even-power polynomials. The generalization performance shown by sub-figures (e)(f) is between that in sub-figures (a)(b) and that in sub-figures (c)(d). This is expected because Type C functions can be viewed as a mix of Type A and Type B functions.

We also consider the situation of finite number of neurons. In Fig. 6, we fix the number of training data and let the x-axis be p (for 2-layer NTK) or p_1 (for 3-layer NTK with fixed $p_2 = 100$). The setup of Fig. 7 is similar to the setup of Fig. 6 except that for 3-layer NTK we fix $p_1 = 100$ and change p_2 . Both in Fig. 6 and Fig. 7, when d is large and the ground-truth function is Type B (i.e., sub-figure (d)), we can see that the curve of 2-layer NTK with normal-bias (the purple curve marked by ◀) is closer to the curve of 2-layer NTK without bias (the red curve marked by ▶). This validates our conjecture in Section 4.2 that 2-layer NTK is more sensitive to different bias settings, especially when d is large.




Figure 5: Curves of MSE with respect to n for 2-layer and 3-layer NTK models when $p, p_1, p_2 \rightarrow \infty$ and $\epsilon = \mathbf{0}$. Let $\mathbf{e}_1 = [1 \ 0 \ 0 \ \dots \ 0]^T \in \mathbb{R}^d$. Type A function is $f(\mathbf{x}) = (\mathbf{x}^T \mathbf{e}_1)^4 - (\mathbf{x}^T \mathbf{e}_1)^2$. Type B function is $f(\mathbf{x}) = \frac{d+2}{3} (\mathbf{x}^T \mathbf{e}_1)^3 - \mathbf{x}^T \mathbf{e}_1$. Type C function is $f(\mathbf{x}) = (\mathbf{x}^T \mathbf{e}_1)^2 + (\mathbf{x}^T \mathbf{e}_1)^3$. Every curve is the average of 10 random simulations.

B Derivation of the learnable set $\mathcal{F}_{(3)}^{\ell_2}$

For the derivation of the learnable set, we assume that the noise ϵ is zero in Eq. (3). We first rewrite Eq. (3) as the sum of terms contributed by each sample. Recall that $\mathbf{H}^T = [\mathbf{H}_1^T \ \dots \ \mathbf{H}_n^T] \in \mathbb{R}^{(p_1 p_2) \times n}$ where $\mathbf{H}_i \in \mathbb{R}^{1 \times (p_1 p_2)}$, $i = 1, 2, \dots, n$. Thus, we have $\mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}} \mathbf{H}^T = \sum_{i=1}^n \left(\mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}} \mathbf{H}_i^T \right) \mathbf{e}_i^T$ where $\mathbf{e}_i \in \mathbb{R}^n$ denotes the i -th standard basis (i.e., the i -th element is 1 while all other elements are 0). Thus, we have

$$\hat{f}^{\ell_2}(\mathbf{x}) = \mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}} \mathbf{H}^T (\mathbf{H} \mathbf{H}^T)^{-1} \mathbf{F}(\mathbf{X}) = \sum_{i=1}^n \left(\frac{1}{p_2} \mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}} \mathbf{H}_i^T \right) p_2 \mathbf{e}_i^T (\mathbf{H} \mathbf{H}^T)^{-1} \mathbf{F}(\mathbf{X}). \quad (10)$$

For any $\mathbf{a}, \mathbf{b} \in \mathbb{R}^{p_1}$, we define a set

$$\mathcal{C}_{\mathbf{a}, \mathbf{b}}^{\mathbf{W}_0} := \{k \in \{1, 2, \dots, p_2\} \mid \mathbf{a}^T \mathbf{W}_0[k] > 0, \mathbf{b}^T \mathbf{W}_0[k] > 0\}, \quad (11)$$

whose cardinality is given by

$$|\mathcal{C}_{\mathbf{a}, \mathbf{b}}^{\mathbf{W}_0}| = \sum_{k=1}^{p_2} \mathbf{1}_{\{\mathbf{a}^T \mathbf{W}_0[k] > 0, \mathbf{b}^T \mathbf{W}_0[k] > 0\}}.$$

Intuitively, $\mathcal{C}_{\mathbf{a}, \mathbf{b}}^{\mathbf{W}_0}$ denotes the indices of the ReLU in the second hidden-layer that are activated both when the output of the first layer is \mathbf{a} and when the output of the first layer is \mathbf{b} . Then, by Eq. (2),




Figure 6: Curves of MSE with respect to p (for 2-layer NTK) or p_1 (for 3-layer NTK where $p_2 = 100$). Other settings such as types of ground-truth functions and $\epsilon = 0$ are the same as those in Fig. 5. Every curve is the average of 10 random simulations.

we have

$$\frac{1}{p_2} \mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}} \mathbf{H}_i^T = (\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V}, \mathbf{X}_i}^{\text{RF}} \frac{\left| \mathcal{C}_{\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}, \mathbf{h}_{\mathbf{V}, \mathbf{X}_i}^{\text{RF}}}^{\mathbf{W}_0} \right|}{p_2}. \quad (12)$$

By Assumption 1, which gives the distribution of \mathbf{W}_0 , we can calculate the limiting value of Eq. (12) when there are an infinite number of neurons in the second hidden-layer. Specifically, since

$$\frac{\left| \mathcal{C}_{\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}, \mathbf{h}_{\mathbf{V}, \mathbf{X}_i}^{\text{RF}}}^{\mathbf{W}_0} \right|}{p_2} \xrightarrow{P} \frac{\pi - \arccos \left(\frac{(\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V}, \mathbf{X}_i}^{\text{RF}}}{\|\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}\|_2 \cdot \|\mathbf{h}_{\mathbf{V}, \mathbf{X}_i}^{\text{RF}}\|_2} \right)}{2\pi}, \text{ as } p_2 \rightarrow \infty, \quad (13)$$

where \xrightarrow{P} denotes convergence in probability, we have

$$\text{Eq. (12)} \xrightarrow{P} \|\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}\|_2 \cdot \|\mathbf{h}_{\mathbf{V}, \mathbf{X}_i}^{\text{RF}}\|_2 \cdot K^{\text{Two}} \left(\frac{(\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V}, \mathbf{X}_i}^{\text{RF}}}{\|\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}\|_2 \cdot \|\mathbf{h}_{\mathbf{V}, \mathbf{X}_i}^{\text{RF}}\|_2} \right), \text{ as } p_2 \rightarrow \infty.$$

Note that K^{Two} is known to be the kernel of 2-layer NTK (Ju et al., 2021). It is natural that K^{Two} appears here, since we can regard the output of the first hidden-layer as the input of a 2-layer network consisting of the top- and middle-layer of the 3-layer network.

To further simplify the above expression, it remains to calculate $(\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V}, \mathbf{X}_i}^{\text{RF}}$. Similar to the derivation above, when the first hidden-layer has an infinite number of neurons, we have

$$(\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V}, \mathbf{X}_i}^{\text{RF}} \xrightarrow{P} K^{\text{RF}}(\mathbf{x}^T \mathbf{X}_i), \text{ as } p_1 \rightarrow \infty. \quad (14)$$

(Eq. (13) and Eq. (14) can be derived from integration over a hyper-sphere, which is shown in Lemma 20 and Lemma 21 in Appendix F.6, respectively.) Note that K^{RF} is also the kernel of the




Figure 7: Curves of MSE with respect to p (for 2-layer NTK) or p_2 (for 3-layer NTK where $p_1 = 100$). Other settings such as types of ground-truth functions and $\epsilon = 0$ are the same as those in Fig. 5. Every curve is the average of 10 random simulations.

random-feature model (Mei & Montanari, 2019). It is natural that K^{RF} appears here since Eq. (14) represents the situation that the bottom-layer has infinite width, which also appears in a random feature model. Notice that $K^{\text{RF}}(\mathbf{X}_i^T \mathbf{X}_i) = K^{\text{RF}}(\mathbf{x}^T \mathbf{x}) = K^{\text{RF}}(1) = \frac{1}{2d}$. Thus, we have

$$\|\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}\|_2 \cdot \|\mathbf{h}_{\mathbf{V}, \mathbf{X}_i}^{\text{RF}}\|_2 \xrightarrow{P} \frac{1}{2d}, \text{ as } p_1 \rightarrow \infty. \quad (15)$$

Plugging Eq. (13)(14)(15) into Eq. (12) and recalling Eq. (6), we thus have

$$\frac{1}{p_2} \mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}} \mathbf{H}_i^T \xrightarrow{P} K^{\text{Three}}(\mathbf{x}^T \mathbf{X}_i), \text{ as } p_1, p_2 \rightarrow \infty.$$

If we let

$$g(\mathbf{z}) = \sum_{i=1}^n p_2 \mathbf{e}_i^T (\mathbf{H}\mathbf{H}^T)^{-1} \mathbf{F}(\mathbf{X}) \delta_{\mathbf{X}_i}(\mathbf{z}),$$

(where $\delta_{\mathbf{z}_0}(\mathbf{z})$ denotes a δ -function, i.e., it has zero value for all $\mathbf{z} \in \mathcal{S}^{d-1} \setminus \{\mathbf{z}_0\}$, but its L_1 -norm is $\|\delta_{\mathbf{z}_0}\|_1 := \int_{\mathcal{S}^{d-1}} \delta_{\mathbf{z}_0}(\mathbf{z}) d\mu(\mathbf{z}) = 1$), then as p_1 and $p_2 \rightarrow \infty$, Eq. (10) approaches $\int_{\mathcal{S}^{d-1}} K^{\text{Three}}(\mathbf{x}^T \mathbf{z}) g(\mathbf{z}) d\mu(\mathbf{z})$, which is in the same form⁸ as functions in $\mathcal{F}_{(3)}^{\ell_2}$.

⁸We acknowledge that the form here is still not exactly the same as $\mathcal{F}_{(3)}^{\ell_2}$ because the δ -function does not satisfy the constrain of finite $\|g\|_\infty$. Nonetheless, $\mathcal{F}_{(3)}^{\ell_2}$ can be relaxed to allow finite $\|g\|_1$, which then includes the δ -function. See footnote 4 on Page 6.

C A Precise Form of the Upper Bound in Theorem 1

We first introduce some extra notations and a condition about large p_1 that will be used later in our upper bound of the generalization error. Define

$$C(n, d, q) := \frac{\pi - 1}{4\pi} \min \left\{ \frac{1}{2}, \left(\frac{(d-1)^2}{8d} \right)^{\frac{1}{d-1}} (qn)^{-\frac{4}{d-1}} \right\}, \quad (16)$$

$$J(n, p_1, p_2, d, q) := \frac{1}{16\pi d} \sqrt{\frac{C(n, d, q)}{\log(4n)}} - \left(qn^2 \sqrt{\frac{2d}{p_1}} + \frac{q^2 n^3 d}{p_1} + \frac{qn^2}{\sqrt{p_2}} \right), \quad (17)$$

$$Q(p_1, d) := 8d \sqrt{\frac{2(d+1) \log(p_1+1)}{p_1}}. \quad (18)$$

Condition 1. (Given n, d , and $q > 0$) p_1 and p_2 are sufficiently large such that $9d \cdot Q(p_1, d) \leq 1$, $p_1 \geq \left(\frac{10dnq\sqrt{2d}}{C(n, d, q)} \right)^2$, and $J(n, p_1, p_2, d, q) > 0$.

Theorem 3. Given a ground-truth function $f(x) = f_g(x) \in \mathcal{F}_{(3)}^{\ell_2}$, for any $q > 0$, under Condition 1, we must have

$$\Pr_{\mathbf{v}, \mathbf{W}_0, \mathbf{X}} \left\{ |\hat{f}^{\ell_2}(\mathbf{x}) - f(\mathbf{x})| \leq \frac{q\|g\|_\infty}{\sqrt{n}} + \frac{q\|g\|_1}{\sqrt{p_2}} + \sqrt{\frac{Q(p_1, d)}{d}} \|g\|_1 + \frac{\sqrt{n}\|g\|_1 \left(\frac{q}{\sqrt{p_2}} + \sqrt{\frac{Q(p_1, d)}{d}} \right) + \|\epsilon\|_2}{\sqrt{J(n, p_1, p_2, d, q)}} \right\} \geq 1 - \frac{10}{q^2} - \frac{2d^2}{(p_1+1)e^{d+1}}.$$

A proof sketch can be found in Appendix E. To better illustrate the meaning of this upper bound, we provide a simplification in Theorem 1 when p_1 and p_2 are much larger than n . If we view d as a constant, we have $C(n, d, q) = O(n^{-\frac{4}{d-1}})$. When p_1 and p_2 are much larger than n , we have $\frac{\sqrt{n}}{\sqrt{J(n, p_1, p_2, d, q)}} = O\left(n^{\frac{2}{d-1} + \frac{1}{2}} \cdot \sqrt{\log(n)}\right)$ and $\sqrt{\frac{Q(p_1, d)}{d}} = O\left(\sqrt[4]{\frac{\log p_1}{p_1}}\right)$. Therefore, when d is fixed and when both p_1 and p_2 are much larger than n , Theorem 3 can be simplified to Eq. (8) (with high probability).

(In the above reduction to Eq. (8), we ignore the stand-alone term $\frac{q\|g\|_1}{\sqrt{p_2}}$ appeared in Theorem 3, since it is much smaller than the product of Term B and Term E. Similarly, we ignore the stand-alone term $\sqrt{\frac{Q(p_1, d)}{d}}\|g\|_1$, since it is much smaller than the product of Term C and Term E.)

D Noise Effect

Before we present the proof of Theorem 3 in Appendix E, we elaborate on how Theorem 3 reveals the impact of noise on the generalization error. Note that in Eq. (8), Term D denotes the average noise power in each training sample, and Term E denotes the extra multiplication factor with which the noise impacts the generalization error. As we see in Theorem 3 in Appendix C, the precise form of Term E is $\frac{\sqrt{n}}{\sqrt{J(n, p_1, p_2, d, q)}}$. Therefore, we will refer to the multiplication of $\|\epsilon\|_2/\sqrt{n}$ with this factor as the ‘‘noise effect’’. Note that although the precise form $\frac{\sqrt{n}}{\sqrt{J(n, p_1, p_2, d, q)}}$ of this factor in Theorem 3 decreases with respect to both p_1 and p_2 by Eq. (17), when p_1 and p_2 are much larger than n , it can be simplified to Term E, which does not depend on p_1 and p_2 .

In the following, we will analyze the relationship between the noise effect and various system parameters. First, we are interested in know how the numbers of neurons in two hidden-layers p_1 and p_2 impact the noise effect. Since Term E is an approximation when p_1 and p_2 are large and it does not contain p_1 or p_2 , we conjecture that even when p_1 and p_2 are extremely large (e.g., $p_1, p_2 \rightarrow \infty$), the noise effect will neither grow dramatically nor go to zero. Further, when p_1 and p_2 are not so large, by Eq. (17), we know that the precise form of Term E in Theorem 3 decreases when p_1 and




Figure 8: Noise effect on the test MSE of 3-layer NTK (no-bias) with respect to n . The noise follows *i.i.d.* Gaussian distribution with zero mean and variance σ^2 . The ground-truth function is $f(\mathbf{x}) = (\mathbf{x}^T \mathbf{e}_1)^2 + (\mathbf{x}^T \mathbf{e}_1)^3$ where $d = 3$. Every curve is the average of 20 random simulations.

p_2 increase, which suggests that the noise will likely contribute more to the test error when the number of neurons is small. An intuitive explanation of such effect is that when p_1 and p_2 are small, the randomness of the initial weights brings some extra “pseudo-noise” to the model, and thus the generalization performance deteriorates.

Second, we are interested in how the noise effect changes with the number of training data n . We notice that Term E increases with n at a speed faster than \sqrt{n} . However, since it is only an upper bound, the actual noise effect may grow much slower than \sqrt{n} . Therefore, precisely estimating the relationship between n and the noise effect of NTK model would be an interesting future research direction.

We then use simulation to study the noise effect and compare them with the implications derived from our upper bound. In Fig. 8, we plot the curves of the test MSE with respect to n . The noise follows *i.i.d.* Gaussian $\mathcal{N}(0, \sigma^2)$. The blue curve denotes the situation where the noise level is $\sigma^2 = 0.01$. The orange curve denotes the noiseless situation. The noise effect (the value of the gap between the blue and the orange curves) is denoted by the dashed black curve. As we can see, when n is large, the value of the black curve in Fig. 8(a) (fix $p_1 = 200, p_2 = 500$) is higher than that in Fig. 8(b) ($p_1, p_2 \rightarrow \infty$), which validates our conjecture that the noise contributes more to the test error when the number of neurons is small. Further, Fig. 8(b) shows that an infinite number of neurons does not make the noise effect diminish or explode for every n , which also confirms our previous analysis on the relationship between the number of neurons and the noise effect. We also notice that the black curve in Fig. 8(b) (where $p_1, p_2 \rightarrow \infty$) does not increase significantly with n , which suggests that our estimate on how fast Term E increases with n could be further improved.

E Proof of Theorem 3

Recall that Theorem 3 is the precise form of Theorem 1, and is stated in Appendix C. To prove Theorem 3, we follow the line of analysis in Ju et al. (2021). We first study the class of the ground-truth functions that can be learned when weights \mathbf{V} and \mathbf{W}_0 are fixed and there is no noise. We refer to them as *pseudo ground-truth* in the following definition, to differentiate them with the set $\mathcal{F}_{(3)}^{\ell_2}$ of learnable functions for random \mathbf{V} and \mathbf{W}_0 .

Definition 2. Given \mathbf{V} and \mathbf{W}_0 , for any learnable ground-truth function $f_g \in \mathcal{F}_{(3)}^{\ell_2}$ with the corresponding function $g(\cdot)$, define the corresponding **pseudo ground-truth** as

$$\begin{aligned} f_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{x}) &:= \int_{\mathcal{S}^{d-1}} \frac{(\mathbf{h}_{\mathbf{V}, \mathbf{W}_0, z}^{\text{Three}})^T \mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}}}{p_1 p_2} g(z) d\mu(z) \\ &= \int_{\mathcal{S}^{d-1}} (\mathbf{h}_{\mathbf{V}, z}^{\text{RF}})^T \mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}} \frac{\mathcal{C}_{\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}, \mathbf{h}_{\mathbf{V}, z}^{\text{RF}}}^{\mathbf{W}_0}}{p_1 p_2} g(z) d\mu(z). \end{aligned} \quad (19)$$

The last equality of Eq. (19) follows from Eq. (2) and Eq. (11). (The form of Eq. (19) can be derived using the similar process shown in Appendix B.)

We prove Theorem 3 in several steps as follows.

Step 1: use pseudo ground-truth as a ‘‘intermediary’’ .

Recall the definition of pseudo ground-truth $f_{\mathbf{V}, \mathbf{W}_0}^g(\cdot)$ in Eq. (19). We define

$$\mathbf{F}_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{X}) := [f_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{X}_1) f_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{X}_2) \cdots f_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{X}_n)]^T \in \mathbb{R}^n. \quad (20)$$

We then have

$$\begin{aligned} \hat{f}^{\ell_2}(\mathbf{x}) &= \mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}} \mathbf{H}^T (\mathbf{H} \mathbf{H}^T)^{-1} (\mathbf{F}(\mathbf{X}) + \boldsymbol{\epsilon}) \quad (\text{by Eq. (3)}) \\ &= \mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}} \mathbf{H}^T (\mathbf{H} \mathbf{H}^T)^{-1} \mathbf{F}_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{X}) + \mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}} \mathbf{H}^T (\mathbf{H} \mathbf{H}^T)^{-1} (\mathbf{F}(\mathbf{X}) - \mathbf{F}_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{X})) \\ &\quad + \mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}} \mathbf{H}^T (\mathbf{H} \mathbf{H}^T)^{-1} \boldsymbol{\epsilon}. \end{aligned} \quad (21)$$

Thus, we have

$$\begin{aligned} &|\hat{f}^{\ell_2}(\mathbf{x}) - f(\mathbf{x})| \\ &= |\hat{f}^{\ell_2}(\mathbf{x}) - f_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{x}) + f_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{x}) - f(\mathbf{x})| \\ &= |\mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}} \mathbf{H}^T (\mathbf{H} \mathbf{H}^T)^{-1} \mathbf{F}_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{X}) - f_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{x}) \\ &\quad + \mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}} \mathbf{H}^T (\mathbf{H} \mathbf{H}^T)^{-1} (\mathbf{F}(\mathbf{X}) - \mathbf{F}_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{X})) \\ &\quad + f_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{x}) - f(\mathbf{x}) + \mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}} \mathbf{H}^T (\mathbf{H} \mathbf{H}^T)^{-1} \boldsymbol{\epsilon}| \quad (\text{by Eq. (21)}) \\ &\leq \underbrace{|\mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}} \mathbf{H}^T (\mathbf{H} \mathbf{H}^T)^{-1} \mathbf{F}_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{X}) - f_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{x})|}_{\text{term A}} \\ &\quad + \underbrace{|\mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}} \mathbf{H}^T (\mathbf{H} \mathbf{H}^T)^{-1} (\mathbf{F}(\mathbf{X}) - \mathbf{F}_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{X}))|}_{\text{term B}} \\ &\quad + \underbrace{|f_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{x}) - f(\mathbf{x})|}_{\text{term C}} + \underbrace{|\mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}} \mathbf{H}^T (\mathbf{H} \mathbf{H}^T)^{-1} \boldsymbol{\epsilon}|}_{\text{term D}}. \end{aligned} \quad (22)$$

In Eq. (22), term A denotes the test error when using the pseudo ground-truth function, term B denotes the effect of replacing the original ground-truth function by the pseudo ground-truth function in the training samples, term C denotes the difference between the original ground-truth function and the pseudo ground-truth function on the test input, term D denotes the noise effect. Next, we bound these terms one by one.

Step 2: estimate term A.

The following proposition gives an upper bound of the test error when the data model is based on the pseudo ground-truth and the NTK model uses exactly the same \mathbf{V} and \mathbf{W}_0 .

Proposition 4. *Assume fixed \mathbf{V} and \mathbf{W}_0 , (thus p_1 , p_2 and d are also fixed), and there is no noise. If the ground-truth function is $f = f_{\mathbf{V}, \mathbf{W}_0}^g$ in Definition 2 and $\|g\|_\infty < \infty$, then for any $\mathbf{x} \in \mathcal{S}^{d-1}$ and $q > 0$, we must have*

$$\Pr_{\mathbf{X}} \left\{ |f_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{x}) - \hat{f}^{\ell_2}(\mathbf{x})| \geq \frac{q \|g\|_\infty}{\sqrt{n}} \right\} \leq \frac{1}{q^2}.$$

The proof of Proposition 4 is in Appendix G. Proposition 4 captures how the test error decreases with the number of training samples n , if the data model is based on a pseudo ground-truth function with the same \mathbf{V} and \mathbf{W}_0 as the NTK. The result shown in Proposition 4 contributes to Term A in Eq. (8). Here we sketch the proof of Proposition 4. By Eq. (19), we can find a vector $\Delta \mathbf{W}^* \in \mathbb{R}^{(p_1 p_2) \times 1}$ and rewrite $f_{\mathbf{V}, \mathbf{W}_0}^g$ as $f_{\mathbf{V}, \mathbf{W}_0}^g = \mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}} \Delta \mathbf{W}^*$. The specific form of $\Delta \mathbf{W}^*$ can be found in Eq. (34) in Appendix G. Then, by Eq. (3), we can see that the learned model is $\hat{f}^{\ell_2}(\mathbf{x}) = \mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}} \mathbf{P} \Delta \mathbf{W}^*$ where $\mathbf{P} := \mathbf{H}^T (\mathbf{H} \mathbf{H}^T)^{-1} \mathbf{H}$ (an orthogonal projection to the row-space of \mathbf{H}). Thus, we have $|f_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{x}) - \hat{f}^{\ell_2}(\mathbf{x})| = |\mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}} (\mathbf{P} - \mathbf{I}) \Delta \mathbf{W}^*| \leq \|\mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}}\|_2 \cdot \|(\mathbf{P} - \mathbf{I}) \Delta \mathbf{W}^*\|_2$. Further, it is easy to show that $\|\mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}}\|_2 \leq \sqrt{p_1 p_2}$. It then remains to estimate $\|(\mathbf{P} - \mathbf{I}) \Delta \mathbf{W}^*\|_2$, which is

upper bounded by $\min_{\mathbf{a} \in \mathbb{R}^n} \|\Delta \mathbf{W}^* - \mathbf{H}^T \mathbf{a}\|_2$ (because \mathbf{P} is an orthogonal projection). The rest of proof focuses on how to choose a vector \mathbf{a} to make $\|\Delta \mathbf{W}^* - \mathbf{H}^T \mathbf{a}\|_2$ as small as possible. Notice that although the similar method of choosing a suitable \mathbf{a} is also used for 2-layer NTK (Ju et al., 2021), the process of estimating $\|\Delta \mathbf{W}^* - \mathbf{H}^T \mathbf{a}\|_2$ is much more complicated than that in Ju et al. (2021), since the feature vector $\mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}}$ of 3-layer NTK involves non-linear activation for two hidden-layers (instead of one in 2-layer NTK).

With Proposition 4, now we are ready to estimate term A of Eq. (22). We have

$$\begin{aligned} & \Pr_{\mathbf{x}, \mathbf{V}, \mathbf{W}_0} \left\{ \text{term A} \geq \frac{q \|g\|_\infty}{\sqrt{n}} \right\} \\ &= \int_{\mathbb{R}^{d p_1}} \int_{\mathbb{R}^{p_1 p_2}} \Pr_{\mathbf{X}} \left\{ \text{term A} \geq \frac{q \|g\|_\infty}{\sqrt{n}} \right\} d\Lambda_w(\mathbf{W}_0) d\Lambda_v(\mathbf{V}) \\ & \quad (\text{where } \Lambda_w(\cdot) \text{ and } \Lambda_v(\cdot) \text{ are probability distribution of } \mathbf{W}_0 \text{ and } \mathbf{V}, \text{ respectively}) \\ &\leq \frac{1}{q^2} \text{ (by Proposition 4).} \end{aligned}$$

Step 3: estimate term C.

Intuitively, when p_1 and p_2 become larger, the randomness brought by \mathbf{V} and \mathbf{W}_0 in the pseudo ground-truth $f_{\mathbf{V}, \mathbf{W}_0}^g$ will be ‘‘averaged out’’, and thus $f_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{x})$ will approach $f(\mathbf{x})$ (i.e., term C will approaches zero). The following proposition makes this statement rigorous.

Proposition 5. *For any $\mathbf{x} \in \mathcal{S}^{d-1}$ and $q > 0$, we must have*

$$\Pr_{\mathbf{V}, \mathbf{W}_0} \left\{ \left| f_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{x}) - f(\mathbf{x}) \right| \geq \frac{q \|g\|_1}{\sqrt{p_2}} + \sqrt{\frac{Q(p_1, d)}{d}} \|g\|_1 \right\} \leq \frac{d^2}{(p_1 + 1)e^{d+1}} + \frac{1}{q^2}.$$

The proof of Proposition 5 is in Appendix I.1. Note that as p_1 and p_2 increase, both $\frac{q \|g\|_1}{\sqrt{p_2}}$ and $\sqrt{\frac{Q(p_1, d)}{d}} \|g\|_1$ decrease, which implies that the pseudo ground-truth $f_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{x})$ approaches $f(\mathbf{x})$ with high probability. The above result thus directly bounds term C.

Step 4: estimate terms B and D.

We note that both terms B and D are of a similar form. Specifically, we can view the difference between $\mathbf{F}(\mathbf{X})$ and $\mathbf{F}_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{X})$ as a special type of ‘‘noise’’ due to random \mathbf{V} and \mathbf{W}_0 (which will approaches zero when $p_1, p_2 \rightarrow \infty$). Then, both terms B and D are the multiplication of $\mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}} \mathbf{H}^T (\mathbf{H}\mathbf{H}^T)^{-1}$ with the noise (either real noise or the special ‘‘noise’’ above). Further, we can show that the magnitude of $\mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}} \mathbf{H}^T (\mathbf{H}\mathbf{H}^T)^{-1}$ can be upper bounded by a quantity inversely proportional to the minimum eigenvalue of $\mathbf{H}\mathbf{H}^T$. Thus, a key step of the proof is to estimate the minimum eigenvalue of $\mathbf{H}\mathbf{H}^T$. We prove the following proposition about $\min \text{eig}(\mathbf{H}\mathbf{H}^T)$ in Appendix H.

Proposition 6. *Recall the definition of $J(\cdot)$ in Eq. (17). For any $q > 0$, when Condition 1 is satisfied, we must have*

$$\Pr_{\mathbf{x}, \mathbf{V}, \mathbf{W}_0} \left\{ \frac{1}{p_1 p_2} \min \text{eig}(\mathbf{H}\mathbf{H}^T) \leq J(n, p_1, p_2, d, q) \right\} \leq \frac{7}{q^2}.$$

Using Proposition 6, we can then bound terms B and D by the following Proposition 7.

Proposition 7. *For any $q > 0$, when Condition 1 is satisfied, we must have*

$$\begin{aligned} & \Pr_{\mathbf{x}, \mathbf{V}, \mathbf{W}_0} \left\{ \text{term D} + \text{term B of Eq. (22)} \geq \frac{\sqrt{n} \|g\|_1 \left(\frac{q}{\sqrt{p_2}} + \sqrt{\frac{Q(p_1, d)}{d}} \right) + \|\epsilon\|_2}{\sqrt{J(n, p_1, p_2, d, q)}} \right\} \\ &\leq \frac{d^2}{(p_1 + 1)e^{d+1}} + \frac{8}{q^2}. \end{aligned}$$

Note that $\sqrt{n}\|g\|_1 \left(\frac{q}{\sqrt{p^2}} + \sqrt{\frac{Q(p_1, d)}{d}} \right)$ and $\|\epsilon\|_2$ correspond to the magnitude of the special ‘‘noise’’ ($\mathbf{F}(\mathbf{X}) - \mathbf{F}_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{X})$) (which can be bounded just like Proposition 5) and the real noise ϵ , respectively. The proof of Proposition 7 is in Appendix I.2.

Plugging the results in Steps 2, 3, and 4 into Eq. (22), the result of Theorem 3 thus follows. Appendices G to I will prove the above propositions, after we present some supporting lemmas in Appendix F.

F Useful Notations and Lemmas

We first collect some useful notations and lemmas, which will be used in the proofs of propositions appeared in Appendix E, as well as the analysis of learnable functions. Let $I(\cdot, \cdot)$ denote the regularized incomplete beta function Dutka (1981). Let $B(\cdot, \cdot)$ denote the beta function Chaudhry et al. (1997). Specifically,

$$B(x, y) := \int_0^1 t^{x-1} (1-t)^{y-1} dt, \quad (23)$$

$$I_x(a, b) := \frac{\int_0^x t^{a-1} (1-t)^{b-1} dt}{B(a, b)}. \quad (24)$$

Define a cap on a unit hyper-sphere \mathcal{S}^{d-1} as the intersection of \mathcal{S}^{d-1} with an open ball in \mathbb{R}^d centered at \mathbf{v}_* with radius r , i.e.,

$$\mathcal{B}_{\mathbf{v}_*}^r := \{ \mathbf{v} \in \mathcal{S}^{d-1} \mid \|\mathbf{v} - \mathbf{v}_*\|_2 < r \}. \quad (25)$$

Remark 2. For ease of exposition, we will sometimes neglect the subscript \mathbf{v}_* of $\mathcal{B}_{\mathbf{v}_*}^r$ and use \mathcal{B}^r instead, when the quantity that we are estimating only depends on r but not \mathbf{v}_* . For example, where we are interested in the area of $\mathcal{B}_{\mathbf{v}_*}^r$, it only depends on r but not \mathbf{v}_* . Thus, we write $\lambda_{d-1}(\mathcal{B}^r)$ instead.

F.1 Quantities related to the area of a cap on a hyper-sphere

The lemmas of this subsection support for the proof of Proposition 6. The following lemma is introduced by Li (2011), which gives the area of a cap on a hyper-sphere with respect to the colatitude angle.

Lemma 8. *Let $\phi \in [0, \frac{\pi}{2}]$ denote the colatitude angle of the smaller cap on the unit hyper-sphere \mathcal{S}^{a-1} , then the area (in the measure of λ_{a-1}) of this hyper-spherical cap is*

$$\frac{1}{2} \lambda_{a-1}(\mathcal{S}^{a-1}) I_{\sin^2 \phi} \left(\frac{a-1}{2}, \frac{1}{2} \right),$$

or equivalently⁹,

$$\lambda_{a-1}(\mathcal{B}^r) = \frac{1}{2} \lambda_{a-1}(\mathcal{S}^{a-1}) I_{r^2(1-\frac{r^2}{4})} \left(\frac{a-1}{2}, \frac{1}{2} \right).$$

where $r \leq \sqrt{2}$.

The following lemma is shown by Lemma 35 of Ju et al. (2021).

Lemma 9. *For any $x \in [0, 1]$, we must have*

$$I_x \left(\frac{a-1}{2}, \frac{1}{2} \right) \in \left[\frac{2x^{\frac{a-1}{2}}}{B(\frac{a-1}{2}, \frac{1}{2}) \cdot (a-1)}, \frac{2x^{\frac{a-1}{2}}}{B(\frac{a-1}{2}, \frac{1}{2}) \cdot (a-1)\sqrt{1-x}} \right].$$

The following lemma is shown by Lemma 32 of Ju et al. (2021).

⁹Proof of this equivalence can be found in Lemma 9 of Ju et al. (2021).

Lemma 10. For any integer $a \geq 2$,

$$B\left(\frac{a-1}{2}, \frac{1}{2}\right) \in \left[\frac{1}{\sqrt{a}}, \pi\right].$$

Further, if $a \geq 5$, we have

$$B\left(\frac{a-1}{2}, \frac{1}{2}\right) \in \left[\frac{1}{\sqrt{a}}, \frac{4}{\sqrt{a-3}}\right].$$

F.2 Estimation of certain norms

In our proofs, we will often need to estimate the norms of the NTK feature vectors. We list some useful lemmas below.

Lemma 11. For any $\mathbf{x} \in \mathcal{S}^{d-1}$, we have

$$\|\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}\|_2 \leq \sqrt{p_1}, \quad \|\mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}}\|_2 \leq \sqrt{p_1 p_2}.$$

Proof. Notice that $\|\mathbf{x}\|_2 = 1$ and $\|\mathbf{V}[j]\|_2 = 1$ for all $j \in \{1, 2, \dots, p_1\}$. By Eq. (1), we have

$$\|\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}\|_2 = \sqrt{\sum_{j=1}^{p_1} ((\mathbf{x}^T \mathbf{V}[j]) \mathbf{1}_{\{\mathbf{x}^T \mathbf{V}[j] > 0\}})^2} \leq \sqrt{p_1}.$$

Thus, by Eq. (2), we have

$$\|\mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}}\|_2 = \sqrt{\sum_{k=1}^{p_2} \|\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}} \mathbf{1}_{\{(\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}})^T \mathbf{W}_0[k] > 0\}}\|_2^2} \leq \sqrt{\sum_{k=1}^{p_2} \|\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}\|_2^2} \leq \sqrt{p_1 p_2}.$$

□

The following lemma is from Lemma 12 of [Ju et al. \(2021\)](#), but we repeat here for the convenience of the readers.

Lemma 12. If $\mathbf{C} = \mathbf{A}\mathbf{B}$, then $\|\mathbf{C}\|_2 \leq \|\mathbf{A}\|_2 \cdot \|\mathbf{B}\|_2$. Here \mathbf{A} , \mathbf{B} , and \mathbf{C} could be scalars, vectors, or matrices.

Proof. This lemma directly follows the definition of matrix norm. □

Remark 3. Note that the (ℓ_2) matrix-norm (i.e., spectral norm) of a vector is exactly its ℓ_2 vector-norm (i.e., Euclidean norm)¹⁰. Therefore, when applying Lemma 12, we do not need to worry about whether \mathbf{A} , \mathbf{B} , and \mathbf{C} are matrices or vectors.

Lemma 13. For any $\mathbf{A}, \mathbf{B} \in \mathbb{R}^{k \times k}$, we must have

$$\|\mathbf{A} - \mathbf{B}\|_2 \leq k \cdot \max_{i,j} |\mathbf{A}_{i,j} - \mathbf{B}_{i,j}|.$$

Consequently, if both \mathbf{A} and \mathbf{B} are positive semi-definite, then

$$|\min \text{eig}(\mathbf{A}) - \min \text{eig}(\mathbf{B})| \leq k \cdot \max_{i,j} |\mathbf{A}_{i,j} - \mathbf{B}_{i,j}|.$$

¹⁰To see this, consider a (row or column) vector \mathbf{a} . The matrix norm of \mathbf{a} is

$$\begin{aligned} & \max_{\|\mathbf{x}\|=1} \|\mathbf{a}\mathbf{x}\|_2 \text{ (when } \mathbf{a} \text{ is a column vector),} \\ & \text{or } \max_{\|\mathbf{x}\|_2=1} \|\mathbf{a}\mathbf{x}\|_2 \text{ (when } \mathbf{a} \text{ is a row vector).} \end{aligned}$$

In both cases, the value of the matrix-norm equals to $\sqrt{\sum a_i^2}$, which is exactly the ℓ_2 -norm (Euclidean norm) of \mathbf{a} .

Proof. Let $\mathbf{C} := \mathbf{A} - \mathbf{B}$. For any $\mathbf{a} \in \mathcal{S}^{k-1}$, we have

$$\begin{aligned} \|\mathbf{C}\mathbf{a}\|_2^2 &= \sum_{i=1}^k \left(\sum_{j=1}^k \mathbf{C}_{i,j} a_j \right)^2 \\ &\leq k (\max_{i,j} \mathbf{C}_{i,j})^2 \left(\sum_{j=1}^k a_j \right)^2 \\ &\leq k^2 (\max_{i,j} \mathbf{C}_{i,j})^2 \sum_{j=1}^k a_j^2 \quad (\text{by Cauchy-Schwarz inequality}) \\ &= k^2 (\max_{i,j} \mathbf{C}_{i,j})^2 \quad (\text{because } \|\mathbf{a}\|_2 = 1). \end{aligned}$$

Because $\|\mathbf{C}\|_2 = \max_{\mathbf{a} \in \mathcal{S}^{k-1}} \|\mathbf{C}\mathbf{a}\|_2$, we have $\|\mathbf{A} - \mathbf{B}\|_2 \leq k \cdot \max_{i,j} |\mathbf{A}_{i,j} - \mathbf{B}_{i,j}|$.

Let $\mathbf{a}^* \in \arg \min_{\mathbf{a} \in \mathcal{S}^{k-1}} \|\mathbf{B}\mathbf{a}\|_2$. We have

$$\begin{aligned} \min \text{eig}(\mathbf{A}) &= \min_{\mathbf{a} \in \mathcal{S}^{k-1}} \|\mathbf{A}\mathbf{a}\|_2 \\ &\leq \|\mathbf{A}\mathbf{a}^*\|_2 \\ &= \|(\mathbf{A} - \mathbf{B})\mathbf{a}^* + \mathbf{B}\mathbf{a}^*\|_2 \\ &\leq \|(\mathbf{A} - \mathbf{B})\mathbf{a}^*\|_2 + \|\mathbf{B}\mathbf{a}^*\|_2 \\ &\leq \|\mathbf{A} - \mathbf{B}\|_2 + \min \text{eig}(\mathbf{B}) \quad (\text{by the definition of } \mathbf{a}^*). \end{aligned}$$

Thus, we have $\min \text{eig}(\mathbf{A}) - \min \text{eig}(\mathbf{B}) \leq \|\mathbf{A} - \mathbf{B}\|_2 \leq k \cdot \max_{i,j} |\mathbf{C}_{i,j}|$. Similarly, we have $\min \text{eig}(\mathbf{B}) - \min \text{eig}(\mathbf{A}) \leq k \cdot \max_{i,j} |\mathbf{C}_{i,j}|$. The result of this lemma thus follows. \square

F.3 Estimates of certain tail probabilities

Lemma 14 (Chebyshev's inequality on the sum of *i.i.d.* random variables/vectors). *Let X_1, X_2, \dots, X_k be i.i.d. random variables and $|X_i| \leq U$ for all $i = 1, 2, \dots, k$. Then, for any $m > 0$,*

$$\Pr \left\{ \left| \left(\frac{1}{k} \sum_{i=1}^k X_i \right) - \mathbb{E} X_1 \right| \geq \frac{mU}{\sqrt{k}} \right\} \leq \frac{1}{m^2}.$$

This inequality also holds when X_1, X_2, \dots, X_k are i.i.d. random vectors and $\|X_i\|_2 \leq U$ for all $i = 1, 2, \dots, k$.

Proof. Because $|X_1| \leq U$, we have

$$\text{Var}[X_1] = \mathbb{E}[(X_1 - \mathbb{E}[X_1])^2] = \mathbb{E}[X_1^2] - (\mathbb{E}[X_1])^2 \leq \mathbb{E}[X_1^2] \leq U^2.$$

Because all X_i 's are *i.i.d.*, we have

$$\text{Var} \left[\frac{1}{k} \sum_{i=1}^k X_i \right] \leq \frac{U^2}{k}, \quad \mathbb{E} \left[\frac{1}{k} \sum_{i=1}^k X_i \right] = \mathbb{E}[X_1].$$

The result of this lemma thus follows by applying Chebyshev's inequality on $\frac{1}{k} \sum_{i=1}^k X_i$. For the situation that X_1, X_2, \dots, X_k are vectors, the proof is the same by using the generalized Chebyshev's inequality for random vectors which we state in Lemma 15 as follows. \square

The following is the Chebyshev's inequality for random vectors that can be found in many textbooks of probability theory (see, e.g., pp. 446-451 of [Laha & Rohatgi \(1979\)](#)).

Lemma 15 (Chebyshev's inequality for random vectors). *For a random vector $\mathbf{w} \in \mathbb{R}^a$ with probability distribution $\Lambda(\cdot)$, for any $\delta > 0$, we must have*

$$\Pr \{ \|\mathbf{w} - \mathbb{E}(\mathbf{w})\|_2 \geq \delta \} \leq \frac{\text{Var}(\mathbf{w})}{\delta^2},$$

where

$$\text{Var}(\mathbf{w}) := \int_{\mathbf{v} \in \mathbb{R}^a} \|\mathbf{v} - \mathbb{E}(\mathbf{w})\|_2^2 d\Lambda(\mathbf{v}).$$

F.4 Estimation about double factorial

Let m be a positive integer. A double factorial can be defined by

$$(2m)!! := \prod_{i=1}^m (2i), \quad (2m-1)!! := \prod_{i=1}^m (2i-1). \quad (26)$$

They are useful in our study of learnable functions. The following lemma is proven by [Chen & Qi \(2005\)](#).

Lemma 16 (Improved Wallis' Inequality). *For all natural numbers k , let $k!!$ denote a double factorial. Then*

$$\frac{1}{\sqrt{\pi \left(k + \frac{4}{\pi} - 1\right)}} \leq \frac{(2k-1)!!}{(2k)!!} < \frac{1}{\sqrt{\pi \left(k + \frac{1}{4}\right)}}.$$

Further, the constants $\frac{4}{\pi} - 1$ and $\frac{1}{4}$ are the best possible.

F.5 Taylor expansion of kernels

The following Taylor expansions are related to the NTK kernel functions, which will also be used in our characterization of the learnable functions.

Lemma 17. *For any $\theta \in [0, \pi]$,*

$$\begin{aligned} \cos \theta \frac{(\pi - \theta)}{2\pi} &= \frac{\cos \theta}{4} + \frac{1}{2\pi} \sum_{k=0}^{\infty} \frac{(2k)!}{(k!)^2} \frac{4}{2k+1} \left(\frac{\cos \theta}{2}\right)^{2k+2}, \\ \frac{\sin \theta + (\pi - \theta) \cos \theta}{\pi} &= \frac{1}{\pi} \left(1 + \frac{\pi}{2} \cos \theta + \sum_{k=0}^{\infty} \frac{2(2k)!}{(k+1)(2k+1)(k!)^2} \left(\frac{\cos \theta}{2}\right)^{2k+2} \right). \end{aligned}$$

Consequently, recalling Eq. (5) and Eq. (4), by letting $a = \cos \theta$, we have

$$\begin{aligned} K^{Two}(a) &= a \frac{\pi - \arccos a}{2\pi} = \frac{a}{4} + \frac{1}{2\pi} \sum_{k=0}^{\infty} \frac{(2k)!}{(k!)^2} \frac{4}{2k+1} \left(\frac{a}{2}\right)^{2k+2}, \\ 2d \cdot K^{RF}(a) &= \frac{\sqrt{1-a^2} + a(\pi - \arccos a)}{\pi} \\ &= \frac{1}{\pi} \left(1 + \frac{\pi}{2} a + \sum_{k=0}^{\infty} \frac{2(2k)!}{(k+1)(2k+1)(k!)^2} \left(\frac{a}{2}\right)^{2k+2} \right). \end{aligned}$$

Proof. Using Taylor expansion on $\arccos x$, we have

$$\arccos(x) = \frac{\pi}{2} - \sum_{k=0}^{\infty} \frac{(2k)!}{2^{2k}(k!)^2} \frac{x^{2k+1}}{2k+1}.$$

We then have

$$\theta = \arccos(\cos \theta) = \frac{\pi}{2} - \sum_{k=0}^{\infty} \frac{(2k)!}{(k!)^2} \frac{2}{2k+1} \left(\frac{\cos \theta}{2}\right)^{2k+1}.$$

Thus, we have

$$\begin{aligned} \cos \theta \frac{(\pi - \theta)}{2\pi} &= \cos \theta \cdot \left(\frac{1}{2} - \frac{1}{2\pi} \left(\frac{\pi}{2} - \sum_{k=0}^{\infty} \frac{(2k)!}{(k!)^2} \frac{2}{2k+1} \left(\frac{\cos \theta}{2}\right)^{2k+1} \right) \right) \\ &= \frac{\cos \theta}{4} + \frac{1}{2\pi} \sum_{k=0}^{\infty} \frac{(2k)!}{(k!)^2} \frac{4}{2k+1} \left(\frac{\cos \theta}{2}\right)^{2k+2}. \end{aligned} \quad (27)$$

Using Taylor expansion on $\sqrt{1+x}$, we have

$$\sqrt{1+x} = 1 - \sum_{k=0}^{\infty} \frac{2}{k+1} \binom{2k}{k} \left(-\frac{x}{4}\right)^{k+1},$$

Replacing x by $-\cos^2 \theta$, we thus have

$$\sin \theta = \sqrt{1 - \cos^2 \theta} = 1 - \sum_{k=0}^{\infty} \frac{2}{k+1} \binom{2k}{k} \left(\frac{\cos \theta}{2}\right)^{2k+2}.$$

Therefore, using Eq. (27) again, we have

$$\begin{aligned} \frac{\sin \theta + (\pi - \theta) \cos \theta}{\pi} &= \frac{1}{\pi} \left(1 + \frac{\pi}{2} \cos \theta + \sum_{k=0}^{\infty} \left(\frac{2}{2k+1} - \frac{1}{k+1} \right) \frac{2(2k)!}{(k!)^2} \left(\frac{\cos \theta}{2}\right)^{2k+2} \right) \\ &= \frac{1}{\pi} \left(1 + \frac{\pi}{2} \cos \theta + \sum_{k=0}^{\infty} \frac{2(2k)!}{(k+1)(2k+1)(k!)^2} \left(\frac{\cos \theta}{2}\right)^{2k+2} \right). \end{aligned}$$

The result of this lemma thus follows. \square

E.6 Calculation of certain integrals

Lemma 18. For any integer $k \geq 2$, we have

$$\int_0^{\pi} \sin^k \varphi \, d\varphi = \frac{k-1}{k} \int_0^{\pi} \sin^{k-2} \varphi \, d\varphi.$$

Proof. We have

$$\begin{aligned} \int_0^{\pi} \sin^k \varphi \, d\varphi &= \int_0^{\pi} \sin \varphi \cdot \sin^{k-1} \varphi \, d\varphi \\ &= -\cos \varphi \cdot \sin^{k-1} \varphi \Big|_0^{\pi} + (k-1) \int_0^{\pi} \cos^2 \varphi \cdot \sin^{k-2} \varphi \, d\varphi \\ &\quad \text{(integration by parts)} \\ &= (k-1) \int_0^{\pi} (1 - \sin^2 \varphi) \sin^{k-2} \varphi \, d\varphi \\ &= (k-1) \int_0^{\pi} \sin^{k-2} \varphi \, d\varphi - (k-1) \int_0^{\pi} \sin^k \varphi \, d\varphi. \end{aligned}$$

Moving the second term of the right hand side to the left hand side, we have

$$k \int_0^{\pi} \sin^k \varphi \, d\varphi = (k-1) \int_0^{\pi} \sin^{k-2} \varphi \, d\varphi.$$

The result of this lemma thus follows. \square

Lemma 19. For any $\theta \in [0, \pi]$,

$$\int_{-\frac{\pi}{2}+\theta}^{\frac{\pi}{2}} \cos(\alpha) \cos(\alpha - \theta) \, d\alpha = \frac{\sin \theta}{2} + \frac{(\pi - \theta) \cos \theta}{2}.$$

Proof. Notice that

$$\begin{aligned} \frac{\partial(\sin(2\alpha - \theta) + 2\alpha \cos \theta)}{\partial \alpha} &= 2 \cos(2\alpha - \theta) + 2 \cos \theta \\ &= 2 \cos(\alpha + (\alpha - \theta)) + 2 \cos(\alpha - (\alpha - \theta)) \\ &= 4 \cos(\alpha) \cos(\alpha - \theta). \end{aligned}$$

Thus, we have

$$\int \cos(\alpha) \cos(\alpha - \theta) d\alpha = \frac{\sin(2\alpha - \theta) + 2\alpha \cos(\theta)}{4} + \text{constant}.$$

Notice that

$$\begin{aligned} \sin(2\alpha - \theta) \Big|_{\alpha = -\frac{\pi}{2} + \theta}^{\frac{\pi}{2}} &= \sin(\pi - \theta) - \sin(\theta - \pi) = 2 \sin \theta, \\ 2\alpha \cos(\theta) \Big|_{\alpha = -\frac{\pi}{2} + \theta}^{\frac{\pi}{2}} &= 2(\pi - \theta) \cos \theta. \end{aligned}$$

The result of this lemma thus follows. \square

Lemma 20. Recall that $\gamma(\cdot)$ denotes the probability density function of $\mathbf{W}_0[k]$ and is unif(\mathcal{S}^{p_1-1}) by Assumption 1. For any $\mathbf{a}, \mathbf{b} \in \mathbb{R}^{p_1}$, we have

$$\int_{\mathcal{S}^{p_1-1}} \mathbf{a}^T \mathbf{b} \cdot \mathbf{1}_{\{\mathbf{a}^T \mathbf{w} > 0, \mathbf{b}^T \mathbf{w} > 0\}} d\gamma(\mathbf{w}) = \mathbf{a}^T \mathbf{b} \frac{\pi - \arccos\left(\frac{\mathbf{a}^T \mathbf{b}}{\|\mathbf{a}\|_2 \|\mathbf{b}\|_2}\right)}{2\pi}.$$

(Although the right hand side is not defined when $\mathbf{a} = \mathbf{0}$ or $\mathbf{b} = \mathbf{0}$, we can artificially re-define the value of the right hand side as 0 when $\mathbf{a} = \mathbf{0}$ or $\mathbf{b} = \mathbf{0}$, so the equation still holds.)

Proof. The result holds trivially when $\mathbf{a} = \mathbf{0}$ or $\mathbf{b} = \mathbf{0}$. When \mathbf{a} and \mathbf{b} are both non-zero, it suffices to prove that

$$\int_{\mathcal{S}^{p_1-1}} \mathbf{1}_{\{\mathbf{a}^T \mathbf{w} > 0, \mathbf{b}^T \mathbf{w} > 0\}} d\gamma(\mathbf{w}) = \frac{\pi - \arccos\left(\frac{\mathbf{a}^T \mathbf{b}}{\|\mathbf{a}\|_2 \|\mathbf{b}\|_2}\right)}{2\pi},$$

which has been proven by Lemma 17 of Ju et al. (2021) (where its geometric explanation is given as well). \square

Lemma 21. For any $\mathbf{x}, \mathbf{z} \in \mathcal{S}^{d-1}$, we have

$$\int_{\mathcal{S}^{d-1}} (\mathbf{x}^T \mathbf{v})(\mathbf{z}^T \mathbf{v}) \mathbf{1}_{\{\mathbf{z}^T \mathbf{v} > 0, \mathbf{x}^T \mathbf{v} > 0\}} d\lambda(\mathbf{v}) = \frac{\sin \theta + (\pi - \theta) \cos \theta}{2d\pi}, \quad (28)$$

where θ denotes the angle between \mathbf{x} and \mathbf{z} , i.e.,

$$\theta = \arccos(\mathbf{x}^T \mathbf{z}) \in [0, \pi]. \quad (29)$$

To help readers understand the correctness of Lemma 21, we first give a simple proof for the special case that $d = 2$, i.e., when vectors \mathbf{x}, \mathbf{z} , and \mathbf{v} are all in the 2-D plane. Then we prove Lemma 21 for the general cases that $d = 2, 3, 4, \dots$.

Proof (of the case when $d = 2$): Without loss of generality, we let

$$\mathbf{v} = \begin{bmatrix} \cos \alpha \\ \sin \alpha \end{bmatrix}, \quad \mathbf{z} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad \text{and } \mathbf{x} = \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix}.$$

Thus, we have

$$\begin{aligned} \text{The left-hand-side of Eq. (28)} &= \frac{1}{2\pi} \int_{(\theta - \frac{\pi}{2}, \theta + \frac{\pi}{2}) \cap (-\frac{\pi}{2}, \frac{\pi}{2})} (\cos \alpha \cos \theta + \sin \alpha \sin \theta) \cos \alpha d\alpha \\ &= \frac{1}{2\pi} \int_{\theta - \frac{\pi}{2}}^{\frac{\pi}{2}} \cos(\alpha - \theta) \cos \alpha d\alpha \quad (\text{since } \theta \in [0, \pi]) \\ &= \frac{\sin \theta + (\pi - \theta) \cos \theta}{4\pi} \quad (\text{by Lemma 19}). \end{aligned}$$

Proof (of the general case). Due to symmetry, we know that the integral in the left-hand-side of Eq. (28) only depends on the angle between \mathbf{x} and \mathbf{z} . Thus, without loss of generality, we let

$$\mathbf{x} = [x_1 \ x_2 \ \cdots \ x_d] = [0 \ 0 \ \cdots \ 0 \ 1 \ 0]^T, \quad \mathbf{z} = [0 \ 0 \ \cdots \ 0 \ \cos \theta \ \sin \theta]^T.$$

Thus, for any $\mathbf{v} = [v_1 \ v_2 \ \cdots \ v_d]^T$, in order for $\mathbf{z}^T \mathbf{v} > 0$ and $\mathbf{x}^T \mathbf{v} > 0$ to hold, it only needs to satisfy

$$[\cos \theta \ \sin \theta] \begin{bmatrix} v_{d-1} \\ v_d \end{bmatrix} > 0, \quad [1 \ 0] \begin{bmatrix} v_{d-1} \\ v_d \end{bmatrix} > 0. \quad (30)$$

We use the spherical coordinate $\varphi_{\mathbf{x}} = [\varphi_1^{\mathbf{x}} \ \varphi_2^{\mathbf{x}} \ \cdots \ \varphi_{d-1}^{\mathbf{x}}]^T$ where $\varphi_1^{\mathbf{x}}, \dots, \varphi_{d-2}^{\mathbf{x}} \in [0, \pi]$ and $\varphi_{d-1}^{\mathbf{x}} \in [0, 2\pi)$ with the convention that

$$\begin{aligned} x_1 &= \cos(\varphi_1^{\mathbf{x}}), \\ x_2 &= \sin(\varphi_1^{\mathbf{x}}) \cos(\varphi_2^{\mathbf{x}}), \\ x_3 &= \sin(\varphi_1^{\mathbf{x}}) \sin(\varphi_2^{\mathbf{x}}) \cos(\varphi_3^{\mathbf{x}}), \\ &\vdots \\ x_{d-1} &= \sin(\varphi_1^{\mathbf{x}}) \sin(\varphi_2^{\mathbf{x}}) \cdots \sin(\varphi_{d-2}^{\mathbf{x}}) \cos(\varphi_{d-1}^{\mathbf{x}}), \\ x_d &= \sin(\varphi_1^{\mathbf{x}}) \sin(\varphi_2^{\mathbf{x}}) \cdots \sin(\varphi_{d-2}^{\mathbf{x}}) \sin(\varphi_{d-1}^{\mathbf{x}}). \end{aligned}$$

Thus, we have $\varphi_{\mathbf{x}} = [\pi/2 \ \pi/2 \ \cdots \ \pi/2 \ 0]^T$. Similarly, the spherical coordinate for \mathbf{z} is $\varphi_{\mathbf{z}} = [\pi/2 \ \pi/2 \ \cdots \ \pi/2 \ \theta]^T$. Let the spherical coordinates for \mathbf{v} be $\varphi_{\mathbf{v}} = [\varphi_1^{\mathbf{v}} \ \varphi_2^{\mathbf{v}} \ \cdots \ \varphi_{d-1}^{\mathbf{v}}]^T$. Thus, Eq. (30) is equivalent to

$$\mathbf{z}^T \mathbf{v} = \sin(\varphi_1^{\mathbf{v}}) \sin(\varphi_2^{\mathbf{v}}) \cdots \sin(\varphi_{d-2}^{\mathbf{v}}) (\cos \theta \cos(\varphi_{d-1}^{\mathbf{v}}) + \sin \theta \sin(\varphi_{d-1}^{\mathbf{v}})) > 0, \quad (31)$$

$$\mathbf{x}^T \mathbf{v} = \sin(\varphi_1^{\mathbf{v}}) \sin(\varphi_2^{\mathbf{v}}) \cdots \sin(\varphi_{d-2}^{\mathbf{v}}) \cos(\varphi_{d-1}^{\mathbf{v}}) > 0. \quad (32)$$

Because $\varphi_1^{\mathbf{v}}, \dots, \varphi_{d-2}^{\mathbf{v}} \in [0, \pi]$ (by the convention of spherical coordinates), we have

$$\sin(\varphi_1^{\mathbf{v}}) \sin(\varphi_2^{\mathbf{v}}) \cdots \sin(\varphi_{d-2}^{\mathbf{v}}) \geq 0.$$

Thus, for Eq. (31) and Eq. (32) to hold, we must have

$$\cos(\theta - \varphi_{d-1}^{\mathbf{v}}) > 0, \quad \cos(\varphi_{d-1}^{\mathbf{v}}) > 0,$$

i.e., $\varphi_{d-1}^{\mathbf{v}} \in (-\pi/2, \pi/2) \cap (\theta - \pi/2, \theta + \pi/2) \pmod{2\pi}$. By Eq. (29), we thus have

$$\varphi_{d-1} \in \left(-\frac{\pi}{2} + \theta, \frac{\pi}{2}\right) \pmod{2\pi}.$$

Let

$$A(\theta, \varphi_{d-1}^{\mathbf{v}}) := (\cos \theta \cos(\varphi_{d-1}^{\mathbf{v}}) + \sin \theta \sin(\varphi_{d-1}^{\mathbf{v}})) \cos(\varphi_{d-1}^{\mathbf{v}}) = \cos(\varphi_{d-1}^{\mathbf{v}} - \theta) \cos \varphi_{d-1}^{\mathbf{v}}.$$

By Eq. (31) and Eq. (32), we have

$$(\mathbf{x}^T \mathbf{v})(\mathbf{z}^T \mathbf{v}) \mathbf{1}_{\{\mathbf{z}^T \mathbf{v} > 0, \mathbf{x}^T \mathbf{v} > 0\}} = \sin^2(\varphi_1^{\mathbf{v}}) \sin^2(\varphi_2^{\mathbf{v}}) \cdots \sin^2(\varphi_{d-2}^{\mathbf{v}}) A(\theta, \varphi_{d-1}^{\mathbf{v}}).$$

Integrating using such spherical coordinates, we have

$$\begin{aligned} & \int_{\mathcal{S}^{d-1}} (\mathbf{x}^T \mathbf{v})(\mathbf{z}^T \mathbf{v}) \mathbf{1}_{\{\mathbf{z}^T \mathbf{v} > 0, \mathbf{x}^T \mathbf{v} > 0\}} d\lambda(\mathbf{v}) \\ &= \frac{\int_{-\frac{\pi}{2} + \theta}^{\frac{\pi}{2}} A(\theta, \varphi_{d-1}^{\mathbf{v}}) \int_0^\pi \cdots \int_0^\pi \sin^d(\varphi_1) \sin^{d-1}(\varphi_2) \cdots \sin^3(\varphi_{d-2}) d\varphi_1 d\varphi_2 \cdots d\varphi_{d-1}}{\int_0^{2\pi} \int_0^\pi \cdots \int_0^\pi \sin^{d-2}(\varphi_1) \sin^{d-3}(\varphi_2) \cdots \sin(\varphi_{d-2}) d\varphi_1 d\varphi_2 \cdots d\varphi_{d-1}} \\ &= \frac{\int_{-\frac{\pi}{2} + \theta}^{\frac{\pi}{2}} A(\theta, \varphi_{d-1}^{\mathbf{v}}) \cdot d\varphi_{d-1}}{\int_0^{2\pi} d\varphi_{d-1}} \cdot \frac{d-1}{d} \frac{d-2}{d-1} \cdots \frac{2}{3} \quad (\text{by Lemma 18}) \\ &= \frac{\sin \theta + (\pi - \theta) \cos \theta}{2d \cdot \pi} \quad (\text{by Lemma 19}). \end{aligned}$$

The result of this lemma thus follows. \square

E.7 Convergence of $\frac{1}{p_1}(\mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V},\mathbf{z}}^{\text{RF}}$ with respect to p_1

Lemma 22 (Theorem 4.2 of [Wainwright \(2015\)](#)). *Let \mathcal{F} be a class of real-valued functions f such that $\|f\|_\infty \leq b$ for all $f \in \mathcal{F}$. Then for all $k \geq 1$ and $\delta \geq 0$, we have*

$$\Pr \left\{ \sup_{f \in \mathcal{F}} \left| \frac{1}{k} \sum_{i=1}^k f(X_i) - \mathbb{E}_{x \sim \mathcal{X}(\cdot)} f(x) \right| \leq 2\mathcal{R}_k(\mathcal{F}) + \delta \right\} \geq 1 - \exp\left(-\frac{k\delta^2}{8b^2}\right),$$

where $\mathcal{R}_k(\mathcal{F})$ denotes the Rademacher complexity, X_1, X_2, \dots, X_k are i.i.d. random variables/vectors that follow the distribution $\mathcal{X}(\cdot)$.

Polynomial discrimination. A class \mathcal{F} of functions with domain \mathcal{X} has polynomial discrimination of order $\nu \geq 1$ if for each positive integer k and collection $X_1^k = \{X_1, \dots, X_k\}$ of k points in \mathcal{X} , the set $\mathcal{F}(X_1^k)$ has cardinality upper bounded by

$$\text{card}(\mathcal{F}(X_1^k)) \leq (k+1)^\nu.$$

Lemma 23 (Lemma 4.1 and Eq. (4.23) of [Wainwright \(2015\)](#)). *Suppose that \mathcal{F} has polynomial discrimination of order ν and $\|f\|_\infty \leq b$ for all $f \in \mathcal{F}$. Then*

$$\mathcal{R}_k(\mathcal{F}) \leq 3\sqrt{\frac{b^2\nu \log(k+1)}{k}} \quad \text{for all } k \geq 10.$$

Given a function $h : \mathcal{S}^{d-1} \mapsto \mathbb{R}$ such that $\|h\|_\infty < \infty$ and given any $\delta > 0$, consider the function class \mathcal{F}_* that consists of functions $h(\mathbf{v})\mathbf{1}_{\{\mathbf{x}^T \mathbf{v} > 0, \mathbf{z}^T \mathbf{v} > 0\}}$, which maps $\mathbf{v} \in \mathcal{S}^{d-1}$ to either 0 or $h(\mathbf{v})$. By Lemma 20 of [Ju et al. \(2021\)](#), we have

$$\text{card}(\mathcal{F}_*(X_1^k)) \leq (k+1)^{2(d+1)}.$$

(Here X_1^k corresponds to $\{\mathbf{V}[1], \dots, \mathbf{V}[k]\}$.) Thus, combined with Lemma 22 and Lemma 23, we have

$$\begin{aligned} & \Pr \left\{ \max_{\mathbf{x}, \mathbf{z}} \left| \frac{1}{p_1} \sum_{j=1}^{p_1} h(\mathbf{V}[j])\mathbf{1}_{\{\mathbf{x}^T \mathbf{V}[j] > 0, \mathbf{z}^T \mathbf{V}[j] > 0\}} - \mathbb{E}_{\mathbf{v}}[h(\mathbf{v})\mathbf{1}_{\{\mathbf{x}^T \mathbf{v} > 0, \mathbf{z}^T \mathbf{v} > 0\}}] \right| \right. \\ & \quad \left. \leq 6\sqrt{\frac{\|h\|_\infty^2 2(d+1) \log(p_1+1)}{p_1}} + \delta \right\} \geq 1 - \exp\left(-\frac{p_1\delta^2}{8\|h\|_\infty^2}\right). \end{aligned}$$

Further, if we let $\delta = 2\sqrt{\frac{\|h\|_\infty^2 2(d+1) \log(p_1+1)}{p_1}}$, we have proven the following lemma.

Lemma 24. *For any given function $h : \mathcal{S}^{d-1} \mapsto \mathbb{R}$ that $\|h\|_\infty < \infty$, when $p_1 \geq 10$, we have*

$$\begin{aligned} & \Pr \left\{ \max_{\mathbf{x}, \mathbf{z}} \left| \frac{1}{p_1} \sum_{j=1}^{p_1} h(\mathbf{V}[j])\mathbf{1}_{\{\mathbf{x}^T \mathbf{V}[j] > 0, \mathbf{z}^T \mathbf{V}[j] > 0\}} - \mathbb{E}_{\mathbf{v}}[h(\mathbf{v})\mathbf{1}_{\{\mathbf{x}^T \mathbf{v} > 0, \mathbf{z}^T \mathbf{v} > 0\}}] \right| \right. \\ & \quad \left. \leq 8\sqrt{\frac{\|h\|_\infty^2 2(d+1) \log(p_1+1)}{p_1}} \right\} \geq 1 - \frac{1}{(p_1+1)e^{d+1}}. \end{aligned}$$

By Eq. (1), we have

$$\begin{aligned} \frac{1}{p_1}(\mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V},\mathbf{z}}^{\text{RF}} &= \frac{1}{p_1} \sum_{j=1}^{p_1} (\mathbf{x}^T \mathbf{V}[j])(\mathbf{V}[j]^T \mathbf{z})\mathbf{1}_{\{\mathbf{x}^T \mathbf{V}[j] > 0, \mathbf{z}^T \mathbf{V}[j] > 0\}} \\ &= \mathbf{x}^T \left(\frac{1}{p_1} \sum_{j=1}^{p_1} (\mathbf{V}[j]\mathbf{V}[j]^T)\mathbf{1}_{\{\mathbf{x}^T \mathbf{V}[j] > 0, \mathbf{z}^T \mathbf{V}[j] > 0\}} \right) \mathbf{z}. \end{aligned}$$

Notice that $\mathbf{V}[j]\mathbf{V}[j]^T$ is a $d \times d$ matrix. Define

$$\mathbf{K}_j := (\mathbf{V}[j]\mathbf{V}[j]^T)\mathbf{1}_{\{\mathbf{x}^T \mathbf{V}[j] > 0, \mathbf{z}^T \mathbf{V}[j] > 0\}} \in \mathbb{R}^{d \times d}.$$

Thus, we have

$$\begin{aligned}
& \max_{\mathbf{x}, \mathbf{z}} \left| \frac{1}{p_1} (\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}} - K^{\text{RF}}(\mathbf{x}, \mathbf{z}) \right| \\
&= \max_{\mathbf{x}, \mathbf{z}} \left| \mathbf{x}^T \left(\frac{1}{p_1} \sum_{j=1}^{p_1} \mathbf{K}_j - \mathbb{E}_{\mathbf{v} \sim \lambda(\cdot)} (\mathbf{v} \mathbf{v}^T) \mathbf{1}_{\{\mathbf{x}^T \mathbf{v} > 0, \mathbf{z}^T \mathbf{v} > 0\}} \right) \mathbf{z} \right| \\
&\leq \max_{\mathbf{x}, \mathbf{z}} \left\| \mathbf{x}^T \right\|_2 \cdot \left\| \left(\frac{1}{p_1} \sum_{j=1}^{p_1} \mathbf{K}_j - \mathbb{E}_{\mathbf{v} \sim \lambda(\cdot)} (\mathbf{v} \mathbf{v}^T) \mathbf{1}_{\{\mathbf{x}^T \mathbf{v} > 0, \mathbf{z}^T \mathbf{v} > 0\}} \right) \right\|_2 \cdot \|\mathbf{z}\|_2 \quad (\text{by Lemma 12}) \\
&= \max_{\mathbf{x}, \mathbf{z}} \left\| \frac{1}{p_1} \sum_{j=1}^{p_1} \mathbf{K}_j - \mathbb{E}_{\mathbf{v} \sim \lambda(\cdot)} (\mathbf{v} \mathbf{v}^T) \mathbf{1}_{\{\mathbf{x}^T \mathbf{v} > 0, \mathbf{z}^T \mathbf{v} > 0\}} \right\|_2 \quad (\text{because } \|\mathbf{x}\|_2 = \|\mathbf{z}\|_2 = 1). \quad (33)
\end{aligned}$$

For any $k, l \in \{1, 2, \dots, d\}$, define the (k, l) -th element of \mathbf{K}_j as $K_{j, k, l}$. Thus, by Lemma 24 (notice that $|K_{j, k, l}| \leq 1$), we have

$$\begin{aligned}
& \Pr_{\mathbf{V}} \left\{ \max_{\mathbf{x}, \mathbf{z}} \left| \frac{1}{p_1} \sum_{j=1}^{p_1} K_{j, k, l} - \left(\mathbb{E}_{\mathbf{v} \sim \lambda(\cdot)} (\mathbf{v} \mathbf{v}^T) \mathbf{1}_{\{\mathbf{x}^T \mathbf{v} > 0, \mathbf{z}^T \mathbf{v} > 0\}} \right)_{k, l} \right| \leq 8 \sqrt{\frac{2(d+1) \log(p_1+1)}{p_1}} \right\} \\
&\geq 1 - \frac{1}{(p_1+1)e^{d+1}}.
\end{aligned}$$

Applying the union bound on all $d \times d$ elements of \mathbf{K}_j and by Lemma 13, we have

$$\begin{aligned}
& \Pr_{\mathbf{V}} \left\{ \max_{\mathbf{x}, \mathbf{z}} \left\| \frac{1}{p_1} \sum_{j=1}^{p_1} \mathbf{K}_j - \mathbb{E}_{\mathbf{v} \sim \lambda(\cdot)} (\mathbf{v} \mathbf{v}^T) \mathbf{1}_{\{\mathbf{x}^T \mathbf{v} > 0, \mathbf{z}^T \mathbf{v} > 0\}} \right\|_2 \leq 8d \sqrt{\frac{2(d+1) \log(p_1+1)}{p_1}} \right\} \\
&\geq 1 - \frac{d^2}{(p_1+1)e^{d+1}}.
\end{aligned}$$

Plugging it into Eq. (33), we thus have proven the following lemma.

Lemma 25. *Recall the definition of $Q(\cdot, \cdot)$ in Eq. (18). When $p_1 \geq 10$, we have*

$$\Pr_{\mathbf{V}} \left\{ \max_{\mathbf{x}, \mathbf{z}} \left| \frac{1}{p_1} (\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}} - K^{\text{RF}}(\mathbf{x}, \mathbf{z}) \right| \leq Q(p_1, d) \right\} \geq 1 - \frac{d^2}{(p_1+1)e^{d+1}}.$$

F.8 Some useful lemmas about multinomial expansion

Lemma 26 (Multinomial theorem (multinomial expansion)). *For any positive integer i and non-negative integer j ,*

$$(x_1 + x_2 + \dots + x_i)^j = \sum_{k_1 + k_2 + \dots + k_i = j} (k_1, k_2, \dots, k_i)! \cdot x_1^{k_1} x_2^{k_2} \dots x_i^{k_i},$$

where

$$(k_1, k_2, \dots, k_i)! = \frac{(k_1 + k_2 + \dots + k_i)!}{k_1! k_2! \dots k_i!}$$

denotes the multinomial coefficient.

Lemma 27. *We have*

$$\left(\sum_{i=0}^{\infty} a_i x^i \right)^j = \sum_{s=0}^{\infty} \left(\sum_{\substack{k_0 + k_1 + \dots + k_s = j \\ k_1 + 2k_2 + \dots + sk_s = s \\ k_0, k_1, \dots, k_s \in \mathbb{Z}_{\geq 0}}} (k_1, k_2, \dots, k_s)! \cdot a_0^{k_0} a_1^{k_1} \dots a_s^{k_s} \right) x^s.$$

Proof. The result directly follows from Lemma 26. Notice that $a_i x^i$ will not contribute to x^s when $i > s$. \square

G Proof of Proposition 4

Define

$$\Delta \mathbf{W}^*[k] := \int_{\mathcal{S}^{d-1}} \mathbf{1}_{\{(\mathbf{h}_{\mathbf{V},z}^{\text{RF}})^T \mathbf{W}_0[k] > 0\}} \mathbf{h}_{\mathbf{V},z}^{\text{RF}} \frac{g(z)}{p_1 p_2} d\mu(z), \quad k = 1, 2, \dots, p_2. \quad (34)$$

Notice that $\Delta \mathbf{W}^*[k]$ is a vector of size $p_1 \times 1$ (same as the size of $\mathbf{h}_{\mathbf{V},z}^{\text{RF}}$ and $\mathbf{h}_{\mathbf{V},x}^{\text{RF}}$). The connection between $\Delta \mathbf{W}^*$ and the pseudo ground-truth $f_{\mathbf{V},\mathbf{W}_0}^g$ is shown by the following lemma.

Lemma 28. *For all $\mathbf{x} \in \mathcal{S}^{d-1}$, we have*

$$\mathbf{h}_{\mathbf{V},\mathbf{W}_0,\mathbf{x}}^{\text{Three}} \cdot \Delta \mathbf{W}^* = f_{\mathbf{V},\mathbf{W}_0}^g(\mathbf{x}).$$

Proof. We have

$$\begin{aligned} & \mathbf{h}_{\mathbf{V},\mathbf{W}_0,\mathbf{x}}^{\text{Three}} \cdot \Delta \mathbf{W}^* \\ &= \sum_{k=1}^{p_2} (\mathbf{h}_{\mathbf{V},\mathbf{W}_0,\mathbf{x}}^{\text{Three}}[k])^T \Delta \mathbf{W}^*[k] \\ &= \sum_{k=1}^{p_2} \int_{\mathcal{S}^{d-1}} \mathbf{1}_{\{(\mathbf{h}_{\mathbf{V},x}^{\text{RF}})^T \mathbf{W}_0[k] > 0, (\mathbf{h}_{\mathbf{V},z}^{\text{RF}})^T \mathbf{W}_0[k] > 0\}} (\mathbf{h}_{\mathbf{V},x}^{\text{RF}})^T \mathbf{h}_{\mathbf{V},z}^{\text{RF}} \frac{g(z)}{p_1 p_2} d\mu(z) \\ & \quad (\text{by Eq. (2) and Eq. (34)}) \\ &= \int_{\mathcal{S}^{d-1}} \sum_{k=1}^{p_2} \frac{\mathbf{1}_{\{(\mathbf{h}_{\mathbf{V},x}^{\text{RF}})^T \mathbf{W}_0[k] > 0, (\mathbf{h}_{\mathbf{V},z}^{\text{RF}})^T \mathbf{W}_0[k] > 0\}}}{p_1 p_2} (\mathbf{h}_{\mathbf{V},x}^{\text{RF}})^T \mathbf{h}_{\mathbf{V},z}^{\text{RF}} g(z) d\mu(z) \\ &= \int_{\mathcal{S}^{d-1}} (\mathbf{h}_{\mathbf{V},z}^{\text{RF}})^T \mathbf{h}_{\mathbf{V},x}^{\text{RF}} \frac{|\mathcal{C}_{\mathbf{h}_{\mathbf{V},x}^{\text{RF}}, \mathbf{h}_{\mathbf{V},z}^{\text{RF}}}^{\mathbf{W}_0}|}{p_1 p_2} g(z) d\mu(z) \quad (\text{by Eq. (11)}) \\ &= f_{\mathbf{V},\mathbf{W}_0}^g(\mathbf{x}) \quad (\text{by Eq. (19)}). \end{aligned}$$

□

The following lemma bounds the test error for the pseudo ground-truth function with respect to the distance between $\Delta \mathbf{W}^*$ and the row-space of \mathbf{H} .

Lemma 29. *For all $\mathbf{a} \in \mathbb{R}^n$, we have*

$$|f_{\mathbf{V},\mathbf{W}_0}^g(\mathbf{x}) - \hat{f}^{\ell_2}(\mathbf{x})| \leq \sqrt{p_1 p_2} \|\Delta \mathbf{W}^* - \mathbf{H}^T \mathbf{a}\|_2.$$

Proof. Define $\mathbf{P} := \mathbf{H}^T (\mathbf{H}\mathbf{H}^T)^{-1} \mathbf{H}$. It is easy to verify that $\mathbf{P}^2 = \mathbf{P} = \mathbf{P}^T$, so \mathbf{P} is an orthogonal projection onto the space spanned by the rows of \mathbf{H} . By Lemma 28 and Eq. (3), when $\epsilon = \mathbf{0}$ and the ground-truth function is $f_{\mathbf{V},\mathbf{W}_0}^g$, we have $\mathbf{F}(\mathbf{X}) = \mathbf{H} \Delta \mathbf{W}^*$ and

$$\hat{f}^{\ell_2}(\mathbf{x}) = \mathbf{h}_{\mathbf{V},\mathbf{W}_0,\mathbf{x}}^{\text{Three}} \mathbf{H}^T (\mathbf{H}\mathbf{H}^T)^{-1} \mathbf{H} \Delta \mathbf{W}^* = \mathbf{h}_{\mathbf{V},\mathbf{W}_0,\mathbf{x}}^{\text{Three}} \mathbf{P} \Delta \mathbf{W}^*.$$

Thus, by Lemma 28, we have

$$|f_{\mathbf{V},\mathbf{W}_0}^g(\mathbf{x}) - \hat{f}^{\ell_2}(\mathbf{x})| = |\mathbf{h}_{\mathbf{V},\mathbf{W}_0,\mathbf{x}}^{\text{Three}} (\mathbf{P} - \mathbf{I}) \Delta \mathbf{W}^*|. \quad (35)$$

Because $\mathbf{P} = \mathbf{H}^T (\mathbf{H}\mathbf{H}^T)^{-1} \mathbf{H}$, we have

$$\mathbf{P}\mathbf{H}^T = \mathbf{H}^T (\mathbf{H}\mathbf{H}^T)^{-1} \mathbf{H}\mathbf{H}^T = \mathbf{H}^T. \quad (36)$$

We then have

$$\begin{aligned} \|\mathbf{P} \Delta \mathbf{W}^*\|_2 &= \|\mathbf{P} \Delta \mathbf{W}^* - \Delta \mathbf{W}^*\|_2 \\ &= \|\mathbf{P}(\mathbf{H}^T \mathbf{a} + \Delta \mathbf{W}^* - \mathbf{H}^T \mathbf{a}) - \Delta \mathbf{W}^*\|_2 \\ &= \|\mathbf{P}\mathbf{H}^T \mathbf{a} + \mathbf{P}(\Delta \mathbf{W}^* - \mathbf{H}^T \mathbf{a}) - \Delta \mathbf{W}^*\|_2 \\ &= \|\mathbf{H}^T \mathbf{a} + \mathbf{P}(\Delta \mathbf{W}^* - \mathbf{H}^T \mathbf{a}) - \Delta \mathbf{W}^*\|_2 \quad (\text{by Eq. (36)}) \\ &= \|(\mathbf{P} - \mathbf{I})(\Delta \mathbf{W}^* - \mathbf{H}^T \mathbf{a})\|_2 \\ &\leq \|\Delta \mathbf{W}^* - \mathbf{H}^T \mathbf{a}\|_2 \quad (\text{because } \mathbf{P} \text{ is an orthogonal projection}). \end{aligned} \quad (37)$$

Therefore, we have

$$\begin{aligned} \|\mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}}(\mathbf{P} - \mathbf{I})\Delta\mathbf{W}^*\| &= \|\mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}}(\mathbf{P} - \mathbf{I})\Delta\mathbf{W}^*\|_2 \\ &\leq \|\mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}}\|_2 \cdot \|(\mathbf{P} - \mathbf{I})\Delta\mathbf{W}^*\|_2 \text{ (by Lemma 12)} \\ &\leq \sqrt{p_1 p_2} \|\Delta\mathbf{W}^* - \mathbf{H}^T \mathbf{a}\|_2 \text{ (by Lemma 11 and Eq. (37)).} \end{aligned}$$

By Eq. (35), the result of this lemma thus follows. \square

Now we are ready to prove Proposition 4.

Define $\mathbf{K}_i \in \mathbb{R}^{(p_1 p_2) \times 1}$ (the same shape as \mathbf{W}_0) as

$$\mathbf{K}_i[k] := \mathbf{h}_{\mathbf{V}, \mathbf{X}_i}^{\text{RF}} \mathbf{1}_{\{(h_{\mathbf{V}, \mathbf{X}_i}^{\text{RF}})^T \mathbf{w}_0[k] > 0\}} \frac{g(\mathbf{X}_i)}{p_1 p_2}, \quad i \in \{1, 2, \dots, n\}, \quad k \in \{1, 2, \dots, p_2\}. \quad (38)$$

It is obvious that $\mathbf{K}_1, \mathbf{K}_2, \dots, \mathbf{K}_n$ are *i.i.d.* with respect to the randomness of \mathbf{X} . By Eq. (34), for all $k = 1, 2, \dots, p_2$, we have

$$\mathbb{E}_{\mathbf{X}_i} [\mathbf{K}_i[k]] = \Delta\mathbf{W}^*[k]. \quad (39)$$

Further, note that

$$\begin{aligned} \|\mathbf{K}_i[k]\|_2 &\leq \frac{\|g\|_\infty}{p_1 p_2} \|\mathbf{h}_{\mathbf{V}, \mathbf{X}_i}^{\text{RF}}\|_2 \text{ (by Lemma 12 and Eq. (38))} \\ &\leq \frac{\|g\|_\infty}{\sqrt{p_1 p_2}} \text{ (by Lemma 11)}. \end{aligned}$$

Thus, we have

$$\|\mathbf{K}_i\|_2 = \sqrt{\sum_{k=1}^{p_2} \|\mathbf{K}_i[k]\|_2^2} \leq \frac{\|g\|_\infty}{\sqrt{p_1 p_2}},$$

i.e.,

$$\sqrt{p_1 p_2} \|\mathbf{K}_i\|_2 \leq \|g\|_\infty. \quad (40)$$

We now construct the vector $\mathbf{a} \in \mathbb{R}^n$ that we will use in Lemma 29. Its i -th element is $\mathbf{a}_i = \frac{g(\mathbf{X}_i)}{n p_1 p_2}$, $i = 1, 2, \dots, n$. Then, for all $k \in \{1, 2, \dots, p_2\}$, we have

$$\begin{aligned} (\mathbf{H}^T \mathbf{a})[k] &= \sum_{i=1}^n \mathbf{H}_i^T[k] \mathbf{a}_i \\ &= \sum_{i=1}^n \mathbf{h}_{\mathbf{V}, \mathbf{X}_i}^{\text{RF}} \mathbf{1}_{\{(h_{\mathbf{V}, \mathbf{X}_i}^{\text{RF}})^T \mathbf{w}_0[k] > 0\}} \frac{g(\mathbf{X}_i)}{n p_1 p_2} \text{ (by Eq. (2))} \\ &= \frac{1}{n} \sum_{i=1}^n \mathbf{K}_i[k] \text{ (by Eq. (38))}, \end{aligned}$$

i.e.,

$$\mathbf{H}^T \mathbf{a} = \frac{1}{n} \sum_{i=1}^n \mathbf{K}_i. \quad (41)$$

Thus, by Lemma 14 (with $X_i = \sqrt{p_1 p_2} \mathbf{K}_i$, $U = \|g\|_\infty$, $m = q$), we have

$$\Pr_{\mathbf{X}} \left\{ \sqrt{p_1 p_2} \left\| \left(\frac{1}{n} \sum_{i=1}^n \mathbf{K}_i \right) - \mathbb{E}_{\mathbf{X}} \mathbf{K}_1 \right\|_2 \geq \frac{q \|g\|_\infty}{\sqrt{n}} \right\} \leq \frac{1}{q^2}.$$

Further, by Eq. (41) and Eq. (39), we have

$$\Pr_{\mathbf{X}} \left\{ \sqrt{p_1 p_2} \|\mathbf{H}^T \mathbf{a} - \Delta\mathbf{W}^*\|_2 \geq \frac{q \|g\|_\infty}{\sqrt{n}} \right\} \leq \frac{1}{q^2}.$$

By Lemma 29, we thus have

$$\Pr_{\mathbf{X}} \left\{ |f_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{x}) - \hat{f}^{\ell_2}(\mathbf{x})| \geq \frac{q \|g\|_\infty}{\sqrt{n}} \right\} \leq \frac{1}{q^2}.$$

The result of Proposition 4 thus follows.

H Proof of Proposition 6 (Minimum Eigenvalue of $\mathbf{H}\mathbf{H}^T$)

Define

$$\begin{aligned}\beta_{i,j} &:= \|\mathbf{h}_{\mathbf{V},\mathbf{X}_i}^{\text{RF}}\|_2 \cdot \|\mathbf{h}_{\mathbf{V},\mathbf{X}_j}^{\text{RF}}\|_2, \\ \theta_{i,j}^{\text{RF}} &:= \arccos\left(\frac{(\mathbf{h}_{\mathbf{V},\mathbf{X}_i}^{\text{RF}})^T \mathbf{h}_{\mathbf{V},\mathbf{X}_j}^{\text{RF}}}{\beta_{i,j}}\right) \in \left[0, \frac{\pi}{2}\right], \\ \theta_{\min}^{\text{RF}} &:= \min_{i \neq j} \theta_{i,j}^{\text{RF}}.\end{aligned}$$

(By Eq. (1), we know that every element of $\mathbf{h}_{\mathbf{V},\mathbf{X}_i}^{\text{RF}}$ and $\mathbf{h}_{\mathbf{V},\mathbf{X}_j}^{\text{RF}}$ are non-negative, and hence $\theta_{i,j}^{\text{RF}} \in [0, \frac{\pi}{2}]$.)

Define $\tilde{\mathbf{H}}^\infty \in \mathbb{R}^{n \times n}$ as

$$\tilde{\mathbf{H}}_{i,j}^\infty := \frac{p_1}{2d} \cos(\theta_{i,j}^{\text{RF}}) \cdot \frac{\pi - \theta_{i,j}^{\text{RF}}}{2\pi}. \quad (42)$$

The following lemma (restated) is from the proof of Lemma 1 of [Satpathi & Srikant \(2021\)](#), which relates $\min \text{eig}(\tilde{\mathbf{H}}^\infty)$ to $\theta_{\min}^{\text{RF}}$. For reader's convenience, we also provide its proof in Appendix H.1.

Lemma 30.

$$\min \text{eig}(\tilde{\mathbf{H}}^\infty) \geq \frac{1}{8\pi} \cdot \frac{p_1}{2d} \cdot \sqrt{\frac{\log(1/\cos \theta_{\min}^{\text{RF}})}{\log(2n/\cos \theta_{\min}^{\text{RF}})}}.$$

We then focus on estimating $\theta_{\min}^{\text{RF}}$.

Lemma 31. *Recall the definition of $C(n, d, q)$ in Eq. (16). For any $q > 0$, when p_1 is sufficient large such that*

$$\frac{10dnq\sqrt{2d}}{\sqrt{p_1}} \leq C(n, d, q). \quad (43)$$

we have

$$\Pr_{\mathbf{V}, \mathbf{X}} \left\{ \cos \theta_{\min}^{\text{RF}} \geq 1 - C(n, d, q) \right\} \leq \frac{4}{q^2}.$$

The proof of Lemma 31 is in Appendix H.2. Intuitively, when n becomes larger, some \mathbf{X}_i 's (together with $\mathbf{h}_{\mathbf{V},\mathbf{X}_i}^{\text{RF}}$'s) will get closer to each other, and thus $\theta_{\min}^{\text{RF}}$ will get closer to zero. Such intuition is captured by Lemma 31 since $C(n, d, q)$ is monotone decreasing with respect to n .

The above lemmas study the minimum eigenvalue of $\tilde{\mathbf{H}}^\infty$. We need to relate it to the minimum eigenvalue of $\mathbf{H}\mathbf{H}^T$, which is achieved by the following lemma.

Lemma 32. *For any $q > 0$,*

$$\Pr_{\mathbf{X}, \mathbf{V}, \mathbf{W}_0} \left\{ \left| \frac{1}{p_2} \min \text{eig}(\mathbf{H}\mathbf{H}^T) - \min \text{eig}(\tilde{\mathbf{H}}^\infty) \right| \geq qn^2 \sqrt{2p_1 d} + q^2 n^3 d + \frac{qn^2 p_1}{\sqrt{p_2}} \right\} \leq \frac{3}{q^2}.$$

The proof of Lemma 32 is in Appendix H.3. From the derivation in Appendix B, we know that each element of $\frac{\mathbf{H}\mathbf{H}^T}{p_1 p_2}$ will approach the corresponding element of $\frac{1}{p_1} \tilde{\mathbf{H}}^\infty$ as p_1 and p_2 get larger. Therefore, it is natural to expect that the minimum eigenvalue of those two matrices will also be closer to each other when p_1 and p_2 becomes larger, which is captured by Lemma 32.

Lemma 33. *For any $a \in (0, 1]$, we have $\log \frac{1}{a} \geq 1 - a$.*

Proof. Consider the function $h(a) := \log(1/a) - 1 + a$. We have $\frac{\partial h(a)}{\partial a} = -\frac{1}{a} + 1 \leq 0$. Thus, we know $h(a)$ is monotone decreasing in $a \in (0, 1]$. Thus, we have $h(a) \geq h(1) = 0$. The result of this lemma thus follows. \square

Now we are ready to prove Proposition 6.

Proof of Proposition 6. We define three events

$$\begin{aligned}\mathcal{J}_1 &:= \left\{ \cos \theta_{\min}^{\text{RF}} \geq 1 - C(n, d, q) \right\}, \\ \mathcal{J}_2 &:= \left\{ \left| \frac{1}{p_2} \min \text{eig}(\mathbf{H}\mathbf{H}^T) - \min \text{eig}(\tilde{\mathbf{H}}^\infty) \right| \geq qn^2\sqrt{2p_1d} + q^2n^3d + \frac{qn^2p_1}{\sqrt{p_2}} \right\}, \\ \mathcal{J}_3 &:= \left\{ \frac{1}{p_2} \min \text{eig}(\mathbf{H}\mathbf{H}^T) \leq p_1 J(n, p_1, p_2, d, q) \triangleq \frac{p_1}{16\pi d} \sqrt{\frac{C(n, d, q)}{\log(4n)}} \right. \\ &\quad \left. - \left(qn^2\sqrt{2p_1d} + q^2n^3d + \frac{qn^2p_1}{\sqrt{p_2}} \right) \right\}.\end{aligned}$$

Step 1: prove $\mathcal{J}_1 \cup \mathcal{J}_2 \supseteq \mathcal{J}_3$.

In order to prove $\mathcal{J}_1 \cup \mathcal{J}_2 \supseteq \mathcal{J}_3$, it is equivalent to prove $\mathcal{J}_1^c \cap \mathcal{J}_2^c \subseteq \mathcal{J}_3^c$. To that end, suppose \mathcal{J}_1^c and \mathcal{J}_2^c happen. Thus, we have

$$\begin{aligned}\log(1/\cos \theta_{\min}^{\text{RF}}) &\geq 1 - \cos \theta_{\min}^{\text{RF}} \quad (\text{by Lemma 33}) \\ &\geq C(n, d, q) \quad (\text{by the event } \mathcal{J}_1^c).\end{aligned}\tag{44}$$

Thus, we have

$$\begin{aligned}&\min \text{eig}(\tilde{\mathbf{H}}^\infty) \\ &\geq \frac{1}{8\pi} \cdot \frac{p_1}{2d} \sqrt{\frac{\log(1/\cos \theta_{\min}^{\text{RF}})}{\log(2n/\cos \theta_{\min}^{\text{RF}})}} \quad (\text{by Lemma 30}) \\ &= \frac{p_1}{16\pi d} \sqrt{\frac{\log(1/\cos \theta_{\min}^{\text{RF}})}{\log(2n) + \log(1/\cos \theta_{\min}^{\text{RF}})}} \\ &\geq \frac{p_1}{16\pi d} \sqrt{\frac{C(n, d, q)}{\log(2n) + C(n, d, q)}} \\ &\quad (\text{by Eq. (44) and } \frac{a}{\log(2n) + a} \text{ is monotone increasing with respect to } a) \\ &\geq \frac{p_1}{16\pi d} \sqrt{\frac{C(n, d, q)}{\log(4n)}} \quad (\text{since } \log(2) \approx 0.7, C(n, d, q) \leq \frac{\pi-1}{4\pi} \cdot \frac{1}{2} \leq \frac{1}{8} \leq \log 2).\end{aligned}$$

Thus, we have

$$\begin{aligned}\frac{1}{p_2} \min \text{eig}(\mathbf{H}\mathbf{H}^T) &\geq \min \text{eig}(\tilde{\mathbf{H}}^\infty) - \left| \frac{1}{p_2} \min \text{eig}(\mathbf{H}\mathbf{H}^T) - \min \text{eig}(\tilde{\mathbf{H}}^\infty) \right| \\ &\quad (\text{by the triangle inequality}) \\ &> \frac{p_1}{16\pi d} \sqrt{\frac{C(n, d, q)}{\log(4n)}} - \left(qn^2\sqrt{2p_1d} + q^2n^3d + \frac{qn^2p_1}{\sqrt{p_2}} \right) \quad (\text{by the event } \mathcal{J}_2^c) \\ &= p_1 J(n, p_1, p_2, d, q) \quad (\text{by Eq. (17)}),\end{aligned}$$

i.e., \mathcal{J}_3^c must then occur. Thus, we have shown that $\mathcal{J}_1^c \cap \mathcal{J}_2^c \subseteq \mathcal{J}_3^c$, which implies that $\mathcal{J}_1 \cup \mathcal{J}_2 \supseteq \mathcal{J}_3$.

Step 2: estimate \mathcal{J}_3

We have

$$\begin{aligned}\Pr_{\mathbf{x}, \mathbf{v}, \mathbf{w}_0} [\mathcal{J}_3] &\leq \Pr_{\mathbf{x}, \mathbf{v}, \mathbf{w}_0} [\mathcal{J}_1 \cup \mathcal{J}_2] \quad (\text{because } \mathcal{J}_1 \cup \mathcal{J}_2 \supseteq \mathcal{J}_3) \\ &\leq \Pr_{\mathbf{x}, \mathbf{v}, \mathbf{w}_0} [\mathcal{J}_1] + \Pr_{\mathbf{x}, \mathbf{v}, \mathbf{w}_0} [\mathcal{J}_2] \quad (\text{by the union bound}) \\ &= \Pr_{\mathbf{x}, \mathbf{v}} [\mathcal{J}_1] + \Pr_{\mathbf{x}, \mathbf{v}, \mathbf{w}_0} [\mathcal{J}_2] \quad (\text{as } \mathcal{J}_1 \text{ is independent of } \mathbf{w}_0) \\ &\leq \frac{7}{q^2} \quad (\text{by Lemma 31 and Lemma 32}).\end{aligned}$$

The result of Proposition 6 thus follows. \square

In the rest of this section, we prove Lemma 30, Lemma 31, and Lemma 32.

H.1 Proof of Lemma 30

Proof. For simplicity of notation, we define $\mathbf{a}_i \in \mathbb{R}^{p_1}$ as

$$\mathbf{a}_i := \frac{\mathbf{h}_{\mathbf{V}, \mathbf{X}_i}^{\text{RF}}}{\|\mathbf{h}_{\mathbf{V}, \mathbf{X}_i}^{\text{RF}}\|_2} \text{ for all } i = 1, 2, \dots, n.$$

Let $\mathbf{a}_i^{\otimes k} \in \mathbb{R}^{p_1 k}$ (a column vector with $p_1 k$ elements) denote the k -time Kronecker product of the vector \mathbf{a}_i with itself. We define

$$\begin{aligned} \mathbf{A} &:= [\mathbf{a}_1 \ \mathbf{a}_2 \ \dots \ \mathbf{a}_n] \in \mathbb{R}^{p_1 \times n}, \\ \mathbf{A}^{(k)} &:= [\mathbf{a}_1^{\otimes k} \ \mathbf{a}_2^{\otimes k} \ \dots \ \mathbf{a}_n^{\otimes k}] \in \mathbb{R}^{(p_1 k) \times n}, \\ \mathbf{B}^{(k)} &:= \left(\mathbf{A}^{(k)}\right)^T \mathbf{A}^{(k)}. \end{aligned}$$

Thus, we have

$$\cos \theta_{i,j}^{\text{RF}} = \mathbf{a}_i^T \mathbf{a}_j. \quad (45)$$

By the definition of Kronecker product, we thus have¹¹

$$\left(\mathbf{a}_i^T \mathbf{a}_j\right)^k = \left(\mathbf{a}_i^{\otimes k}\right)^T \left(\mathbf{a}_j^{\otimes k}\right). \quad (46)$$

Thus, by Lemma 17, we have

$$\begin{aligned} \cos(\theta_{i,j}^{\text{RF}}) \cdot \frac{\pi - \theta_{i,j}^{\text{RF}}}{2\pi} &= \frac{\cos \theta_{i,j}^{\text{RF}}}{4} + \frac{1}{2\pi} \sum_{k=0}^{\infty} \frac{(2k)!}{(k!)^2} \frac{4}{2k+1} \left(\frac{\cos \theta_{i,j}^{\text{RF}}}{2}\right)^{2k+2} \\ &= \frac{\mathbf{a}_i^T \mathbf{a}_j}{4} + \frac{1}{2\pi} \sum_{k=0}^{\infty} \frac{(2k)!}{(k!)^2} \frac{4}{2k+1} \left(\frac{\mathbf{a}_i^T \mathbf{a}_j}{2}\right)^{2k+2} \\ &= \frac{\mathbf{a}_i^T \mathbf{a}_j}{4} + \frac{1}{2\pi} \sum_{k=0}^{\infty} \frac{(2k)!}{(k!)^2} \frac{4}{2k+1} \left(\frac{1}{2}\right)^{2k+2} \left(\mathbf{a}_i^{\otimes 2k+2}\right)^T \mathbf{a}_j^{\otimes 2k+2}. \end{aligned}$$

Using Eq. (42), we then have

$$\tilde{\mathbf{H}}^\infty = \frac{p_1}{2d} \left(\frac{\mathbf{A}^T \mathbf{A}}{4} + \frac{1}{2\pi} \sum_{k=0}^{\infty} \frac{(2k)!}{(k!)^2} \frac{4}{2k+1} \left(\frac{1}{2}\right)^{2k+2} \mathbf{B}^{(2k+2)} \right).$$

Thus, we have

$$\begin{aligned} \min \text{eig}(\tilde{\mathbf{H}}^\infty) &= \min_{\mathbf{u}: \|\mathbf{u}\|_2=1} \mathbf{u}^T \tilde{\mathbf{H}}^\infty \mathbf{u} \\ &\geq \frac{p_1}{2d} \cdot \frac{1}{2\pi} \sum_{k=0}^{\infty} \frac{(2k)!}{(k!)^2} \frac{4}{2k+1} \left(\frac{1}{2}\right)^{2k+2} \min_{\mathbf{u}: \|\mathbf{u}\|_2=1} \mathbf{u}^T \mathbf{B}^{(2k+2)} \mathbf{u} \\ &= \frac{p_1}{2d} \cdot \frac{1}{2\pi} \sum_{k=0}^{\infty} \frac{(2k)!}{(k!)^2} \frac{4}{2k+1} \left(\frac{1}{2}\right)^{2k+2} \min \text{eig}(\mathbf{B}^{(2k+2)}). \end{aligned} \quad (47)$$

¹¹To help readers understand the correctness of Eq. (46), we give a toy example as follows. We have

$$\left(\begin{bmatrix} a & b \end{bmatrix} \begin{bmatrix} c \\ d \end{bmatrix} \right)^2 = (ac + bd)^2 = a^2 c^2 + 2abcd + b^2 d^2.$$

We also have

$$\begin{bmatrix} a \\ b \end{bmatrix}^{\otimes 2} = \begin{bmatrix} aa \\ ab \\ ba \\ bb \end{bmatrix}, \quad \begin{bmatrix} c \\ d \end{bmatrix}^{\otimes 2} = \begin{bmatrix} cc \\ cd \\ dc \\ dd \end{bmatrix} \implies \left(\begin{bmatrix} a \\ b \end{bmatrix}^{\otimes 2} \right)^T \left(\begin{bmatrix} c \\ d \end{bmatrix}^{\otimes 2} \right) = a^2 c^2 + 2abcd + b^2 d^2.$$

Thus, we have shown that $\left(\begin{bmatrix} a & b \end{bmatrix} \begin{bmatrix} c \\ d \end{bmatrix} \right)^2 = \left(\begin{bmatrix} a \\ b \end{bmatrix}^{\otimes 2} \right)^T \left(\begin{bmatrix} c \\ d \end{bmatrix}^{\otimes 2} \right)$.

Notice that all diagonal elements of $\mathbf{B}^{(2k+2)}$ equal to 1. Thus, by Gershgorin circle theorem (Bell, 1965), we have

$$\min \text{eig}(\mathbf{B}^{(2k+2)}) \geq 1 - \max_i \sum_{j \neq i} \mathbf{B}_{ij}^{(2k+2)}, \quad (48)$$

where $\mathbf{B}_{ij}^{(2k+2)}$ denotes the (i, j) -th element of $\mathbf{B}^{(2k+2)}$. Notice that

$$\begin{aligned} \max_i \sum_{j \neq i} \mathbf{B}_{ij}^{(2k+2)} &= \max_i \sum_{j \neq i} (\mathbf{a}_i^{\otimes 2k+2})^T (\mathbf{a}_j^{\otimes 2k+2}) \\ &= \max_i \sum_{j \neq i} (\cos \theta_{i,j}^{\text{RF}})^{2k+2} \quad (\text{by Eq. (46) and Eq. (45)}) \\ &\leq (n-1) (\cos \theta_{\min}^{\text{RF}})^{2k+2}. \end{aligned}$$

Note that, when $k \geq k^* := \frac{\log(2n-2)}{2 \log(1/\cos \theta_{\min}^{\text{RF}})} - 1$, we have

$$\begin{aligned} 2k+2 &\geq \frac{\log(2n-2)}{\log(1/\cos \theta_{\min}^{\text{RF}})} \\ \implies (2k+2) \log(1/\cos \theta_{\min}^{\text{RF}}) &\geq \log(2n-2) \\ \implies (n-1) (\cos \theta_{\min}^{\text{RF}})^{2k+2} &\leq \frac{1}{2}. \end{aligned}$$

Therefore, we have

$$\max_i \sum_{j \neq i} \mathbf{B}_{ij}^{(2k+2)} \leq \frac{1}{2}, \text{ for all } k \geq k^*.$$

By Eq. (47) and Eq. (48), we thus have

$$\begin{aligned} \min \text{eig}(\tilde{\mathbf{H}}^\infty) &\geq \frac{p_1}{8d\pi} \sum_{k \geq k^*} \frac{(2k)!}{(k!)^2} \frac{4}{2k+1} \left(\frac{1}{2}\right)^{2k+2} \\ &= \frac{p_1}{8d\pi} \sum_{k \geq \lceil k^* \rceil} \frac{(2k-1)!!}{(2k)!!} \frac{1}{2k+1} \\ &\geq \frac{p_1}{8d\pi} \sum_{k \geq \lceil k^* \rceil} \frac{1}{\sqrt{\pi} (k + \frac{4}{\pi} - 1)} \frac{1}{2k+1} \quad (\text{by Lemma 16}) \\ &\geq \frac{p_1}{8d\pi} \int_{k^*+1}^{\infty} \frac{1}{\sqrt{\pi} (x + \frac{4}{\pi} - 1)} \frac{1}{2x+1} dx \\ &\geq \frac{p_1}{8d\pi} \int_{k^*+1}^{\infty} \frac{1}{2\sqrt{\pi}} (x+1)^{-\frac{3}{2}} dx \quad (\text{notice that } \frac{4}{\pi} - 1 \approx 0.27 \leq 1) \\ &= \frac{p_1}{8d\pi} \frac{1}{\sqrt{\pi}} \frac{1}{\sqrt{k^*+2}}. \end{aligned}$$

Notice that $\frac{1}{\sqrt{\pi}} \geq \frac{1}{2}$ and

$$k^* + 2 = \frac{\log(2n-2)}{2 \log(1/\cos \theta_{\min}^{\text{RF}})} + 1 \leq \frac{\log(2n)}{\log(1/\cos \theta_{\min}^{\text{RF}})} + 1 = \frac{\log(2n/\cos \theta_{\min}^{\text{RF}})}{\log(1/\cos \theta_{\min}^{\text{RF}})}.$$

We thus have

$$\min \text{eig}(\tilde{\mathbf{H}}^\infty) \geq \frac{p_1}{16d\pi} \sqrt{\frac{\log(1/\cos \theta_{\min}^{\text{RF}})}{\log(2n/\cos \theta_{\min}^{\text{RF}})}}.$$

□

H.2 Proof of Lemma 31

We first show some useful lemmas.

Lemma 34. For any $\theta \in [0, \pi]$, we have

$$1 - \frac{\sin \theta + (\pi - \theta) \cos \theta}{\pi} \geq \frac{\pi - 1}{2\pi} \sin^2 \theta,$$

$$\lim_{\theta \rightarrow 0^+} \frac{1 - \frac{\sin \theta + (\pi - \theta) \cos \theta}{\pi}}{\sin^2 \theta} = \frac{1}{2}.$$

Proof. To prove the first part, we have

$$\begin{aligned} \frac{\sin \theta + (\pi - \theta) \cos \theta}{\pi} &\leq \frac{\sin \theta + (\pi - \theta) \sqrt{1 - \sin^2 \theta}}{\pi} \\ &\quad (\text{although } \cos \theta \text{ could be negative, we always have } \cos \theta \leq \sqrt{1 - \sin^2 \theta}) \\ &\leq \frac{\sin \theta + (\pi - \theta) \sqrt{1 - \sin^2 \theta + \frac{1}{4} \sin^4 \theta}}{\pi} \\ &= \frac{\sin \theta + (\pi - \theta) (1 - \frac{1}{2} \sin^2 \theta)}{\pi} \\ &\leq \frac{\sin \theta + (\pi - \sin \theta) (1 - \frac{1}{2} \sin^2 \theta)}{\pi} \quad (\text{because } \sin \theta \leq \theta) \\ &= 1 - \frac{\pi - \sin \theta}{2\pi} \sin^2 \theta \\ &\leq 1 - \frac{\pi - 1}{2\pi} \sin^2 \theta \quad (\text{because } \sin \theta \leq 1), \end{aligned}$$

i.e.,

$$1 - \frac{\sin \theta + (\pi - \theta) \cos \theta}{\pi} \geq \frac{\pi - 1}{2\pi} \sin^2 \theta.$$

To prove the second part, we have

$$\begin{aligned} \lim_{\theta \rightarrow 0^+} \frac{1 - \frac{\sin \theta + (\pi - \theta) \cos \theta}{\pi}}{\sin^2 \theta} &= \lim_{\theta \rightarrow 0^+} \frac{\frac{\partial}{\partial \theta} \left(1 - \frac{\sin \theta + (\pi - \theta) \cos \theta}{\pi} \right)}{\frac{\partial \sin^2 \theta}{\partial \theta}} \quad (\text{by L'Hospital's rule}) \\ &= \lim_{\theta \rightarrow 0^+} \frac{-\cos \theta + (\pi - \theta) \sin \theta + \cos \theta}{2\pi \sin \theta \cos \theta} \\ &= \lim_{\theta \rightarrow 0^+} \frac{\pi - \theta}{2\pi \cos \theta} \\ &= \frac{1}{2}. \end{aligned}$$

□

Lemma 35. Consider $a \geq 0$ and $b > 0$. Let $\delta := |b - 1|$. If $\delta \in [0, 0.5]$, we then have

$$a - a\delta \leq \frac{a}{b} \leq a + 2a\delta.$$

Therefore, for any $c \in \mathbb{R}$, we have

$$\left| \frac{a}{b} - c \right| \leq |a - c| + 2a\delta.$$

Further, if we know the upper bound of a , we have the following conclusion: (i) if $a \leq 1$, we must have $\frac{a}{b} \leq a + 2\delta$; (ii) if $a \leq 1.5$, we must have $\left| \frac{a}{b} - c \right| \leq |a - c| + 3a$.

Proof. We have

$$\begin{aligned}
\frac{a}{b} &\leq a \frac{1}{1-\delta} \quad (\text{because } b \geq 1 - |1-b| = 1-\delta) \\
&\leq a \frac{1+(1-2\delta)\delta}{1-\delta} \quad (\text{by } (1-2\delta) \geq 0 \text{ because } \delta \in [0, 0.5]) \\
&= a \frac{1+\delta-2\delta^2}{1-\delta} \\
&= a \frac{(1-\delta)(1+2\delta)}{1-\delta} \\
&= a + 2a\delta.
\end{aligned}$$

We also have

$$\begin{aligned}
\frac{a}{b} &\geq a \frac{1}{1+\delta} \quad (\text{because } b \leq 1 + |1-b| = 1+\delta) \\
&\geq a \frac{1-\delta^2}{1+\delta} \\
&= a - a\delta.
\end{aligned}$$

The result of this lemma thus follows. \square

Lemma 36. *If the condition in Eq. (43) is satisfied, then*

$$\frac{1 - 2C(n, d, q) + \frac{2dnq}{\sqrt{p_1}}}{1 - \frac{4dnq\sqrt{2d}}{\sqrt{p_1}} - \frac{4d^2n^2q^2}{p_1}} \leq 1 - C(n, d, q).$$

Proof. By Eq. (43) and the definition of $C(n, d, q)$ in Eq. (16), we have

$$\frac{10dnq\sqrt{2d}}{\sqrt{p_1}} \leq \frac{\pi-1}{4\pi} \cdot \frac{1}{2} \leq \frac{1}{8} \leq \frac{1}{2} \quad (49)$$

$$\Rightarrow \frac{dnq\sqrt{2d}}{\sqrt{p_1}} \leq \frac{1}{20}, \text{ and } \frac{d^2n^2q^2 \cdot 2d}{p_1} \leq \left(\frac{1}{20}\right)^2 = \frac{1}{400}$$

$$\Rightarrow \frac{4dnq\sqrt{2d}}{\sqrt{p_1}} + \frac{4d^2n^2q^2}{p_1} \leq \frac{1}{5} + \frac{1}{200d} \leq 0.5. \quad (50)$$

We also have

$$\begin{aligned}
&\frac{2dnq}{\sqrt{p_1}} + \frac{8dnq\sqrt{2d}}{\sqrt{p_1}} + \frac{8d^2n^2q^2}{p_1} \\
&\leq \frac{dnq\sqrt{2d}}{\sqrt{p_1}} + \frac{8dnq\sqrt{2d}}{\sqrt{p_1}} + \frac{1}{2} \sqrt{\frac{8d^2n^2q^2}{p_1}} \quad (\text{because } 2 \leq \sqrt{2d} \text{ and } \frac{8d^2n^2q^2}{p_1} \leq \frac{1}{100d} \leq \frac{1}{4}) \\
&= \frac{9dnq\sqrt{2d}}{\sqrt{p_1}} + \frac{dnq\sqrt{2}}{\sqrt{p_1}} \\
&\leq \frac{10dnq\sqrt{2d}}{\sqrt{p_1}} \\
&\leq C(n, d, q) \quad (\text{by Eq. (43)}). \quad (51)
\end{aligned}$$

Thus, we have

$$\begin{aligned}
\frac{2dnq}{\sqrt{p_1}} &\leq C(n, d, q) \quad (\text{by Eq. (43)}) \\
\Rightarrow 1 - 2C(n, d, q) + \frac{2dnq}{\sqrt{p_1}} &\in [0, 1] \quad (\text{since } C(n, d, q) \leq \frac{1}{8} \text{ by Eq. (49)}). \quad (52)
\end{aligned}$$

By Eq. (50), Eq. (52) and applying Lemma 35(i) (where $a = 1 - 2C(n, d, q) + \frac{2dnq}{\sqrt{p_1}}$, $b = 1 - \frac{4dnq\sqrt{2d}}{\sqrt{p_1}} - \frac{4d^2n^2q^2}{p_1}$, and $\delta = \frac{4dnq\sqrt{2d}}{\sqrt{p_1}} + \frac{4d^2n^2q^2}{p_1}$), we thus have

$$\begin{aligned} & \frac{1 - 2C(n, d, q) + \frac{2dnq}{\sqrt{p_1}}}{1 - \frac{4dnq\sqrt{2d}}{\sqrt{p_1}} - \frac{4d^2n^2q^2}{p_1}} \\ & \leq 1 - 2C(n, d, q) + \frac{2dnq}{\sqrt{p_1}} + \frac{8dnq\sqrt{2d}}{\sqrt{p_1}} + \frac{8d^2n^2q^2}{p_1} \\ & \leq 1 - C(n, d, q) \text{ (by Eq. (51)).} \end{aligned}$$

□

Lemma 37. Given \mathbf{X} , for any $m > 0$,

$$\Pr_{\mathbf{V}} \left\{ \left| \beta_{i,j} - \frac{p_1}{2d} \right| \geq 2m\sqrt{2p_1d} + 2m^2d \right\} \leq \frac{2}{m^2}.$$

In other words, given any $\mathbf{x}, \mathbf{z} \in \mathcal{S}^{d-1}$ and for any $q > 0$,

$$\Pr_{\mathbf{V}} \left\{ \left| \|\mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}}\|_2 \cdot \|\mathbf{h}_{\mathbf{V},\mathbf{z}}^{\text{RF}}\|_2 - \frac{p_1}{2d} \right| \geq 2m\sqrt{2p_1d} + 2m^2d \right\} \leq \frac{2}{m^2}.$$

Proof. Define

$$Q_k^i := (\mathbf{X}_i^T \mathbf{V}[k])(\mathbf{X}_i^T \mathbf{V}[k]) \mathbf{1}_{\{\mathbf{X}_i^T \mathbf{V}[k] > 0\}}.$$

By Eq. (1), we have

$$\frac{1}{p_1} \|\mathbf{h}_{\mathbf{V},\mathbf{x}_i}^{\text{RF}}\|_2^2 = \frac{1}{p_1} (\mathbf{h}_{\mathbf{V},\mathbf{x}_i}^{\text{RF}})^T \cdot \mathbf{h}_{\mathbf{V},\mathbf{x}_i}^{\text{RF}} = \frac{1}{p_1} \sum_{k=1}^{p_1} Q_k^i.$$

Note that

$$|Q_k^i| \leq \|\mathbf{X}_i\|_2 \cdot \|\mathbf{V}[k]\|_2 \cdot \|\mathbf{X}_i\|_2 \cdot \|\mathbf{V}[k]\|_2 = 1 \text{ (by Assumption 1).}$$

Further, note that

$$\begin{aligned} \mathbb{E}_{\mathbf{V}}[Q_1^i] &= \int_{\mathcal{S}^{d-1}} (\mathbf{X}_i^T \mathbf{v})(\mathbf{X}_i^T \mathbf{v}) \mathbf{1}_{\{\mathbf{X}_i^T \mathbf{v} > 0\}} d\lambda(\mathbf{v}) \text{ (by Eq. (1))} \\ &= \frac{\sin 0 + \pi \cos 0}{2d\pi} \text{ (by Lemma 21)} \\ &= \frac{1}{2d}. \end{aligned}$$

By Lemma 14, we thus have

$$\Pr_{\mathbf{V}} \left\{ \left| \frac{1}{p_1} \|\mathbf{h}_{\mathbf{V},\mathbf{x}_i}^{\text{RF}}\|_2^2 - \frac{1}{2d} \right| \geq \frac{m}{\sqrt{p_1}} \right\} \leq \frac{1}{m^2}. \quad (53)$$

Notice that

$$\left| \|\mathbf{h}_{\mathbf{V},\mathbf{x}_i}^{\text{RF}}\|_2 - \frac{p_1}{2d} \right| = \left(\|\mathbf{h}_{\mathbf{V},\mathbf{x}_i}^{\text{RF}}\|_2 + \sqrt{\frac{p_1}{2d}} \right) \cdot \left| \|\mathbf{h}_{\mathbf{V},\mathbf{x}_i}^{\text{RF}}\|_2 - \sqrt{\frac{p_1}{2d}} \right| \geq \sqrt{\frac{p_1}{2d}} \cdot \left| \|\mathbf{h}_{\mathbf{V},\mathbf{x}_i}^{\text{RF}}\|_2 - \sqrt{\frac{p_1}{2d}} \right|. \quad (54)$$

Combining Eq. (53) and Eq. (54), we then have

$$\Pr_{\mathbf{V}} \left\{ \left| \|\mathbf{h}_{\mathbf{V},\mathbf{x}_i}^{\text{RF}}\|_2 - \sqrt{\frac{p_1}{2d}} \right| \geq m\sqrt{2d} \right\} \leq \Pr_{\mathbf{V}} \left\{ \left| \|\mathbf{h}_{\mathbf{V},\mathbf{x}_i}^{\text{RF}}\|_2 - \frac{p_1}{2d} \right| \geq m\sqrt{p_1} \right\} \leq \frac{1}{m^2}.$$

Finally, notice that

$$\begin{aligned}
& \left| \beta_{i,j} - \frac{p_1}{2d} \right| \\
&= \left| \|\mathbf{h}_{\mathbf{V},\mathbf{x}_i}^{\text{RF}}\|_2 \left(\|\mathbf{h}_{\mathbf{V},\mathbf{x}_j}^{\text{RF}}\|_2 - \sqrt{\frac{p_1}{2d}} \right) + \|\mathbf{h}_{\mathbf{V},\mathbf{x}_j}^{\text{RF}}\|_2 \left(\|\mathbf{h}_{\mathbf{V},\mathbf{x}_i}^{\text{RF}}\|_2 - \sqrt{\frac{p_1}{2d}} \right) \right. \\
&\quad \left. - \left(\|\mathbf{h}_{\mathbf{V},\mathbf{x}_j}^{\text{RF}}\|_2 - \sqrt{\frac{p_1}{2d}} \right) \left(\|\mathbf{h}_{\mathbf{V},\mathbf{x}_i}^{\text{RF}}\|_2 - \sqrt{\frac{p_1}{2d}} \right) \right| \\
&\leq \sqrt{p_1} \left| \|\mathbf{h}_{\mathbf{V},\mathbf{x}_j}^{\text{RF}}\|_2 - \sqrt{\frac{p_1}{2d}} \right| + \sqrt{p_1} \left| \|\mathbf{h}_{\mathbf{V},\mathbf{x}_i}^{\text{RF}}\|_2 - \sqrt{\frac{p_1}{2d}} \right| \\
&\quad + \left| \|\mathbf{h}_{\mathbf{V},\mathbf{x}_j}^{\text{RF}}\|_2 - \sqrt{\frac{p_1}{2d}} \right| \cdot \left| \|\mathbf{h}_{\mathbf{V},\mathbf{x}_i}^{\text{RF}}\|_2 - \sqrt{\frac{p_1}{2d}} \right| \text{ (by the triangle inequality and Lemma 11)}.
\end{aligned}$$

Thus, we have

$$\begin{aligned}
& \left\{ \left| \|\mathbf{h}_{\mathbf{V},\mathbf{x}_i}^{\text{RF}}\|_2 - \sqrt{\frac{p_1}{2d}} \right| \geq m\sqrt{2d} \right\} \cup \left\{ \left| \|\mathbf{h}_{\mathbf{V},\mathbf{x}_j}^{\text{RF}}\|_2 - \sqrt{\frac{p_1}{2d}} \right| \geq m\sqrt{2d} \right\} \\
&\supseteq \left\{ \left| \beta_{i,j} - \frac{p_1}{2d} \right| \geq 2m\sqrt{2p_1d} + 2m^2d \right\}.
\end{aligned}$$

Applying the union bound, we thus have

$$\Pr_{\mathbf{V}} \left\{ \left| \beta_{i,j} - \frac{p_1}{2d} \right| \geq 2m\sqrt{2p_1d} + 2m^2d \right\} \leq \frac{2}{m^2}.$$

□

Now we are ready to prove Lemma 31.

Proof of Lemma 31. Define three events as

$$\begin{aligned}
\mathcal{J}_{1,i,j} &:= \left\{ \frac{2d}{p_1} (\mathbf{h}_{\mathbf{V},\mathbf{x}_i}^{\text{RF}})^T \cdot \mathbf{h}_{\mathbf{V},\mathbf{x}_j}^{\text{RF}} \geq 1 - 2C(n, d, q) + \frac{2dnq}{\sqrt{p_1}} \right\}, \\
\mathcal{J}_{2,i,j} &:= \left\{ \beta_{i,j} \leq \frac{p_1}{2d} - 2nq\sqrt{2p_1d} - 2n^2q^2d \right\}, \\
\mathcal{J}_{3,i,j} &:= \left\{ \cos \theta_{i,j}^{\text{RF}} \geq 1 - 2C(n, d, q) \right\}.
\end{aligned}$$

We take a few steps as follows to finish the proof.

Step 1: estimate $\mathcal{J}_{1,i,j}$.

Define

$$Q_k^{i,j} := (\mathbf{X}_i^T \mathbf{V}[k]) (\mathbf{X}_j^T \mathbf{V}[k]) \mathbf{1}_{\{\mathbf{x}_i^T \mathbf{V}[k] > 0, \mathbf{x}_j^T \mathbf{V}[k] > 0\}}, \quad k = 1, 2, \dots, p_1.$$

By Eq. (1) and the definition of $Q_k^{i,j}$, we have

$$(\mathbf{h}_{\mathbf{V},\mathbf{x}_i}^{\text{RF}})^T \cdot \mathbf{h}_{\mathbf{V},\mathbf{x}_j}^{\text{RF}} = \sum_{k=1}^{p_1} Q_k^{i,j}. \tag{55}$$

Note that

$$|Q_k^{i,j}| \leq \|\mathbf{X}_i\|_2 \cdot \|\mathbf{V}[k]\|_2 \cdot \|\mathbf{X}_j\|_2 \cdot \|\mathbf{V}[k]\|_2 = 1 \text{ (by Assumption 1 and Lemma 12)}.$$

By Lemma 14, we then have

$$\Pr_{\mathbf{V}} \left\{ \left| \frac{1}{p_1} \sum_{k=1}^{p_1} Q_k^{i,j} - \mathbb{E}_{\mathbf{V}}[Q_1^{i,j}] \right| \geq \frac{m}{\sqrt{p_1}} \right\} \leq \frac{1}{m^2}.$$

Let $\theta_{i,j} = \arccos(\mathbf{X}_i^T \mathbf{X}_j) \in [0, \pi]$ denote the angle between \mathbf{X}_i and \mathbf{X}_j , where $i \neq j$ and $i, j \in \{1, 2, \dots, n\}$. Notice that

$$\begin{aligned} \mathbb{E}_{\mathbf{V}}[Q_1^{i,j}] &= \int_{\mathcal{S}^{d-1}} (\mathbf{X}_i^T \mathbf{v})(\mathbf{X}_j^T \mathbf{v}) \mathbf{1}_{\{\mathbf{X}_i^T \mathbf{v} > 0, \mathbf{X}_j^T \mathbf{v} > 0\}} d\lambda(\mathbf{v}) \text{ (by Eq. (1))} \\ &= \frac{\sin \theta_{i,j} + (\pi - \theta_{i,j}) \cos \theta_{i,j}}{2d\pi} \text{ (by Lemma 21)} \\ &\leq \frac{1}{2d} \left(1 - \frac{\pi - 1}{2\pi} \sin^2 \theta_{i,j} \right) \text{ (by Lemma 34).} \end{aligned}$$

Thus, we have

$$\Pr_{\mathbf{V}} \left\{ \frac{1}{p_1} \sum_{k=1}^{p_1} Q_k^{i,j} \geq \frac{1}{2d} \left(1 - \frac{\pi - 1}{2\pi} \sin^2 \theta_{i,j} \right) + \frac{m}{\sqrt{p_1}} \right\} \leq \frac{1}{m^2}.$$

For any $\alpha \in [0, 1]$, we have

$$\begin{aligned} &\Pr_{\mathbf{X}} \{ \sin^2 \theta_{i,j} \leq \alpha \} \\ &= \Pr_{\mathbf{X}} \{ \theta_{i,j} \leq \arcsin(\sqrt{\alpha}) \text{ OR } \pi - \theta_{i,j} \leq \arcsin(\sqrt{\alpha}) \} \\ &\leq \Pr_{\mathbf{X}} \{ \theta_{i,j} \leq \arcsin(\sqrt{\alpha}) \} + \Pr_{\mathbf{X}} \{ \pi - \theta_{i,j} \leq \arcsin(\sqrt{\alpha}) \} \text{ (by the union bound)} \\ &= I_{\alpha} \left(\frac{d-1}{2}, \frac{1}{2} \right) \text{ (area of two caps, by Lemma 8 and Assumption 1)} \\ &\leq \frac{2\sqrt{d}\alpha^{\frac{d-1}{2}}}{(d-1)\sqrt{1-\alpha}} \text{ (by Lemma 9 and Lemma 10).} \end{aligned}$$

Further, because

$$\begin{aligned} &\{ \sin^2 \theta_{i,j} > \alpha \} \cap \left\{ \frac{1}{p_1} \sum_{k=1}^{p_1} Q_k^{i,j} < \frac{1}{2d} \left(1 - \frac{\pi - 1}{2\pi} \sin^2 \theta_{i,j} \right) + \frac{m}{\sqrt{p_1}} \right\} \\ &\subseteq \left\{ \frac{2d}{p_1} \sum_{k=1}^{p_1} Q_k^{i,j} < 1 - \frac{\pi - 1}{2\pi} \alpha + \frac{2dm}{\sqrt{p_1}} \right\}, \end{aligned}$$

we have

$$\begin{aligned} &\{ \sin^2 \theta_{i,j} \leq \alpha \} \cup \left\{ \frac{1}{p_1} \sum_{k=1}^{p_1} Q_k^{i,j} \geq \frac{1}{2d} \left(1 - \frac{\pi - 1}{2\pi} \sin^2 \theta_{i,j} \right) + \frac{m}{\sqrt{p_1}} \right\} \\ &\supseteq \left\{ \frac{2d}{p_1} \sum_{k=1}^{p_1} Q_k^{i,j} \geq 1 - \frac{\pi - 1}{2\pi} \alpha + \frac{2dm}{\sqrt{p_1}} \right\}. \end{aligned}$$

Thus, by the union bound and Eq. (55), we have

$$\Pr_{\mathbf{V}, \mathbf{X}} \left\{ \frac{2d}{p_1} (\mathbf{h}_{\mathbf{V}, \mathbf{X}_i}^{\text{RF}})^T \cdot \mathbf{h}_{\mathbf{V}, \mathbf{X}_j}^{\text{RF}} \geq 1 - \frac{\pi - 1}{2\pi} \alpha + \frac{2dm}{\sqrt{p_1}} \right\} \leq \frac{1}{m^2} + \frac{2\sqrt{d}\alpha^{\frac{d-1}{2}}}{(d-1)\sqrt{1-\alpha}}. \quad (56)$$

By letting

$$\alpha = \min \left\{ \frac{1}{2}, \left(\frac{(d-1)^2}{8d} \right)^{\frac{1}{d-1}} (qn)^{-\frac{4}{d-1}} \right\}, \text{ and } m = qn,$$

we have

$$\begin{aligned} \frac{2\sqrt{d}\alpha^{\frac{d-1}{2}}}{(d-1)\sqrt{1-\alpha}} &\leq \frac{2\sqrt{2}\sqrt{d}\alpha^{\frac{d-1}{2}}}{(d-1)} \text{ (because } \alpha \leq \frac{1}{2}) \\ &\leq \frac{2\sqrt{2}\sqrt{d}\sqrt{\frac{(d-1)^2}{8d} \frac{1}{q^2 n^2}}}{d-1} \text{ (because } \alpha \leq \left(\frac{(d-1)^2}{8d} \right)^{\frac{1}{d-1}} (qn)^{-\frac{4}{d-1}}) \\ &= \frac{1}{q^2 n^2}. \end{aligned}$$

Thus, by Eq. (56), we have

$$\Pr_{\mathbf{V}, \mathbf{X}}[\mathcal{J}_{1,i,j}] \leq \frac{2}{q^2 n^2}. \quad (57)$$

Step 2: estimate $\mathcal{J}_{2,i,j}$. By Lemma 37, we have

$$\Pr_{\mathbf{V}} \left\{ \beta_{i,j} \leq \frac{p_1}{2d} - 2m\sqrt{2p_1 d} - 2m^2 d \right\} \leq \frac{2}{m^2}.$$

Letting $m = qn$, we then have

$$\Pr_{\mathbf{V}}[\mathcal{J}_{2,i,j}] \leq \frac{2}{q^2 n^2}. \quad (58)$$

Step 3: prove $\mathcal{J}_{3,i,j} \subseteq \mathcal{J}_{1,i,j} \cup \mathcal{J}_{2,i,j}$.

In order to show $\mathcal{J}_{3,i,j} \subseteq \mathcal{J}_{1,i,j} \cup \mathcal{J}_{2,i,j}$, it suffices to show $\mathcal{J}_{3,i,j}^c \supseteq \mathcal{J}_{1,i,j}^c \cap \mathcal{J}_{2,i,j}^c$. When $\mathcal{J}_{1,i,j}^c \cap \mathcal{J}_{2,i,j}^c$ happens, we have

$$\begin{aligned} \frac{2d}{p_1} \beta_{i,j} &> 1 - \frac{2d}{p_1} \cdot 2nq\sqrt{2p_1 d} - \frac{2d}{p_1} \cdot 2n^2 q^2 d = 1 - \frac{4dnq\sqrt{2d}}{\sqrt{p_1}} - \frac{4d^2 n^2 q^2}{p_1}, \\ \frac{2d}{p_1} (\mathbf{h}_{\mathbf{V}, \mathbf{X}_i}^{\text{RF}})^T \cdot \mathbf{h}_{\mathbf{V}, \mathbf{X}_j}^{\text{RF}} &< 1 - 2C(n, d, q) + \frac{2dnq}{\sqrt{p_1}}. \end{aligned}$$

Thus, we have

$$\begin{aligned} \cos \theta_{i,j}^{\text{RF}} &= \frac{(\mathbf{h}_{\mathbf{V}, \mathbf{X}_i}^{\text{RF}})^T \cdot \mathbf{h}_{\mathbf{V}, \mathbf{X}_j}^{\text{RF}}}{\beta_{i,j}} \\ &< \frac{1 - 2C(n, d, q) + \frac{2dnq}{\sqrt{p_1}}}{1 - \frac{4dnq\sqrt{2d}}{\sqrt{p_1}} - \frac{4d^2 n^2 q^2}{p_1}} \\ &\leq 1 - C(n, d, q) \text{ (by Lemma 36)} \end{aligned}$$

i.e., the event $\mathcal{J}_{3,i,j}^c$ happens. To sum up, we have proven that $\mathcal{J}_{3,i,j}^c \supseteq \mathcal{J}_{1,i,j}^c \cap \mathcal{J}_{2,i,j}^c$, which implies $\mathcal{J}_{3,i,j} \subseteq \mathcal{J}_{1,i,j} \cup \mathcal{J}_{2,i,j}$.

Step 4: estimate $\mathcal{J}_{3,i,j}$. We have

$$\begin{aligned} \Pr_{\mathbf{V}, \mathbf{X}}[\mathcal{J}_{3,i,j}] &\leq \Pr_{\mathbf{V}, \mathbf{X}}[\mathcal{J}_{1,i,j}] + \Pr_{\mathbf{V}, \mathbf{X}}[\mathcal{J}_{2,i,j}] \text{ (by } \mathcal{J}_{3,i,j} \subseteq \mathcal{J}_{1,i,j} \cup \mathcal{J}_{2,i,j} \text{ and the union bound)} \\ &\leq \frac{4}{q^2 n^2} \text{ (by Eq. (57) and Eq. (58)).} \end{aligned} \quad (59)$$

Step 5: estimate $\cos \theta_{\min}^{\text{RF}}$. We have

$$\begin{aligned} &\Pr_{\mathbf{V}, \mathbf{X}} \left\{ \cos \theta_{\min}^{\text{RF}} \geq 1 - C(n, d, q) \right\} \\ &= \Pr_{\mathbf{V}, \mathbf{X}} \left[\bigcup_{i \neq j} \mathcal{J}_{3,i,j} \right] \\ &\leq n(n-1) \Pr_{\mathbf{V}, \mathbf{X}}[\mathcal{J}_{3,i,j}] \text{ (by the union bound)} \\ &\leq \frac{4}{q^2} \text{ (by Eq. (59)).} \end{aligned}$$

The result of Lemma 31 thus follows. \square

H.3 Proof of Lemma 32

We first introduce two useful lemmas. Define $\mathbf{H}^\infty \in \mathbb{R}^{n \times n}$ as

$$\mathbf{H}_{i,j}^\infty := \beta_{i,j} \cos(\theta_{i,j}^{\text{RF}}) \cdot \frac{\pi - \theta_{i,j}^{\text{RF}}}{2\pi}.$$

Lemma 38. Given \mathbf{X} and \mathbf{V} , for any $q > 0$, we have

$$\Pr_{\mathbf{W}_0} \left\{ \max_{i,j} \left| \frac{1}{p_2} (\mathbf{H}\mathbf{H}^T)_{i,j} - \mathbf{H}_{i,j}^\infty \right| \geq \frac{qnp_1}{\sqrt{p_2}} \right\} \leq \frac{1}{q^2}.$$

Thus, we also have

$$\Pr_{\mathbf{W}_0, \mathbf{X}, \mathbf{V}} \left\{ \max_{i,j} \left| \frac{1}{p_2} (\mathbf{H}\mathbf{H}^T)_{i,j} - \mathbf{H}_{i,j}^\infty \right| \geq \frac{qnp_1}{\sqrt{p_2}} \right\} \leq \frac{1}{q^2}.$$

Proof. For notation simplicity, given any $i, j \in \{1, 2, \dots, n\}$, we define

$$Q_k^{i,j} := (\mathbf{h}_{\mathbf{V}, \mathbf{X}_i}^{\text{RF}})^T \mathbf{h}_{\mathbf{V}, \mathbf{X}_j}^{\text{RF}} \mathbf{1}_{\{(\mathbf{h}_{\mathbf{V}, \mathbf{X}_i}^{\text{RF}})^T \mathbf{W}_0[k] > 0, (\mathbf{h}_{\mathbf{V}, \mathbf{X}_j}^{\text{RF}})^T \mathbf{W}_0[k]\}} \text{ for all } k \in \{1, 2, \dots, p_2\}.$$

By Eq. (2), we thus have

$$\begin{aligned} (\mathbf{H}\mathbf{H}^T)_{i,j} &= (\mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{X}_i}^{\text{Three}})^T \mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{X}_j}^{\text{Three}} = \sum_{k=1}^{p_2} (\mathbf{h}_{\mathbf{V}, \mathbf{X}_i}^{\text{RF}})^T \mathbf{h}_{\mathbf{V}, \mathbf{X}_j}^{\text{RF}} \mathbf{1}_{\{(\mathbf{h}_{\mathbf{V}, \mathbf{X}_i}^{\text{RF}})^T \mathbf{W}_0[k] > 0, (\mathbf{h}_{\mathbf{V}, \mathbf{X}_j}^{\text{RF}})^T \mathbf{W}_0[k]\}} \\ &= \sum_{k=1}^{p_2} Q_k^{i,j}. \end{aligned}$$

By Lemma 20 and recalling Eq. (42), we have

$$\mathbb{E}_{\mathbf{W}_0} [Q_k^{i,j}] = \mathbf{H}_{i,j}^\infty.$$

By Lemma 11 and Lemma 12, we have

$$|Q_k^{i,j}| \leq \|\mathbf{h}_{\mathbf{V}, \mathbf{X}_i}^{\text{RF}}\|_2 \cdot \|\mathbf{h}_{\mathbf{V}, \mathbf{X}_j}^{\text{RF}}\|_2 \leq p_1.$$

Note that $Q_k^{i,j}$ are independent across k . By Lemma 14, for any $m > 0$, we have

$$\Pr_{\mathbf{W}_0} \left\{ \left| \frac{1}{p_2} (\mathbf{H}\mathbf{H}^T)_{i,j} - \mathbf{H}_{i,j}^\infty \right| \geq m \frac{p_1}{\sqrt{p_2}} \right\} \leq \frac{1}{m^2}. \quad (60)$$

The result of this lemma thus follows by letting $m = qn$ and the union bound, i.e.,

$$\begin{aligned} \Pr_{\mathbf{W}_0} \left\{ \max_{i,j} \left| \frac{1}{p_2} (\mathbf{H}\mathbf{H}^T)_{i,j} - \mathbf{H}_{i,j}^\infty \right| \geq \frac{qnp_1}{\sqrt{p_2}} \right\} &= \Pr_{\mathbf{W}_0} \left\{ \bigcup_{i,j} \left\{ \left| \frac{1}{p_2} (\mathbf{H}\mathbf{H}^T)_{i,j} - \mathbf{H}_{i,j}^\infty \right| \geq \frac{qnp_1}{\sqrt{p_2}} \right\} \right\} \\ &\leq \sum_{i,j} \Pr_{\mathbf{W}_0} \left\{ \left| \frac{1}{p_2} (\mathbf{H}\mathbf{H}^T)_{i,j} - \mathbf{H}_{i,j}^\infty \right| \geq \frac{qnp_1}{\sqrt{p_2}} \right\} \\ &\quad \text{(by the union bound)} \\ &\leq \sum_{i,j} \frac{1}{q^2 n^2} \text{ (by letting } m = qn \text{ in Eq. (60))} \\ &= \frac{1}{q^2}. \end{aligned}$$

□

Lemma 39. Given \mathbf{X} , for any $q > 0$, we must have

$$\Pr_{\mathbf{V}} \left\{ \max_{i,j} \left| \mathbf{H}_{i,j}^\infty - \tilde{\mathbf{H}}_{i,j}^\infty \right| \geq qn\sqrt{2p_1 d} + q^2 n^2 d \right\} \leq \frac{2}{q^2}.$$

Thus, we also have

$$\Pr_{\mathbf{V}, \mathbf{X}, \mathbf{W}_0} \left\{ \max_{i,j} \left| \mathbf{H}_{i,j}^\infty - \tilde{\mathbf{H}}_{i,j}^\infty \right| \geq qn\sqrt{2p_1 d} + q^2 n^2 d \right\} \leq \frac{2}{q^2}.$$

Proof. We have

$$\begin{aligned}
\left| \mathbf{H}_{i,j}^\infty - \tilde{\mathbf{H}}_{i,j}^\infty \right| &= \left| \left(\beta_{i,j} - \frac{p_1}{2d} \right) \cos(\theta_{i,j}^{\text{RF}}) \cdot \frac{\pi - \theta_{i,j}^{\text{RF}}}{2\pi} \right| \quad (\text{by Eq. (42)}) \\
&\leq \left| \beta_{i,j} - \frac{p_1}{2d} \right| \cdot \left| \cos(\theta_{i,j}^{\text{RF}}) \right| \cdot \left| \frac{\pi - \theta_{i,j}^{\text{RF}}}{2\pi} \right| \quad (\text{by Lemma 12}) \\
&\leq \frac{1}{2} \left| \beta_{i,j} - \frac{p_1}{2d} \right| \quad (\text{since } 0 \leq \theta_{i,j}^{\text{RF}} \leq \frac{\pi}{2}).
\end{aligned}$$

The result of this lemma thus follows by letting $m = qn$ in Lemma 37 and the union bound, i.e.,

$$\begin{aligned}
&\Pr_{\mathbf{V}} \left\{ \max_{i,j} \left| \mathbf{H}_{i,j}^\infty - \tilde{\mathbf{H}}_{i,j}^\infty \right| \geq qn\sqrt{2p_1d} + q^2n^2d \right\} \\
&\leq \Pr_{\mathbf{V}} \left\{ \max_{i,j} \left| \beta_{i,j} - \frac{p_1}{2d} \right| \geq 2qn\sqrt{2p_1d} + 2q^2n^2d \right\} \\
&= \Pr_{\mathbf{V}} \left\{ \bigcup_{i,j} \left\{ \left| \beta_{i,j} - \frac{p_1}{2d} \right| \geq 2qn\sqrt{2p_1d} + 2q^2n^2d \right\} \right\} \\
&\leq \sum_{i,j} \Pr_{\mathbf{V}} \left\{ \left| \beta_{i,j} - \frac{p_1}{2d} \right| \geq 2qn\sqrt{2p_1d} + 2q^2n^2d \right\} \quad (\text{by the union bound}) \\
&\leq \sum_{i,j} \frac{2}{q^2n^2} \quad (\text{by letting } m = qn \text{ in Lemma 37}) \\
&= \frac{2}{q^2}.
\end{aligned}$$

□

Now we are ready to prove Lemma 32.

Proof of Lemma 32. By the triangle inequality, we have

$$\left| \frac{1}{p_2} (\mathbf{H}\mathbf{H}^T)_{i,j} - \tilde{\mathbf{H}}_{i,j}^\infty \right| \leq \left| \mathbf{H}_{i,j}^\infty - \tilde{\mathbf{H}}_{i,j}^\infty \right| + \left| \frac{1}{p_2} (\mathbf{H}\mathbf{H}^T)_{i,j} - \mathbf{H}_{i,j}^\infty \right|.$$

Thus, we have

$$\begin{aligned}
&\Pr_{\mathbf{x}, \mathbf{V}, \mathbf{W}_0} \left\{ \max_{i,j} \left| \frac{1}{p_2} (\mathbf{H}\mathbf{H}^T)_{i,j} - \tilde{\mathbf{H}}_{i,j}^\infty \right| \geq qn\sqrt{2p_1d} + q^2n^2d + \frac{qnp_1}{\sqrt{p_2}} \right\} \\
&\leq \Pr_{\mathbf{x}, \mathbf{V}, \mathbf{W}_0} \left\{ \left\{ \max_{i,j} \left| \mathbf{H}_{i,j}^\infty - \tilde{\mathbf{H}}_{i,j}^\infty \right| \geq qn\sqrt{2p_1d} + q^2n^2d \right\} \right. \\
&\quad \left. \cup \left\{ \max_{i,j} \left| \frac{1}{p_2} (\mathbf{H}\mathbf{H}^T)_{i,j} - \mathbf{H}_{i,j}^\infty \right| \geq \frac{qnp_1}{\sqrt{p_2}} \right\} \right\} \\
&\leq \Pr_{\mathbf{x}, \mathbf{V}, \mathbf{W}_0} \left\{ \max_{i,j} \left| \mathbf{H}_{i,j}^\infty - \tilde{\mathbf{H}}_{i,j}^\infty \right| \geq qn\sqrt{2p_1d} + q^2n^2d \right\} \\
&\quad + \Pr_{\mathbf{x}, \mathbf{V}, \mathbf{W}_0} \left\{ \max_{i,j} \left| \frac{1}{p_2} (\mathbf{H}\mathbf{H}^T)_{i,j} - \mathbf{H}_{i,j}^\infty \right| \geq \frac{qnp_1}{\sqrt{p_2}} \right\} \quad (\text{by the union bound}) \\
&\leq \frac{3}{q^2} \quad (\text{by Lemma 38 and Lemma 39}).
\end{aligned}$$

The result of Lemma 32 thus follows by Lemma 13 (where $k = n$).

□

I Proof of Proposition 5 and Proposition 7

We first provide some useful lemmas.

Lemma 40. For any $\varphi \in [0, 2\pi]$, we must have $\sin \varphi \leq \varphi$. For any $\varphi \in [0, \pi/2]$, we must have $\varphi \leq \frac{\pi}{2} \sin \varphi$.

Proof. See Lemma 41 of Ju et al. (2021). □

Lemma 41. For any $a_1, a_2 \in [-1, 1]$ that $|a_1 - a_2| \leq 1$, we must have

$$|\arccos(a_1) - \arccos(a_2)| \leq \frac{\sqrt{2}\pi}{2} \sqrt{|a_1 - a_2|}.$$

Proof. Without loss of generality, we assume $a_2 \geq a_1$ and let $\delta := a_2 - a_1 \in [0, 1]$. Because $\frac{\partial \arccos x}{\partial x} = -\frac{1}{\sqrt{1-x^2}}$, we have

$$\begin{aligned} \frac{\partial(\arccos(a_1) - \arccos(a_1 + \delta))}{\partial a_1} &= -\frac{1}{\sqrt{1-a_1^2}} + \frac{1}{\sqrt{1-(a_1 + \delta)^2}} \\ &\begin{cases} \leq 0, & \text{when } a_1 \in [-1, -\frac{\delta}{2}] \\ \geq 0, & \text{when } a_1 \in [-\frac{\delta}{2}, 1 - \delta] \end{cases}. \end{aligned}$$

Thus, we know the largest value of $\arccos(a_1) - \arccos(a_1 + \delta)$ can only be achieved at either $a_1 = -1$ or $a_1 = 1 - \delta$, i.e.,

$$\arccos(a_1) - \arccos(a_1 + \delta) \leq \max\{\pi - \arccos(-1 + \delta), \arccos(1 - \delta)\} = \arccos(1 - \delta). \quad (61)$$

(The last equality is because $\arccos(-x) = \pi - \arccos x$.) It remains to show that $\arccos(1 - \delta) \leq \frac{\sqrt{2}\pi}{2} \sqrt{\delta}$. To that end, it suffices to prove $\cos(\frac{\sqrt{2}\pi}{2} \sqrt{\delta}) \leq 1 - \delta$. Let $\theta := \frac{\sqrt{2}\pi}{2} \sqrt{\delta}$, i.e., $\delta = \frac{2}{\pi^2} \theta^2$. When $\theta > \frac{\pi}{2}$, we have $\cos(\frac{\sqrt{2}\pi}{2} \sqrt{\delta}) = \cos \theta < 0 < 1 - \delta$ (since $\delta \in [0, 1]$). When $\theta \in [0, \frac{\pi}{2}]$, we have

$$\begin{aligned} \cos(\frac{\sqrt{2}\pi}{2} \sqrt{\delta}) &= \cos \theta = \sqrt{1 - \sin^2 \theta} \leq \sqrt{1 - \sin^2 \theta + \frac{1}{4} \sin^4 \theta} = 1 - \frac{1}{2} \sin^2 \theta \\ &\leq 1 - \frac{1}{2} \left(\frac{2}{\pi} \theta\right)^2 \text{ (by Lemma 40)} \\ &= 1 - \delta. \end{aligned}$$

Therefore, we have proven that $\arccos(1 - \delta) \leq \frac{\sqrt{2}\pi}{2} \sqrt{\delta}$ for all $\delta \in [0, 1]$. By Eq. (61), the result of this lemma thus follows. □

Lemma 42. For any real number a_1, a_2, δ_1 , and δ_2 such that $a_2 \in [-1, 1]$, $a_2 + \delta_2 \in [-1, 1]$, and $|\delta_2| \leq 1$, we must have

$$\left| (a_1 + \delta_1) \frac{\pi - \arccos(a_2 + \delta_2)}{2\pi} - a_1 \frac{\pi - \arccos(a_2)}{2\pi} \right| \leq \frac{1}{2} |\delta_1| + \frac{\sqrt{2}|a_1| \sqrt{|\delta_2|}}{4}.$$

Proof. Define

$$b := a_1 \frac{\pi - \arccos(a_2 + \delta_2)}{2\pi}.$$

we have

$$\begin{aligned}
& \left| (a_1 + \delta_1) \frac{\pi - \arccos(a_2 + \delta_2)}{2\pi} - a_1 \frac{\pi - \arccos(a_2)}{2\pi} \right| \\
&= \left| (a_1 + \delta_1) \frac{\pi - \arccos(a_2 + \delta_2)}{2\pi} - b + b - a_1 \frac{\pi - \arccos(a_2)}{2\pi} \right| \\
&\leq \left| (a_1 + \delta_1) \frac{\pi - \arccos(a_2 + \delta_2)}{2\pi} - b \right| + \left| b - a_1 \frac{\pi - \arccos(a_2)}{2\pi} \right| \\
&= |\delta_1| \cdot \left| \frac{\pi - \arccos(a_2 + \delta_2)}{2\pi} \right| + |a_1| \cdot \left| \frac{\arccos(a_2 + \delta_2) - \arccos(a_2)}{2\pi} \right| \\
&\leq \frac{1}{2} |\delta_1| + \frac{\sqrt{2}|a_1|\sqrt{|\delta_2|}}{4} \quad (\text{since } \arccos(\cdot) \in [0, \pi] \text{ and by Lemma 41}).
\end{aligned}$$

□

Lemma 43. For any $\theta \in [0, \pi]$, we have

$$\frac{\sin \theta + (\pi - \theta) \cos \theta}{\pi} \in [0, 1].$$

Proof. We have

$$\frac{\partial(\sin \theta + (\pi - \theta) \cos \theta)}{\partial \theta} = -(\pi - \theta) \sin \theta \leq 0.$$

Thus, $\sin \theta + (\pi - \theta) \cos \theta$ is monotone decreasing. The result of this lemma thus follows by plugging $\theta = 0$ and $\theta = \pi$ into the expression. □

Lemma 44. Recall the definition of $K^{\text{Three}}(\cdot)$ in Eq. (6) and the definition of $Q(p_1, d)$ in Eq. (18). When p_1 is large enough such that $9d \cdot Q(p_1, d) \leq 1$, we must have

$$\begin{aligned}
& \Pr_{\mathbf{V}} \left\{ \max_{\mathbf{x}, \mathbf{z}} \left| \frac{1}{p_1} (\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}} \frac{\pi - \arccos \left(\frac{(\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}}{\|\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}}\|_2 \cdot \|\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}\|_2} \right)}{2\pi} - K^{\text{Three}}(\mathbf{x}^T \mathbf{z}) \right| \geq \sqrt{\frac{Q(p_1, d)}{d}} \right\} \\
&\leq \frac{d^2}{(p_1 + 1)e^{d+1}}.
\end{aligned}$$

Proof. Because $9d \cdot Q(p_1, d) \leq 1$, we have

$$Q(p_1, d) = \sqrt{Q(p_1, d)} \sqrt{Q(p_1, d)} \leq \sqrt{Q(p_1, d)} \sqrt{\frac{1}{9d}} = \sqrt{\frac{Q(p_1, d)}{9d}}. \quad (62)$$

We also have

$$2d \cdot Q(p_1, d) \leq \frac{2}{9} \leq 0.5. \quad (63)$$

Define two events

$$\begin{aligned}
\mathcal{J}_1 &:= \left\{ \max_{\mathbf{x}, \mathbf{z}} \left| \frac{1}{p_1} (\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}} - K^{\text{RF}}(\mathbf{x}^T \mathbf{z}) \right| \geq Q(p_1, d) \right\}, \\
\mathcal{J}_2 &:= \left\{ \max_{\mathbf{x}, \mathbf{z}} \left| a - K^{\text{Three}}(\mathbf{x}^T \mathbf{z}) \right| \geq \sqrt{\frac{Q(p_1, d)}{d}} \right\}.
\end{aligned}$$

Notice that the randomness of those events is on \mathbf{V} . We first show $\mathcal{J}_1 \supseteq \mathcal{J}_2$, i.e., $\mathcal{J}_1^c \subseteq \mathcal{J}_2^c$. To that end, suppose \mathcal{J}_1^c happens. Because of \mathcal{J}_1^c , we have

$$\max_{\mathbf{x}, \mathbf{z}} \left| \frac{2d}{p_1} (\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}} - 2d \cdot K^{\text{RF}}(\mathbf{x}^T \mathbf{z}) \right| \leq 2d \cdot Q(p_1, d). \quad (64)$$

By Eq. (4), we have

$$K^{\text{RF}}(\mathbf{x}^T \mathbf{x}) = K^{\text{RF}}(1) = \frac{1}{2d}. \quad (65)$$

Thus, we have

$$\begin{aligned} & \max_{\mathbf{x}, \mathbf{z}} \left| \frac{2d}{p_1} \|\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}\|_2 \cdot \|\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}}\|_2 - 1 \right| \\ &= \max_{\mathbf{x}} \left| \frac{2d}{p_1} \|\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}\|_2^2 - 1 \right| \quad (\text{the max value is achieved when } \|\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}\|_2 = \|\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}}\|_2) \\ &= \max_{\mathbf{x}} \left| \frac{2d}{p_1} \|\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}\|_2^2 - 2dK^{\text{RF}}(\mathbf{x}^T \mathbf{x}) \right| \quad (\text{by Eq. (65)}) \\ &\leq \max_{\mathbf{x}, \mathbf{z}} \left| \frac{2d}{p_1} \|\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}\|_2 \|\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}}\|_2 - 2dK^{\text{RF}}(\mathbf{x}^T \mathbf{z}) \right| \quad (\text{since we could set } \mathbf{z} = \mathbf{x} \text{ on the right hand side}) \\ &\leq 2d \cdot Q(p_1, d) \quad (\text{because of } \mathcal{J}_1^c). \end{aligned} \quad (66)$$

By Eq. (66), Eq. (64), and Eq. (63), we thus have

$$\left| \frac{2d}{p_1} \|\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}\|_2 \cdot \|\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}}\|_2 - 1 \right| \leq 0.5 \text{ for all } \mathbf{x} \text{ and } \mathbf{z}. \quad (67)$$

Thus, we then have $\frac{2d}{p_1} (\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}} \leq \frac{2d}{p_1} \|\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}\|_2 \cdot \|\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}}\|_2 \leq 1.5$. Besides, we have $\frac{2d}{p_1} (\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}} \geq 0$ because all elements of $\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}$ and $\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}}$ are non-negative by Eq. (1). In other words, we have

$$\frac{2d}{p_1} (\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}} \in [0, 1.5] \text{ for all } \mathbf{x} \text{ and } \mathbf{z}. \quad (68)$$

Therefore, we then have

$$\begin{aligned} & \max_{\mathbf{x}, \mathbf{z}} \left| \frac{(\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}}{\|\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}}\|_2 \cdot \|\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}\|_2} - 2d \cdot K^{\text{RF}}(\mathbf{x}, \mathbf{z}) \right| \\ &= \max_{\mathbf{x}, \mathbf{z}} \left| \frac{\frac{2d}{p_1} (\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}}{\frac{2d}{p_1} \|\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}}\|_2 \cdot \|\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}\|_2} - 2d \cdot K^{\text{RF}}(\mathbf{x}, \mathbf{z}) \right| \\ &\leq \max_{\mathbf{x}, \mathbf{z}} \left| \frac{2d}{p_1} (\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}} - 2d \cdot K^{\text{RF}}(\mathbf{x}, \mathbf{z}) \right| + 3 \max_{\mathbf{x}, \mathbf{z}} \left| \frac{2d}{p_1} \|\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}\|_2 \cdot \|\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}}\|_2 - 1 \right| \\ &\quad (\text{by Lemma 35(ii) where } a = \frac{2d}{p_1} (\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}} \in [0, 1.5] \text{ by Eq. (68),} \\ &\quad b = \frac{2d}{p_1} \|\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}\|_2 \cdot \|\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}}\|_2, \text{ and } \delta = \left| \frac{2d}{p_1} \|\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}\|_2 \cdot \|\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}}\|_2 - 1 \right| \in [0, 0.5] \text{ by Eq. (67).)} \\ &\leq 9d \cdot Q(p_1, d) \quad (\text{by Eq. (64) and Eq. (66)}). \end{aligned} \quad (69)$$

Now we apply Lemma 42 by letting $\delta_1 = \frac{1}{p_1} (\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}} - K^{\text{RF}}(\mathbf{x}^T \mathbf{z})$, $\delta_2 = \frac{(\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}}{\|\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}}\|_2 \cdot \|\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}\|_2} - 2d \cdot K^{\text{RF}}(\mathbf{x}^T \mathbf{z})$, $a_1 = K^{\text{RF}}(\mathbf{x}^T \mathbf{z})$, and $a_2 = 2d \cdot K^{\text{RF}}(\mathbf{x}^T \mathbf{z})$. We first check the conditions required by Lemma 42. By Eq. (4) and Lemma 43, we have

$$a_2 = 2d \cdot K^{\text{RF}}(\mathbf{x}^T \mathbf{z}) \in [0, 1] \subseteq [-1, 1].$$

Because $|(\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}| \leq \|\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}}\|_2 \cdot \|\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}\|_2$, we have

$$a_2 + \delta_2 = \frac{(\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}}{\|\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}}\|_2 \cdot \|\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}\|_2} \in [-1, 1].$$

By Eq. (69) and $9d \cdot Q(p_1, d) \leq 1$ (the condition of this lemma), we have

$$|\delta_2| \leq 9d \cdot Q(p_1, d) \leq 1.$$

Therefore, all conditions of Lemma 42 are satisfied. According to Lemma 42, we then have

$$\begin{aligned}
& \left| (a_1 + \delta_1) \frac{\pi - \arccos(a_2 + \delta_2)}{2\pi} - a_1 \frac{\pi - \arccos(a_2)}{2\pi} \right| \leq \frac{1}{2} |\delta_1| + \frac{\sqrt{2} |a_1| \sqrt{|\delta_2|}}{4} \\
\Rightarrow & \left| \frac{1}{p_1} (\mathbf{h}_{\mathbf{V},z}^{\text{RF}})^T \mathbf{h}_{\mathbf{V},x}^{\text{RF}} \frac{\pi - \arccos\left(\frac{(\mathbf{h}_{\mathbf{V},z}^{\text{RF}})^T \mathbf{h}_{\mathbf{V},x}^{\text{RF}}}{\|\mathbf{h}_{\mathbf{V},z}^{\text{RF}}\|_2 \cdot \|\mathbf{h}_{\mathbf{V},x}^{\text{RF}}\|_2}\right)}{2\pi} \right. \\
& \left. - K^{\text{RF}}(\mathbf{x}^T \mathbf{z}) \frac{\pi - \arccos(2d \cdot K^{\text{RF}}(\mathbf{x}^T \mathbf{z}))}{2\pi} \right| \leq \frac{1}{2} \left| \frac{1}{p_1} (\mathbf{h}_{\mathbf{V},x}^{\text{RF}})^T \mathbf{h}_{\mathbf{V},z}^{\text{RF}} - K^{\text{RF}}(\mathbf{x}^T \mathbf{z}) \right| \\
& + \frac{\sqrt{2} |K^{\text{RF}}(\mathbf{x}^T \mathbf{z})| \sqrt{\left| \frac{(\mathbf{h}_{\mathbf{V},z}^{\text{RF}})^T \mathbf{h}_{\mathbf{V},x}^{\text{RF}}}{\|\mathbf{h}_{\mathbf{V},z}^{\text{RF}}\|_2 \cdot \|\mathbf{h}_{\mathbf{V},x}^{\text{RF}}\|_2} - 2d \cdot K^{\text{RF}}(\mathbf{x}^T \mathbf{z}) \right|}}{4}.
\end{aligned}$$

By \mathcal{J}_1^c and Eq. (69), we thus have

$$\begin{aligned}
& \max_{\mathbf{x}, \mathbf{z}} \left| \frac{1}{p_1} (\mathbf{h}_{\mathbf{V},z}^{\text{RF}})^T \mathbf{h}_{\mathbf{V},x}^{\text{RF}} \frac{\pi - \arccos\left(\frac{(\mathbf{h}_{\mathbf{V},z}^{\text{RF}})^T \mathbf{h}_{\mathbf{V},x}^{\text{RF}}}{\|\mathbf{h}_{\mathbf{V},z}^{\text{RF}}\|_2 \cdot \|\mathbf{h}_{\mathbf{V},x}^{\text{RF}}\|_2}\right)}{2\pi} - K^{\text{Three}}(\mathbf{x}^T \mathbf{z}) \right| \\
& \leq \frac{Q(p_1, d)}{2} + \frac{\sqrt{2} |K^{\text{RF}}(\mathbf{x}, \mathbf{z})| \sqrt{9d \cdot Q(p_1, d)}}{4} \\
& \leq \frac{Q(p_1, d)}{2} + \sqrt{\frac{9}{32d}} \cdot Q(p_1, d) \text{ (because } K^{\text{RF}}(\mathbf{x}, \mathbf{z}) \in \left[0, \frac{1}{2d}\right] \text{ by Lemma 43)} \\
& \leq \left(\sqrt{\frac{1}{36d}} + \sqrt{\frac{9}{32d}} \right) \sqrt{Q(p_1, d)} \text{ (by Eq. (62))} \\
& \leq \sqrt{\frac{Q(p_1, d)}{d}} \text{ (since } \sqrt{\frac{1}{36}} + \sqrt{\frac{9}{32}} \approx \frac{1}{6} + 0.53 \leq 1),
\end{aligned}$$

i.e., \mathcal{J}_2^c happens. We next estimate the probability of \mathcal{J}_2 . We have

$$\begin{aligned}
\Pr_{\mathbf{V}}[\mathcal{J}_2] & \leq \Pr[\mathcal{J}_1] \text{ (because } \mathcal{J}_2 \subseteq \mathcal{J}_1) \\
& \leq \frac{d^2}{(p_1 + 1)e^{d+1}} \text{ (by Lemma 25, noticing that } 9d \cdot Q(p_1, d) \leq 1 \implies p_1 \geq 10).
\end{aligned}$$

The result of this lemma thus follows. \square

Lemma 45. *We have*

$$\|\mathbf{H}^T (\mathbf{H}\mathbf{H}^T)^{-1}\|_2 \leq \frac{1}{\sqrt{\min \text{eig}(\mathbf{H}\mathbf{H}^T)}}.$$

Proof. For any $\mathbf{a} \in \mathbb{R}^n$, we have

$$\begin{aligned}
\|\mathbf{H}^T (\mathbf{H}\mathbf{H}^T)^{-1} \mathbf{a}\|_2 & = \sqrt{(\mathbf{H}^T (\mathbf{H}\mathbf{H}^T)^{-1} \mathbf{a})^T \mathbf{H}^T (\mathbf{H}\mathbf{H}^T)^{-1} \mathbf{a}} = \sqrt{\mathbf{a}^T (\mathbf{H}\mathbf{H}^T)^{-1} \mathbf{a}} \\
& \leq \frac{\|\mathbf{a}\|_2}{\sqrt{\min \text{eig}(\mathbf{H}\mathbf{H}^T)}}.
\end{aligned}$$

The result of this lemma thus follows. \square

We are now ready to prove Proposition 5 and Proposition 7.

I.1 Proof of Proposition 5

Proof. For $k = 1, 2, \dots, p_2$, define

$$K_k = \int_{S^{d-1}} \frac{(\mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V},\mathbf{z}}^{\text{RF}}}{p_1} \mathbf{1}_{\{(\mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}})^T \mathbf{W}_0[k] > 0, (\mathbf{h}_{\mathbf{V},\mathbf{z}}^{\text{RF}})^T \mathbf{W}_0[k]\}} g(\mathbf{z}) d\mu(\mathbf{z}). \quad (70)$$

It is obvious that K_1, K_2, \dots, K_{p_2} are *i.i.d.* (when randomness is on \mathbf{W}_0). By Eq. (19) and Eq. (2), we have

$$f_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{x}) = \frac{1}{p_2} \sum_{k=1}^{p_2} K_k. \quad (71)$$

Notice that

$$\begin{aligned} |K_k| &\leq \int_{S^{d-1}} \left| \frac{(\mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V},\mathbf{z}}^{\text{RF}}}{p_1} \right| \cdot |g(\mathbf{z})| d\mu(\mathbf{z}) \\ &\leq \int_{S^{d-1}} |g(\mathbf{z})| d\mu(\mathbf{z}) \text{ (by Lemma 11)} \\ &= \|g\|_1. \end{aligned} \quad (72)$$

Thus, by Lemma 14, we have

$$\Pr_{\mathbf{W}_0} \left\{ \left| \frac{1}{p_2} \sum_{k=1}^{p_2} K_k - \mathbb{E}_{\mathbf{W}_0} [K_1] \right| \geq \frac{q \|g\|_1}{\sqrt{p_2}} \right\} \leq \frac{1}{q^2}. \quad (73)$$

For any $k \in \{1, 2, \dots, p_2\}$, we have

$$\begin{aligned} &\mathbb{E}_{\mathbf{W}_0} [K_k] \\ &= \int_{S^{d-1}} \mathbb{E}_{\mathbf{W}_0} \left[\frac{(\mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V},\mathbf{z}}^{\text{RF}}}{p_1} \mathbf{1}_{\{(\mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}})^T \mathbf{W}_0[k] > 0, (\mathbf{h}_{\mathbf{V},\mathbf{z}}^{\text{RF}})^T \mathbf{W}_0[k]\}} g(\mathbf{z}) d\mu(\mathbf{z}) \right] \\ &= \int_{S^{d-1}} \frac{(\mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V},\mathbf{z}}^{\text{RF}}}{p_1} \cdot \frac{\pi - \arccos \left(\frac{(\mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V},\mathbf{z}}^{\text{RF}}}{\|\mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}}\|_2 \cdot \|\mathbf{h}_{\mathbf{V},\mathbf{z}}^{\text{RF}}\|_2} \right)}{2\pi} g(\mathbf{z}) d\mu(\mathbf{z}) \text{ (by Lemma 20)} \\ &= f(\mathbf{x}) + \int_{S^{d-1}} \left(\frac{(\mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V},\mathbf{z}}^{\text{RF}}}{p_1} \cdot \frac{\pi - \arccos \left(\frac{(\mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V},\mathbf{z}}^{\text{RF}}}{\|\mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}}\|_2 \cdot \|\mathbf{h}_{\mathbf{V},\mathbf{z}}^{\text{RF}}\|_2} \right)}{2\pi} - K^{\text{Three}}(\mathbf{x}^T \mathbf{z}) \right) g(\mathbf{z}) d\mu(\mathbf{z}) \\ &\text{(by } f = f_g \text{ and Eq. (7)).} \end{aligned}$$

Thus, we have

$$\begin{aligned} &\left| \mathbb{E}_{\mathbf{W}_0} [K_k] - f(\mathbf{x}) \right| \\ &\leq \max_{\mathbf{x}, \mathbf{z}} \left| \frac{(\mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V},\mathbf{z}}^{\text{RF}}}{p_1} \cdot \frac{\pi - \arccos \left(\frac{(\mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V},\mathbf{z}}^{\text{RF}}}{\|\mathbf{h}_{\mathbf{V},\mathbf{x}}^{\text{RF}}\|_2 \cdot \|\mathbf{h}_{\mathbf{V},\mathbf{z}}^{\text{RF}}\|_2} \right)}{2\pi} - K^{\text{Three}}(\mathbf{x}^T \mathbf{z}) \right| \cdot \|g\|_1. \end{aligned} \quad (74)$$

Applying Lemma 44, we then have

$$\Pr_{\mathbf{V}} \left\{ \left| \mathbb{E}_{\mathbf{W}_0} [K_k] - f(\mathbf{x}) \right| \geq \sqrt{\frac{Q(p_1, d)}{d}} \|g\|_1 \right\} \leq \frac{d^2}{(p_1 + 1)e^{d+1}}. \quad (75)$$

Notice that

$$\begin{aligned} \left| f_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{x}) - f(\mathbf{x}) \right| &= \left| \frac{1}{p_2} \sum_{k=1}^{p_2} K_k - f(\mathbf{x}) \right| \quad (\text{by Eq. (71)}) \\ &\leq \left| \frac{1}{p_2} \sum_{k=1}^{p_2} K_k - \mathbb{E}_{\mathbf{W}_0} [K_1] \right| + \left| \mathbb{E}_{\mathbf{W}_0} [K_1] - f(\mathbf{x}) \right| \quad (\text{by the triangle inequality}). \end{aligned}$$

Combining Eq. (73) and Eq. (75) by the union bound, we thus have

$$\begin{aligned} &\Pr_{\mathbf{V}, \mathbf{W}_0} \left\{ \left| f_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{x}) - f(\mathbf{x}) \right| \geq \frac{q\|g\|_1}{\sqrt{p_2}} + \sqrt{\frac{Q(p_1, d)}{d}} \|g\|_1 \right\} \\ &\leq \Pr_{\mathbf{V}, \mathbf{W}_0} \left\{ \left| \frac{1}{p_2} \sum_{k=1}^{p_2} K_k - \mathbb{E}_{\mathbf{W}_0} [K_1] \right| \geq \frac{q\|g\|_1}{\sqrt{p_2}} \right\} + \Pr_{\mathbf{V}, \mathbf{W}_0} \left\{ \left| \mathbb{E}_{\mathbf{W}_0} [K_k] - f(\mathbf{x}) \right| \geq \sqrt{\frac{Q(p_1, d)}{d}} \|g\|_1 \right\} \\ &\leq \frac{d^2}{(p_1 + 1)e^{d+1}} + \frac{1}{q^2}. \end{aligned} \quad (76)$$

□

I.2 Proof of Proposition 7

Proof. For $k = 1, 2, \dots, p_2$, define $\mathbf{K}_k \in \mathbb{R}^n$ whose i -th element is

$$\mathbf{K}_{k,i} := \int_{S^{d-1}} \frac{(\mathbf{h}_{\mathbf{V}, \mathbf{X}_i}^{\text{RF}})^T \mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}}}{p_1} \mathbf{1}_{\{(\mathbf{h}_{\mathbf{V}, \mathbf{X}_i}^{\text{RF}})^T \mathbf{w}_0[k] > 0, (\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}})^T \mathbf{w}_0[k]\}} g(\mathbf{z}) d\mu(\mathbf{z}).$$

Note that $\mathbf{K}_{k,i}$ is similar to K_k in Eq. (70), with the only difference that the former is defined with respect to \mathbf{X}_i and the latter is defined with respect to \mathbf{x} . Thus, we use a similar strategy to work with $\mathbf{K}_{k,i}$. By Eq. (20) and Eq. (19), we have

$$\mathbf{F}_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{X}) = \frac{1}{p_2} \sum_{k=1}^{p_2} \mathbf{K}_k.$$

Similar to Eq. (72), we have

$$|\mathbf{K}_{k,i}| \leq \|g\|_1, \text{ for all } i = 1, 2, \dots, n.$$

Thus, we have

$$\|\mathbf{K}_k\|_2 = \sqrt{\sum_{i=1}^n \mathbf{K}_{k,i}^2} \leq \sqrt{n} \|g\|_1.$$

By Lemma 14, we thus have

$$\Pr_{\mathbf{W}_0} \left\{ \left\| \frac{1}{p_2} \sum_{k=1}^{p_2} \mathbf{K}_k - \mathbb{E}_{\mathbf{W}_0} [\mathbf{K}_1] \right\|_2 \geq \frac{q\sqrt{n}\|g\|_1}{\sqrt{p_2}} \right\} \leq \frac{1}{q^2}.$$

Similar to Eq. (74), we have

$$\begin{aligned} &\left\| \mathbb{E}_{\mathbf{W}_0} [\mathbf{K}_1] - \mathbf{F}(\mathbf{X}) \right\|_2 \\ &\leq \sqrt{n} \max_{\mathbf{x}, \mathbf{z}} \left| \frac{(\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}}}{p_1} \cdot \frac{\pi - \arccos \left(\frac{(\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}})^T \mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}}}{\|\mathbf{h}_{\mathbf{V}, \mathbf{x}}^{\text{RF}}\|_2 \cdot \|\mathbf{h}_{\mathbf{V}, \mathbf{z}}^{\text{RF}}\|_2} \right)}{2\pi} - K^{\text{Three}}(\mathbf{x}, \mathbf{z}) \right| \cdot \|g\|_1. \end{aligned}$$

Thus, similar to Eq. (76), we have

$$\Pr_{\mathbf{V}, \mathbf{W}_0} \left\{ \left\| \mathbf{F}(\mathbf{X}) - \mathbf{F}_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{X}) \right\|_2 \geq \frac{q\sqrt{n}\|g\|_1}{\sqrt{p_2}} + \sqrt{\frac{Q(p_1, d)}{d}} \sqrt{n}\|g\|_1 \right\} \leq \frac{d^2}{(p_1 + 1)e^{d+1}} + \frac{1}{q^2}. \quad (77)$$

We note that

$$\begin{aligned}
& \text{term D} + \text{term B of Eq. (22)} \\
& \leq \| \mathbf{h}_{\mathbf{V}, \mathbf{W}_0, \mathbf{x}}^{\text{Three}} \|_2 \| \mathbf{H}^T (\mathbf{H} \mathbf{H}^T)^{-1} \|_2 \cdot \left(\| \mathbf{F}(\mathbf{X}) - \mathbf{F}_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{X}) \|_2 + \| \boldsymbol{\epsilon} \|_2 \right) \quad (\text{by Lemma 12}) \\
& \leq \frac{\sqrt{p_1 p_2} \cdot \left(\| \mathbf{F}(\mathbf{X}) - \mathbf{F}_{\mathbf{V}, \mathbf{W}_0}^g(\mathbf{X}) \|_2 + \| \boldsymbol{\epsilon} \|_2 \right)}{\sqrt{\min \text{eig}(\mathbf{H} \mathbf{H}^T)}} \quad (\text{by Lemma 45 and Lemma 11}).
\end{aligned}$$

Combining Eq. (77) and Proposition 6 by the union bound, we thus have

$$\begin{aligned}
& \Pr_{\mathbf{x}, \mathbf{V}, \mathbf{W}_0} \left\{ \text{term D} + \text{term B of Eq. (22)} \geq \frac{\sqrt{n} \|g\|_1 \left(\frac{q}{\sqrt{p_2}} + \sqrt{\frac{Q(p_1, d)}{d}} \right) + \| \boldsymbol{\epsilon} \|_2}{\sqrt{J(n, p_1, p_2, d, q)}} \right\} \\
& \leq \frac{d^2}{(p_1 + 1)e^{d+1}} + \frac{8}{q^2}.
\end{aligned}$$

□

J Details Related to Learnable Set

In this part, we first restate Proposition 2 in a more precise way, i.e., Proposition 46 in Appendix J.1 and Proposition 47 in Appendix J.2. Then, in Appendix J.3 we discuss the generalization performance of ground-truth functions outside the learnable set.

J.1 $\mathcal{F}_{(3)}^{\ell_2}$ contains all polynomials with finite degree

By the following proposition, we show that $\mathcal{F}_{(3)}^{\ell_2}$ contains all polynomials with finite degree. We formally state it in the following proposition.

Proposition 46. *Let k be a finite non-negative integer. For any $f(\mathbf{x}) = \sum_{i=0}^k c_i (\mathbf{x}^T \mathbf{a}_i)^i$ where $c_i \in \mathbb{R}$ and $\mathbf{a}_i \in \mathbb{R}^d$, we must have $f \in \mathcal{F}_{(3)}^{\ell_2}$.*

We prove Proposition 46 in Appendix K. Although Proposition 46 is only for no-bias situation of 3-layer NTK, we can easily prove the similar results for the biased 3-layer NTK with the same proof technique.

J.2 $\mathcal{F}_{(3)}^{\ell_2}$ is a superset of $\mathcal{F}_{(2), b}^{\ell_2}$ (recall the definition of $\mathcal{F}_{(2), b}^{\ell_2}$ in Section 4.2)

The learnable sets of both 3-layer and 2-layer NTK models also contain polynomials with *infinite* degree. Notice that not all infinite-degree polynomials belong to the learnable sets, because the norm of the corresponding function g may not be finite. As we mentioned in footnote 4, the constrain $\|g\|_\infty < \infty$ can be relaxed to $\|g\|_1 < \infty$. However, with $\|g\|_1 < \infty$, the comparison among those learnable sets becomes more difficult. For convenience, we just relax the constraint to $\|g\|_2 < \infty$ (instead of $\|g\|_1 < \infty$) in the following result.

Proposition 47. *Under the constraint of $\|g\|_2 < \infty$, the learnable set of the 3-layer NTK (no bias) is at least as large as the 2-layer NTK (both with and without bias), i.e., $\mathcal{F}_{(2)}^{\ell_2} \cup \mathcal{F}_{(2), b}^{\ell_2} \subseteq \mathcal{F}_{(3)}^{\ell_2}$. The learnable set of 2-layer NTK with bias is larger than that of 2-layer NTK without bias i.e., $\mathcal{F}_{(2)}^{\ell_2} \subset \mathcal{F}_{(2), b}^{\ell_2}$. The learnable sets of 2-layer NTK with different bias settings are the same i.e., $\mathcal{F}_{(2), b_1}^{\ell_2} = \mathcal{F}_{(2), b_2}^{\ell_2}$ for any $b_1, b_2 \in (0, 1)$.*

We prove Proposition 47 in Appendix L. An important message conveyed by Proposition 47 is that, 3-layer NTK can at least learn all learnable functions for 2-layer NTK under the constraint $\|g\|_2 < \infty$. We conjecture that the same result may also hold for $\|g\|_1 < \infty$, which we leave for future work.




Figure 9: Curves of test MSE of 2-layer NTK with normal bias with respect to n for the ground-truth functions $f_{k,\alpha}(\mathbf{x})$ where $p_1, p_2 \rightarrow \infty$, $d = 2$, and $\epsilon = \mathbf{0}$. Every curve is the average of 10 simulation runs.

J.3 Generalization performance of ground-truth functions outside the learnable set

One may wonder what happens to the generalization performance for functions outside the learnable set. Notice that although we have proven that ground-truth functions inside the learnable set can be learned, it is possible that some functions outside the learnable set could still be learnable. For 2-layer NTK models without bias, [Ju et al. \(2021\)](#) shows that if a ground-truth function has a positive distance away from the learnable set, then such distance becomes the lower bound of the generalization error. Such ground-truth functions with positive distance exist for 2-layer NTK, e.g., $(\mathbf{x}^T \mathbf{e}_1)^3$, because $\mathcal{F}_{(2)}^{\ell_2}$ does not contain odd power polynomials except linear functions. However, for 2-layer NTK with bias or 3-layer NTK, there do not exist such ground-truth functions with a positive distance away from the learnable set. In other words, functions outside the learnable set is still in the closure of the corresponding learnable set. Thus, it is unclear whether or not those functions have a very different generalization performance compared with functions inside the learnable set.

We now use simulation results in [Fig. 9](#) to show that functions outside the learnable set may indeed exhibit qualitatively different generalization performance (and thus [Proposition 47](#) will be meaningful in capturing ground-truth functions with good generalization performance). We construct an example of functions inside and outside the learnable set (in the sense of finite $\|g\|_2$, consistent with [Proposition 47](#)). For simplicity, we focus on $\mathcal{F}_{(2),\text{NLB}}^{\ell_2}$, which is the learnable set for the 2-layer NTK with normal bias. We then consider a specific type of normalized ground-truth functions $f_{k,\alpha} := \bar{f}_{k,\alpha} / \|\bar{f}_{k,\alpha}\|_2$ where $\bar{f}_{k,\alpha}(\mathbf{x}) := \sum_{i=1}^k i^\alpha (\mathbf{x}^T \mathbf{e}_d)^i$. By previous discussion, we have already known that if k is finite, then $f_{k,\alpha} \in \mathcal{F}_{(2),\text{NLB}}^{\ell_2}$. However, when $k = \infty$, then whether $f_{\infty,\alpha} \in \mathcal{F}_{(2),\text{NLB}}^{\ell_2}$ or not is determined by the value of α . We let $d = 2$ and choose the value of α to be -3.5 , -3 , and -1.1 , respectively. It can be verified that $f_{k,\alpha} \in \mathcal{F}_{(2),\text{NLB}}^{\ell_2}$ when $\alpha = -3.5$ or $\alpha = -3$, while $f_{k,\alpha} \notin \mathcal{F}_{(2),\text{NLB}}^{\ell_2}$ when $\alpha = -1.1$. In numerical experiments, it is difficult to directly calculate $f_{\infty,\alpha}$, as we do not know the close form of $f_{\infty,\alpha}$. Therefore, we use $f_{k,\alpha}$ to approach $f_{\infty,\alpha}$ by increasing k . In [Fig. 9\(a\)](#), we let $\alpha = -3.5$ and plot the test MSE with respect to n when $k = 3$ (blue curve), $k = 10^2$ (orange curve), and $k = 10^5$ (green curve), respectively. We can see that these three curves almost overlap with each other, which implies that increasing k does not alter the test error significantly. (Similar phenomenon also appears in [Fig. 9\(b\)](#) where $\alpha = -3$.) In contrast, when we let $\alpha = -1.1$ in [Fig. 9\(c\)](#), larger k leads to a much flatter curve. This phenomenon suggests that when $k \rightarrow \infty$, providing more training data becomes less effective in lowering the test error. Besides, by comparing the curve of $k = 10^5$ in [Fig. 9\(a\)](#) and (c), we can see that the curve in [Fig. 9\(c\)](#) is higher than the one in [Fig. 9\(a\)](#) by several orders of magnitude. Therefore, we can tell that the functions inside and outside the learnable set could have very different generalization performance.

The setup of [Fig. 10](#) is the same as that of [Fig. 9](#) except that here we let x-axis be k . In [Fig. 10](#), we can see that the curves of $\alpha = -3.5$ and $\alpha = -3$ (finite $\|g\|_2$) in all sub-figures (a)(b)(c)(d) are almost flat with respect to k . In contrast, the curves of $\alpha = -1.1$ (infinite $\|g\|_2$) keep increasing with respect to k , and have much higher generalization error when k is large than those with finite $\|g\|_2$. This also validates our conjecture that the functions inside and outside the learnable set could have very different generalization performance.




Figure 10: Curves of test MSE of 2-layer NTK with normal bias with respect to k for the ground-truth functions $f_{k,\alpha}(\mathbf{x})$ where $p_1, p_2 \rightarrow \infty$, $d = 2$, and $\epsilon = 0$. Every curve is the average of 10 simulation runs.

K Proof of Proposition 46

Proof. We prove Proposition 46 by using similar methods as in Ju et al. (2021). For any $f_g \in \mathcal{F}_{(3)}^{\ell_2}$, we have

$$f_g(\mathbf{x}) = g \otimes h^{(3)}(\mathbf{x}) := \int_{\text{SO}(d)} g(\mathbf{S}e_d)h^{(3)}(\mathbf{S}^{-1}\mathbf{x})d\mathbf{S}, \quad (78)$$

$$h^{(3)}(\mathbf{x}) := K^{\text{Three}}(\mathbf{x}^T \mathbf{e}_d), \quad (79)$$

where $\mathbf{e}_d := [0 \ 0 \ \dots \ 0 \ 1]^T \in \mathbb{R}^d$, and \mathbf{S} is a $d \times d$ orthogonal matrix that denotes a rotation in S^{d-1} , chosen from the set $\text{SO}(d)$ of all rotations. An important property of the convolution Eq. (78) is that it corresponds to multiplication in the frequency domain, similar to Fourier coefficients. To define such a transformation to the frequency domain, we use a set of hyper-spherical harmonics $\Xi_{\mathbf{K}}^l$ (Vilenkin, 1968; Dokmanic & Petrinovic, 2009) when $d \geq 3$, which forms an orthonormal basis for functions on S^{d-1} . These harmonics are indexed by l and \mathbf{K} , where $\mathbf{K} = (k_1, k_2, \dots, k_{d-2})$ and $l = k_0 \geq k_1 \geq k_2 \geq \dots \geq k_{d-2} \geq 0$ (those k_i 's and l are all non-negative integers). Any function $f \in L^2(S^{d-1} \mapsto \mathbb{R})$ (including even δ -functions (Li & Wong, 2013)) can be decomposed uniquely into these harmonics, i.e., $f(\mathbf{x}) = \sum_l \sum_{\mathbf{K}} c_f(l, \mathbf{K}) \Xi_{\mathbf{K}}^l(\mathbf{x})$, where $c_f(\cdot, \cdot)$ are projections of f onto the basis function.

In Eq. (78), let $c_g(\cdot, \cdot)$ and $c_h(\cdot, \cdot)$ denote the coefficients corresponding to the decompositions of g and h , respectively. Then, we must have (Dokmanic & Petrinovic, 2009)

$$c_{f_g}(l, \mathbf{K}) = \Lambda \cdot c_g(l, \mathbf{K})c_{h^{(3)}}(l, \mathbf{0}), \quad (80)$$

where Λ is some normalization constant.

Eq. (80) describes an interesting “filtering” interpretation on $\mathcal{F}_{(3)}^{\ell_2}$. Specifically, $h^{(3)}$ and $c_{h^{(3)}}$ work like a channel or a filter in a wireless communication system, where c_g denotes the transmitted signal and c_{f_g} denotes the received signal. Therefore, for any basis function $f(\mathbf{x}) = \Xi_{\mathbf{K}}^l(\mathbf{x})$, as long as $c_{h^{(3)}}(l, \mathbf{0}) \neq 0$, we must have $f = f_g \in \mathcal{F}_{(3)}^{\ell_2}$ where the corresponding $g(\cdot)$ can simply be chosen as $g(\mathbf{z}) = \frac{\Xi_{\mathbf{K}}^l(\mathbf{z})}{\Lambda c_{h^{(3)}}(l, \mathbf{0})}$. Indeed, we have the following proposition about values of $c_{h^{(3)}}(l, \mathbf{0})$.

Proposition 48. $c_{h^{(3)}}(l, \mathbf{0}) > 0$ for all $l = 0, 1, 2, \dots$.

We provide its proof in Appendix K.1.

By Proposition 48, we know that all harmonics $\Xi_{\mathbf{K}}^l \in \mathcal{F}_{(3)}^{\ell_2}$. Notice that the set $\mathcal{F}_{(3)}^{\ell_2}$ is invariant under addition and scale operation¹². Therefore, any finite sum of $\Xi_{\mathbf{K}}^l$ belongs to $\mathcal{F}_{(3)}^{\ell_2}$. Notice that for any non-negative integer i and a real-valued vector $\mathbf{a} \in \mathbb{R}^d$, a polynomial $(\mathbf{x}^T \mathbf{a})^i$ consists of a finite sum of harmonic basis. Thus, $\mathcal{F}_{(3)}^{\ell_2}$ contains any polynomials $(\mathbf{x}^T \mathbf{a})^l$ for all $l = 0, 1, 2, \dots$. Proposition 46 thus follows. \square

K.1 Proof of Proposition 48

It is relatively easy to prove the result when $d = 2$, which is omitted here. We focus on the general case when $d \geq 3$. By Eq. (115) of Ju et al. (2021), the harmonics Ξ_0^l can be expressed by

$$\Xi_0^l(\mathbf{x}) = A_0^l \sum_{k=0}^{\lfloor \frac{l}{2} \rfloor} (-1)^k \frac{\Gamma(l - k + \frac{d-2}{2})}{\Gamma(\frac{d-2}{2})k!(l-2k)!} (2\mathbf{x}^T \mathbf{e}_d)^{l-2k}, \quad (81)$$

where A_0^l is a positive number as the normalization factor of Ξ_0^l . We give a few examples of Ξ_0^l as follows.

$$\begin{aligned} \Xi_0^0(\mathbf{x}) &= A_0^0, \\ \Xi_0^1(\mathbf{x}) &= A_0^1(d-2)\mathbf{x}^T \mathbf{e}_d, \\ \Xi_0^2(\mathbf{x}) &= A_0^2 \frac{d-2}{2} (d(\mathbf{x}^T \mathbf{e}_d)^2 - 1), \\ \Xi_0^3(\mathbf{x}) &= A_0^3 \frac{d-2}{2} \cdot d \cdot ((\mathbf{x}^T \mathbf{e}_d)^3 - \mathbf{x}^T \mathbf{e}_d). \end{aligned}$$

Recalling Eq. (6), we perform a Taylor expansion of $K^{\text{Three}}(\cdot)$. Let u_0, u_1, \dots denote the Taylor expansion coefficients of $2d \cdot K^{\text{Three}}$, i.e.,

$$2d \cdot K^{\text{Three}}(a) = \sum_{k=0}^{\infty} u_k a^k. \quad (82)$$

The following lemma shows that all coefficients in Eq. (82) are positive.

Lemma 49. For all $k = 0, 1, 2, \dots$, we have $u_k > 0$ in Eq. (82).

Proof. By Lemma 17, for any $a, b \in [0, 1]$, we have

$$2d \cdot K^{\text{RF}}(a) = \frac{1}{\pi} \left(1 + \frac{\pi}{2}a + \sum_{k=0}^{\infty} \frac{2(2k)!}{(k+1)(2k+1)(k!)^2} \left(\frac{a}{2}\right)^{2k+2} \right), \quad (83)$$

$$K^{\text{Two}}(b) = \frac{b}{4} + \frac{1}{2\pi} \sum_{k=0}^{\infty} \frac{(2k)!}{(k!)^2} \frac{4}{2k+1} \left(\frac{b}{2}\right)^{2k+2}. \quad (84)$$

By Lemma 43, we know that $2d \cdot K^{\text{RF}}(a) \in [0, 1]$. Thus, we can let $b = 2d \cdot K^{\text{RF}}(a)$ in Eq. (84) and then apply Eq. (83), i.e.,

$$\begin{aligned} & K^{\text{Two}}(2d \cdot K^{\text{RF}}(a)) \\ &= \frac{1}{4\pi} \left(1 + \frac{\pi}{2}a + \sum_{k=0}^{\infty} \frac{2(2k)!}{(k+1)(2k+1)(k!)^2} \left(\frac{a}{2}\right)^{2k+2} \right) \\ &+ \frac{1}{2\pi} \sum_{l=0}^{\infty} \frac{(2l)!}{(l!)^2} \frac{4}{2l+1} \left(\frac{1}{2\pi} \left(1 + \frac{\pi}{2}a + \sum_{k=0}^{\infty} \frac{2(2k)!}{(k+1)(2k+1)(k!)^2} \left(\frac{a}{2}\right)^{2k+2} \right) \right)^{2l+2}. \quad (85) \end{aligned}$$

¹²Specifically, if $f_{g_1}, f_{g_2} \in \mathcal{F}_{(3)}^{\ell_2}$, then $f_{g_1+g_2} := f_{g_1} + f_{g_2} \in \mathcal{F}_{(3)}^{\ell_2}$ and $f_{\alpha g_1} := \alpha f_{g_1} \in \mathcal{F}_{(3)}^{\ell_2}$.

By Eq. (6) and Eq. (82), we know that u_k is the coefficient of a^k in Eq. (85). In order to know the sign of u_k , it remains to combine similar terms in Eq. (85). To that end, we apply Lemma 27 and have

$$\begin{aligned}
u_0 &= \frac{1}{4\pi} + \frac{1}{2\pi} \sum_{l=0}^{\infty} \frac{(2l)!}{(l!)^2} \frac{4}{2l+1} \left(\frac{1}{2\pi}\right)^{2l+2}, \\
u_1 &= \frac{1}{8} + \frac{1}{2\pi} \sum_{l=0}^{\infty} \frac{(2l)!}{(l!)^2} \frac{4}{2l+1} \left(\frac{1}{2\pi}\right)^{2l+2} (2l+2) \left(\frac{\pi}{2}\right)^{2l+1}, \\
u_{2i+1} &= \frac{1}{2\pi} \sum_{l=0}^{\infty} \frac{(2l)!}{(l!)^2} \frac{4}{2l+1} \left(\frac{1}{2\pi}\right)^{2l+2} \cdot \sum_{\substack{k_0+k_1+k_2+k_4+\dots+k_{2i}=2l+2 \\ k_1+2k_2+4k_4+\dots+2ik_{2i}=2i+1}} (k_0, k_1, k_2, k_4, \dots, k_{2i})! \left(\frac{\pi}{2}\right)^{k_1} \prod_{j=0}^{i-1} \left(\frac{2(2j)!}{(j+1)(2j+1)(j!)^2} \left(\frac{1}{2}\right)^{2j+2}\right)^{k_{2j+2}}, \\
u_{2i+2} &= \frac{1}{4\pi} \frac{2(2i)!}{(i+1)(2i+1)(i!)^2} \left(\frac{1}{2}\right)^{2i+2} \\
&\quad + \frac{1}{2\pi} \sum_{l=0}^{\infty} \frac{(2l)!}{(l!)^2} \frac{4}{2l+1} \left(\frac{1}{2\pi}\right)^{2l+2} \cdot \sum_{\substack{k_0+k_1+k_2+k_4+\dots+k_{2i}=2l+2 \\ k_1+2k_2+4k_4+\dots+2ik_{2i}=2i+2}} (k_0, k_1, k_2, k_4, \dots, k_{2i})! \left(\frac{\pi}{2}\right)^{k_1} \prod_{j=0}^{i-1} \left(\frac{2(2j)!}{(j+1)(2j+1)(j!)^2} \left(\frac{1}{2}\right)^{2j+2}\right)^{k_{2j+2}}.
\end{aligned}$$

As we can see, every term in those expressions of u_0, u_1, \dots is positive, which implies that $u_k > 0$ for all $k = 0, 1, \dots$. \square

From Eq. (82), we have $2d \cdot K^{\text{Three}}(\mathbf{x}^T \mathbf{e}_d) = \sum_{k=0}^{\infty} u_k (\mathbf{x}^T \mathbf{e}_d)^k$. We now consider the decomposition of each $(\mathbf{x}^T \mathbf{e}_d)^k$ into harmonics.

Lemma 50. *Let a and b be two non-negative integers. Define the function*

$$Q(a, b) := \int_{\mathcal{S}^{d-1}} (\mathbf{x}^T \mathbf{e}_d)^a \cdot \Xi_0^b(\mathbf{x}) d\mu(\mathbf{x}). \quad (86)$$

We must have

$$Q(2k, 2m+1) = Q(2k+1, 2m) = 0, \quad (87)$$

$$Q(2k, 2m) \begin{cases} > 0, & \text{if } m \leq k, \\ = 0, & \text{if } m > k, \end{cases} \quad (88)$$

and

$$Q(2k+1, 2m+1) \begin{cases} > 0, & \text{if } m \leq k, \\ = 0, & \text{if } m > k. \end{cases} \quad (89)$$

Proof. By Eq. (81), we have $\Xi_0^b(-\mathbf{x}) = (-1)^b \Xi_0^b(\mathbf{x})$. Thus, when $a+b$ is odd, the function $(\mathbf{x}^T \mathbf{e}_d)^a \cdot \Xi_0^b(\mathbf{x})$ is an odd function with respect to \mathbf{x} . By symmetry of \mathcal{S}^{d-1} , we then have $Q(a, b) = 0$ when $a+b$ is odd, i.e., Eq. (87) holds. Eq. (88) has been proved in Lemma 53 of Ju et al. (2021) by mathematical induction. Here we prove Eq. (89).

By Eq. (118) of Ju et al. (2021), for any a , we have

$$Q(a+1, l+1) = q_{l,1} \cdot Q(a, l+2) + q_{l,2} \cdot Q(a, l), \quad (90)$$

where $q_{l,1} > 0$ and $q_{l,2} > 0$. Applying Eq. (90) for $a = 2k$ and $l = 2m$, we have

$$Q(2k+1, 2m+1) = q_{2m,1} \cdot Q(2k, 2m+2) + q_{2m,2} \cdot Q(2k, 2m).$$

By Eq. (88), the result of Eq. (89) thus follows. \square

Notice that

$$\begin{aligned}
c_{h^{(3)}}(l, \mathbf{0}) &= \int_{\mathcal{S}^{d-1}} h^{(3)}(\mathbf{x}) \Xi_{\mathbf{0}}^l(\mathbf{x}) d\mu(\mathbf{x}) \\
&= \frac{1}{2d} \int_{\mathcal{S}^{d-1}} 2d \cdot K^{\text{Three}}(\mathbf{x}^T \mathbf{e}_d) \Xi_{\mathbf{0}}^l(\mathbf{x}) d\mu(\mathbf{x}) \text{ (by Eq. (79))} \\
&= \frac{1}{2d} \sum_{k=0}^{\infty} u_k \cdot Q(k, l) \text{ (by Eq. (82) and Eq. (86))} \\
&> 0 \text{ (by Lemma 49 and Lemma 50).}
\end{aligned}$$

The result of Proposition 48 thus follows.

L Proof of Proposition 47

Proof. Using similar decomposition in Eq. (78), we define filter functions $h^{(2)}$ (for 2-layer NTK, no-bias) and $h_b^{(2)}$ (for 2-layer NTK, with bias). The corresponding harmonic coefficients are denoted by $c_{h^{(2)}}$ and $c_{h_b^{(2)}}$. We have the following result about the magnitude of those harmonic coefficients.

Lemma 51. *For any $b_1, b_2 \in (0, 1)$, we must have $c_{h_{b_1}^{(2)}}(2k, \mathbf{0}) = \Theta(c_{h^{(2)}}(2k, \mathbf{0}))$, $c_{h_{b_1}^{(2)}}(k, \mathbf{0}) = \Theta(c_{h_{b_2}^{(2)}}(k, \mathbf{0}))$, and $c_{h^{(3)}}(k, \mathbf{0}) = \Omega(c_{h_{b_1}^{(2)}}(k, \mathbf{0}))$. Here, $\Theta(\cdot)$ and $\Omega(\cdot)$ denote the orders as k becomes large.*

We prove Lemma 51 in Appendix M.

Lemma 51 has the following implications for the magnitude of the harmonics coefficients when the leading index of harmonics is large (i.e., k in Lemma 51 is large). The first statement states that, for 2-layer NTK, the setting with bias and the setting without bias have the same order of harmonics coefficients for even terms. (For odd terms, recall that for 2-layer NTK without bias, the coefficients of odd terms except linear term are zero. In contrast, for 2-layer NTK with bias, the coefficients of odd terms are not zero (Ju et al., 2021). Hence, the first statement does not hold for odd terms). The second statement states that, the coefficients of harmonics for 2-layer NTK have the same order with respect to k for all non-zero bias. The third statement states that, the coefficients for 3-layer NTK even without bias is not smaller (in order) than 2-layer NTK with bias.

By comparing the magnitude of these filter coefficients, we can then compare whether polynomials with infinite degree belong to each of the learnable sets. Specifically, consider an infinite-degree polynomial with the form $f_g(\mathbf{x}) = \sum_{l, \mathbf{K}} \alpha_{l, \mathbf{K}} \cdot \Xi_{\mathbf{K}}^l(\mathbf{x})$. By Eq. (80), we have $g(\mathbf{z}) = \sum_{l, \mathbf{K}} \frac{\alpha_{l, \mathbf{K}} \cdot \Xi_{\mathbf{K}}^l(\mathbf{z})}{\Lambda \cdot c_h(l, \mathbf{0})}$, where h can be $h^{(3)}$, $h^{(2)}$, or $h_b^{(2)}$. Thus, the magnitude of $c_h(l, \mathbf{0})$ determines the norm of g . Specifically, we have $\|g\|_2 = \sum_{l, \mathbf{K}} \frac{\alpha_{l, \mathbf{K}}}{\Lambda \cdot c_h(l, \mathbf{0})}$ due to the orthogonality of harmonics $\Xi_{\mathbf{K}}^l$. Note that $c_{h_{b_1}^{(2)}}(k, \mathbf{0}) = \Theta(c_{h_{b_2}^{(2)}}(k, \mathbf{0}))$ by Lemma 51. Thus, if $\|g\|_2$ is finite for 2-layer NTK with bias $b_1 > 0$, then it must also be finite for 2-layer NTK with a different bias $b_2 > 0$. This implies that The learnable sets of 2-layer NTK with different bias settings are the same i.e., $\mathcal{F}_{(2), b_1}^{\ell_2} = \mathcal{F}_{(2), b_2}^{\ell_2}$ for any $b_1, b_2 \in (0, 1)$. Similarly, for 3-layer NTK, by Lemma 51, we can also show that $\mathcal{F}_{(2)}^{\ell_2} \cup \mathcal{F}_{(2), b}^{\ell_2} \subseteq \mathcal{F}_{(3)}^{\ell_2}$ and $\mathcal{F}_{(2)}^{\ell_2} \subset \mathcal{F}_{(2), b}^{\ell_2}$. Therefore, the result of Proposition 47 follows. \square

M Proof of Lemma 51

The following lemma shows the relationship between harmonic coefficients and Taylor coefficients.

Lemma 52. *Consider two polynomial functions $h_{\alpha}(\mathbf{x}) := \sum_{k=0}^{\infty} u_{\alpha, k} (\mathbf{x}^T \mathbf{e}_d)^k$ and $h_{\beta}(\mathbf{x}) := \sum_{k=0}^{\infty} u_{\beta, k} (\mathbf{x}^T \mathbf{e}_d)^k$ where $u_{\alpha, k} \geq 0$ and $u_{\beta, k} \geq 0$ for all k . Let $c_{h_{\alpha}}$ and $c_{h_{\beta}}$ denote their harmonic coefficients. If $u_{\alpha, k} = O(u_{\beta, k})$ (where $O(\cdot)$ denotes the order when k is large), then $c_{h_{\alpha}}(l, \mathbf{0}) =$*

$O(c_{h_\beta}(l, \mathbf{0}))$ for large l . The same is true if we restrict to only even harmonics, i.e., if $u_{\alpha, 2k} = O(u_{\beta, 2k})$, then $c_{h_\alpha}(2l, \mathbf{0}) = O(c_{h_\beta}(2l, \mathbf{0}))$ for large l . This lemma also holds if $O(\cdot)$ is replaced by $\Omega(\cdot)$ or $\Theta(\cdot)$.

Proof. Notice that

$$\begin{aligned} c_{h_\alpha}(l, \mathbf{0}) &= \int_{\mathcal{S}^{d-1}} h(\mathbf{x}) \Xi_{\mathbf{0}}^l(\mathbf{x}) d\mu(\mathbf{x}) \\ &= \sum_{k=0}^{\infty} u_{\alpha, k} \cdot Q(k, l) \text{ (by Eq. (86))} \\ &= \sum_{k=l}^{\infty} u_{\alpha, k} \cdot Q(k, l) \text{ (by Lemma 50).} \end{aligned}$$

and

$$\begin{aligned} c_{h_\alpha}(2l, \mathbf{0}) &= \int_{\mathcal{S}^{d-1}} h(\mathbf{x}) \Xi_{\mathbf{0}}^{2l}(\mathbf{x}) d\mu(\mathbf{x}) \\ &= \sum_{k=0}^{\infty} u_{\alpha, k} \cdot Q(k, 2l) \text{ (by Eq. (86))} \\ &= \sum_{k=l}^{\infty} u_{\alpha, 2k} \cdot Q(2k, 2l) \\ &\quad (Q(k, 2l) \text{ is non-zero only when } k \text{ is even and not smaller than } 2l \text{ by Lemma 50).} \end{aligned}$$

Similarly, we have

$$\begin{aligned} c_{h_\beta}(l, \mathbf{0}) &= \sum_{k=l}^{\infty} u_{\beta, k} \cdot Q(k, l), \\ c_{h_\beta}(2l, \mathbf{0}) &= \sum_{k=l}^{\infty} u_{\beta, 2k} \cdot Q(2k, 2l). \end{aligned}$$

Notice that all $Q(\cdot, \cdot)$ and $u_{\cdot, \cdot}$ are all non-negative. Thus, we have

$$\frac{c_{h_\alpha}(l, \mathbf{0})}{c_{h_\beta}(l, \mathbf{0})} \in \left[\min_{k \geq l} \frac{u_{\alpha, k}}{u_{\beta, k}}, \max_{k \geq l} \frac{u_{\alpha, k}}{u_{\beta, k}} \right], \quad \frac{c_{h_\alpha}(2l, \mathbf{0})}{c_{h_\beta}(2l, \mathbf{0})} \in \left[\min_{k \geq l} \frac{u_{\alpha, 2k}}{u_{\beta, 2k}}, \max_{k \geq l} \frac{u_{\alpha, 2k}}{u_{\beta, 2k}} \right].$$

The result of this lemma thus follows. \square

With Lemma 52, in order to show Lemma 51, it is equivalent to compare Taylor coefficients of the expression of different kernels $h^{(3)}$, $h^{(2)}$, and $h_b^{(2)}$. Specifically, we are looking at the Taylor coefficients of the following expression:

$$T(x) := K(x) \frac{\pi - \arccos(K(x))}{2\pi}. \quad (91)$$

For 2-layer NTK, $K(x) = (1-a)x + a$ where $a \in [0, 1]$ corresponds to different choices of bias ($a = 0$ corresponds to no bias). For 3-layer NTK (no bias), we have $K(x) = 2d \cdot K^{\text{RF}}(x)$ (we neglect the constant $1/(2d)$ in K^{Three} , which does not change its order.) In other words, we have

$$T(x) = \begin{cases} K^{\text{Two}}(x), & \text{if } K(x) = x \text{ (i.e., 2-layer NTK no bias),} \\ K^{\text{Two}}((1-a)x + a), & \text{if } K(x) = (1-a)x + a \text{ (i.e., 2-layer NTK with bias } a > 0), \\ 2d \cdot K^{\text{Three}}(x), & \text{if } K(x) = 2d \cdot K^{\text{RF}}(x) \text{ (i.e., 3-layer NTK no bias).} \end{cases}$$

When $K(x) = x$, we already have the exact form of the Taylor expansion of $T(x)$ by Lemma 17. However, when $K(x)$ is a polynomial, it is not easy to get the close form of Taylor coefficients. We will first estimate the Taylor coefficients when $K(x) = (1-a)x + a$ in Appendix M.1. Second, we will estimate the Taylor coefficients when $K(x)$ is a polynomial with finite degree in Appendix M.2. Last, we will estimate the case of $K(x) = 2d \cdot K^{\text{RF}}(x)$ in Appendix M.3. (By Lemma 17, we know that $K^{\text{RF}}(x)$ is a polynomial with infinite degree.) The result of Lemma 51 then follows from these estimates.

M.1 Harmonic coefficients for 2-layer NTK with bias

After adding bias, the kernel of 2-layer NTK changes from $x \frac{\pi - \arccos x}{2\pi}$ to $((1-a)x + a) \frac{\pi - \arccos((1-a)x + a)}{2\pi}$. Here $a > 0$ denotes the bias setting ($a = 0$ corresponds to the no-bias setting). By Lemma 52, we only need to investigate the relationship between the Taylor coefficients. We define $u_{a,m}$ as the Taylor coefficients under the bias setting, i.e.,

$$((1-a)x + a) \frac{\pi - \arccos((1-a)x + a)}{2\pi} = \sum_{m=0}^{\infty} u_{a,m} \cdot x^m.$$

When $a = 0$, $u_{a,m}$ becomes $u_{0,m}$ and corresponds to the no-bias setting.

Lemma 53. For any $k \in \{2, 3, \dots\}$ and any $a \in [0, 1)$, we must have $\frac{u_{a,2k}}{u_{0,2k}} \in$

$$\left[\frac{1}{\left(2 + 2 \left(\frac{2a}{1-a} + 1\right)\right)^2} \frac{1}{1 + \frac{\left(\frac{1+a}{2}\right)^2}{1 - \left(\frac{1+a}{2}\right)^2}} \frac{1}{1-a^2} \cdot \frac{1 + \left(\frac{1-a}{1+a}\right)^{2k+1}}{1 + \frac{1-a}{1+a}}, \frac{1}{1-a^2} \cdot \frac{1 + \left(\frac{1-a}{1+a}\right)^{2k+1}}{1 + \frac{1-a}{1+a}} \right],$$

and $\frac{u_{a,2k-1}}{u_{0,2k}} \in$

$$\left[\frac{1}{\left(2 + 2 \left(\frac{2a}{1-a} + 1\right)\right)^2} \frac{1}{1 + \frac{\left(\frac{1+a}{2}\right)^2}{1 - \left(\frac{1+a}{2}\right)^2}} \frac{1}{1-a^2} \cdot \frac{1 - \left(\frac{1-a}{1+a}\right)^{2k}}{1 + \frac{1-a}{1+a}}, \frac{1}{1-a^2} \cdot \frac{1 - \left(\frac{1-a}{1+a}\right)^{2k}}{1 + \frac{1-a}{1+a}} \right].$$

We prove Lemma 53 in Appendix M.1.1.

Note that when $k \rightarrow \infty$, the terms that depend on k (i.e., $\left(\frac{1-a}{1+a}\right)^{2k+1}$ and $\left(\frac{1-a}{1+a}\right)^{2k}$) all approach 0. In other words, as k becomes larger, $u_{a,2k}$ (as well as $u_{a,2k+1}$) approaches (approximately) a constant (that only depends on a) multiple of $u_{0,2k}$. Therefore, by Lemma 53, we can then conclude that $u_{b_1,2k} = \Theta(u_{0,2k})$ and $u_{b_1,k} = \Theta(u_{b_2,k})$ when k is large for any $b_1 \in (0, 1)$ and $b_2 \in (0, 1)$. By Lemma 52, it immediately implies that $c_{h_{b_1}^{(2)}}(2k, \mathbf{0}) = \Theta(c_{h_{b_2}^{(2)}}(2k, \mathbf{0}))$,

$c_{h_{b_1}^{(2)}}(k, \mathbf{0}) = \Theta\left(c_{h_{b_2}^{(2)}}(k, \mathbf{0})\right)$. This proves the first and second statements of Lemma 51.

M.1.1 Proof of Lemma 53

We first write the form of $u_{0,l}$, i.e., the Taylor coefficients under no-bias setting. By Lemma 17, we have

$$x \frac{\pi - \arccos x}{2\pi} = \frac{x}{4} + \frac{1}{2\pi} \sum_{k=0}^{\infty} \frac{(2k)!}{(k!)^2} \frac{4}{2k+1} \left(\frac{x}{2}\right)^{2k+2}.$$

Thus, for $k \geq 1$, we have

$$u_{0,2k} = \frac{1}{2\pi} \frac{(2k-2)!}{((k-1)!)^2} \frac{4}{2k-1} \frac{1}{2^{2k}}, \quad (92)$$

$$u_{0,2k+1} = 0. \quad (93)$$

Next, we write the expression of $u_{a,l}$. To that end, we define

$$d_{a,2k,i} := \binom{2k+2i}{2k} (1-a)^{2k} a^{2i}, \quad (94)$$

$$d_{a,2k+1,i} := \binom{2k+2i+2}{2k+1} (1-a)^{2k+1} a^{2i+1}. \quad (95)$$

The following lemma provides the expression of $u_{a,l}$.

Lemma 54. For any $k \geq 1$, we must have

$$u_{a,2k} = \sum_{i=0}^{\infty} u_{0,2(k+i)} d_{a,2k,i},$$

$$u_{a,2k+1} = \sum_{i=0}^{\infty} u_{0,2(k+i+1)} d_{a,2k+1,i}.$$

We prove Lemma 54 in Appendix M.1.2.

Although we have the expression of $u_{a,k}$ by Lemma 54, it is not easy to directly estimate its value because terms like $u_{0,2(k+i)} d_{a,2k,i}$ have a very complicated form. Fortunately, some properties of $u_{0,2k}$ is very helpful. Specifically, by Eq. (92), we have $\frac{u_{0,2(k+1)}}{u_{0,2k}} = \frac{(2k-1)^2 2k}{k^2(2k+1) \cdot 4} = \frac{(2k-1)^2}{2k(2k+1)}$, whose value approaches 1 when $k \rightarrow \infty$. In other words, $u_{0,2k}$ has a very slow changing speed when k is large. Therefore, we can approximate the tail of $\sum_{i=0}^{\infty} u_{0,2(k+i)} d_{a,2k,i}$ by treating $u_{0,2(k+i)}$ as a constant. This allows us to focus our attention on estimating $\sum_{i=0}^{\infty} d_{a,2k,i}$ (and its tail $\sum_{i=l}^{\infty} d_{a,2k,i}$), whose value can be calculated by examining the coefficients of the Taylor expansion of $\frac{1}{1-((1-a)x+a)^2}$ (i.e., the sum of a geometric sequence $1, ((1-a)x+a)^2, ((1-a)x+a)^4, \dots$). The latter is much easier to study. We show these steps in detail as the following lemmas.

Define

$$l := \max \left\{ k, \left\lceil \frac{2a}{1-a} k \right\rceil \right\}. \quad (96)$$

The following lemma estimates the target ratio $\frac{u_{a,2k}}{u_{0,2k}}$ and $\frac{u_{a,2k-1}}{u_{0,2k}}$ in terms of $\sum_{i=0}^l d_{a,2k,i}$ and $\sum_{i=0}^{\infty} d_{a,2k,i}$.

Lemma 55. For any $k \in \{2, 3, \dots\}$, we must have

$$\frac{u_{a,2k}}{u_{0,2k}} \in \left[\frac{1}{\left(2 + 2 \left(\frac{2a}{1-a} + 1\right)\right)^2} \sum_{i=0}^l d_{a,2k,i}, \sum_{i=0}^{\infty} d_{a,2k,i} \right],$$

$$\frac{u_{a,2k-1}}{u_{0,2k}} \in \left[\frac{1}{\left(2 + 2 \left(\frac{2a}{1-a} + 1\right)\right)^2} \sum_{i=0}^l d_{a,2k-1,i}, \sum_{i=0}^{\infty} d_{a,2k-1,i} \right].$$

We prove Lemma 55 in Appendix M.1.3.

In order to finish the proof of Lemma 53, it only remains to estimate $\sum_{i=0}^l d_{a,2k,i}$, $\sum_{i=0}^l d_{a,2k-1,i}$, $\sum_{i=0}^{\infty} d_{a,2k,i}$, $\sum_{i=0}^{\infty} d_{a,2k-1,i}$, which are shown by the following two lemmas.

Lemma 56. For any $k \in \{2, 3, \dots\}$, we must have

$$\sum_{i=0}^{\infty} d_{a,2k,i} = \frac{1}{1-a^2} \cdot \frac{1 + \left(\frac{1-a}{1+a}\right)^{2k+1}}{1 + \frac{1-a}{1+a}}, \quad \sum_{i=0}^{\infty} d_{a,2k-1,i} = \frac{1}{1-a^2} \cdot \frac{1 - \left(\frac{1-a}{1+a}\right)^{2k}}{1 + \frac{1-a}{1+a}}.$$

We prove Lemma 56 in Appendix M.1.4

Lemma 57. Recall that l is defined in Eq. (96). For any $k \in \{2, 3, \dots\}$, we must have

$$\frac{\sum_{i=0}^l d_{a,2k,i}}{\sum_{i=0}^{\infty} d_{a,2k,i}}, \frac{\sum_{i=0}^l d_{a,2k-1,i}}{\sum_{i=0}^{\infty} d_{a,2k-1,i}} \in \left[\frac{1}{1 + \frac{\left(\frac{1+a}{2}\right)^2}{1 - \left(\frac{1+a}{2}\right)^2}}, 1 \right].$$

We prove Lemma 57 in Appendix M.1.5

The result of Lemma 53 follows by combining Lemma 56, Lemma 57, and Lemma 55.

M.1.2 Proof of Lemma 54

Proof. We have

$$\begin{aligned}
& ((1-a)x+a) \frac{\pi - \arccos((1-a)x+a)}{2\pi} \\
&= \sum_{m=0}^{\infty} u_{0,m} ((1-a)x+a)^m \\
&= \sum_{m=0}^{\infty} u_{0,m} \sum_{i=0}^m \binom{m}{i} (1-a)^i a^{m-i} x^i \\
&= \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} u_{0,i+j} \binom{i+j}{i} (1-a)^i a^j x^i \text{ (replace } m \text{ by } i+j \text{ and reorganize terms).}
\end{aligned}$$

Thus, we have

$$u_{a,i} = \sum_{j=0}^{\infty} u_{0,i+j} \binom{i+j}{i} (1-a)^i a^j.$$

Letting $i = 2k$, we have

$$\begin{aligned}
u_{a,2k} &= \sum_{j=0}^{\infty} u_{0,2k+j} \binom{2k+j}{2k} (1-a)^{2k} a^j \\
&= \sum_{i=0}^{\infty} u_{0,2(k+i)} \binom{2k+2i}{2k} (1-a)^{2k} a^{2i} \text{ (by Eq. (93) and letting } j = 2i) \\
&= \sum_{i=0}^{\infty} u_{0,2(k+i)} d_{a,2k,i} \text{ (by Eq. (94)).}
\end{aligned}$$

Similarly, we have

$$\begin{aligned}
u_{a,2k+1} &= \sum_{j=0}^{\infty} u_{0,2k+1+j} \binom{2k+1+j}{2k+1} (1-a)^{2k+1} a^j \\
&= \sum_{i=0}^{\infty} u_{0,2(k+i+1)} \binom{2k+2i+2}{2k+1} (1-a)^{2k+1} a^{2i+1} \text{ (by Eq. (93) and letting } j = 2i+1) \\
&= \sum_{i=0}^{\infty} u_{0,2(k+i+1)} d_{a,2k+1,i} \text{ (by Eq. (95)).}
\end{aligned}$$

The result of this lemma thus follows. \square

M.1.3 Proof of Lemma 55

Proof. By Eq. (92), we have

$$\begin{aligned}
\frac{u_{0,2(k+1)}}{u_{0,2k}} &= \frac{(2k-1)^2 2k}{k^2(2k+1) \cdot 4} = \frac{(2k-1)^2}{2k(2k+1)} \\
&\leq 1.
\end{aligned} \tag{97}$$

By iterating the above inequality, we have $u_{0,2(k+i)} \leq u_{0,2k}$ for all $i \geq 0$. By Lemma 54, we thus have

$$\frac{u_{a,2k}}{u_{0,2k}} = \frac{\sum_{i=0}^{\infty} u_{0,2(k+i)} d_{a,2k,i}}{u_{0,2k}} \leq \sum_{i=0}^{\infty} d_{a,2k,i}.$$

Similarly, we have

$$\frac{u_{a,2k-1}}{u_{0,2k}} = \frac{\sum_{i=0}^{\infty} u_{0,2(k+i)} d_{a,2k-1,i}}{u_{0,2k}} \leq \sum_{i=0}^{\infty} d_{a,2k-1,i}.$$

These prove the upper bounds in Lemma 55. To prove the lower bounds, note that for any $m \in \{0, 1, \dots, l\}$ (recall that l is defined in Eq. (96)), we must have

$$\begin{aligned}
\frac{u_{0,2(k+m)}}{u_{0,2k}} &= \prod_{i=0}^{m-1} \frac{(2k+2i-1)^2}{(2k+2i)(2k+2i+1)} \quad (\text{by Eq. (97)}) \\
&\geq \prod_{i=0}^{m-1} \frac{(2k+2i-1)^2}{(2k+2i+1)^2} \\
&= \frac{(2k-1)^2}{(2k+2m-1)^2} \\
&\geq \frac{k^2}{(2k+2m)^2} \quad (\text{using } 2k-1 \geq k, \text{ which is true because } k \geq 1) \\
&= \frac{1}{\left(2 + \frac{2m}{k}\right)^2} \\
&\geq \frac{1}{\left(2 + 2\left(\frac{2a}{1-a} + 1\right)\right)^2} \quad (\text{because } m \leq l \leq \frac{2a}{1-a}k + k \text{ for } k \geq 2).
\end{aligned}$$

Thus, we have

$$\begin{aligned}
\frac{u_{a,2k}}{u_{0,2k}} &= \frac{\sum_{i=0}^{\infty} u_{0,2(k+i)} d_{a,2k,i}}{u_{0,2k}} \quad (\text{by Lemma 54}) \\
&\geq \frac{\sum_{i=0}^l u_{0,2(k+i)} d_{a,2k,i}}{u_{0,2k}} \\
&\geq \frac{1}{\left(2 + 2\left(\frac{2a}{1-a} + 1\right)\right)^2} \sum_{i=0}^l d_{a,2k,i}.
\end{aligned}$$

Similarly, we have

$$\frac{u_{a,2k-1}}{u_{0,2k}} \geq \frac{1}{\left(2 + 2\left(\frac{2a}{1-a} + 1\right)\right)^2} \sum_{i=0}^l d_{a,2k-1,i}.$$

The result of this lemma thus follows. □

M.1.4 Proof of Lemma 56

We first state a useful fact.

Lemma 58. *For any $|r| < 1$, we have*

$$\frac{1}{1-r} = \sum_{i=0}^{\infty} r^i.$$

Proof. The result of this lemma directly follows the sum of a geometric series (noticing that $\lim_{i \rightarrow \infty} r^i = 0$ when $|r| < 1$). □

Proof of Lemma 56. The proof idea is to express the coefficients of the Taylor expansion of $\frac{1}{1 - ((1-a)x+a)^2}$ (where $|x| < 1$) in two different ways. On the one hand, we have

$$\begin{aligned}
& \frac{1}{1 - ((1-a)x+a)^2} \\
&= \sum_{m=0}^{\infty} ((1-a)x+a)^{2m} \text{ (by letting } r = (1-a)x+a \text{ in Lemma 58)} \\
&= \sum_{m=0}^{\infty} \sum_{j=0}^{2m} \binom{2m}{j} (1-a)^j a^{2m-j} x^j \\
&= \sum_{k=0}^{\infty} \left(\left(\sum_{i=0}^{\infty} \binom{2k+2i}{2k} (1-a)^{2k} a^{2i} \right) x^{2k} + \left(\sum_{i=0}^{\infty} \binom{2k+2i+2}{2k+1} (1-a)^{2k+1} a^{2i+1} \right) x^{2k+1} \right) \\
&\quad \text{(by letting } j = 2k, 2m = 2k + 2i \text{ for } x^{2k} \text{ and letting } j = 2k + 1, 2m = 2k + 2i + 2 \text{ for } x^{2k+1}) \\
&= \sum_{k=0}^{\infty} \left(\left(\sum_{i=0}^{\infty} d_{a,2k,i} \right) x^{2k} + \left(\sum_{i=0}^{\infty} d_{a,2k+1,i} \right) x^{2k+1} \right) \text{ (by Eq. (94) and Eq. (95)).} \tag{98}
\end{aligned}$$

On the other hand, we have

$$\begin{aligned}
& \frac{1}{1 - ((1-a)x+a)^2} \\
&= \frac{1}{1 - ((1-a)x+a)} \cdot \frac{1}{1 + ((1-a)x+a)} \\
&= \frac{1}{1-a} \cdot \frac{1}{1-x} \cdot \frac{1}{1+a} \cdot \frac{1}{1 + \frac{1-a}{1+a}x} \\
&= \frac{1}{1-a^2} \left(\sum_{i=0}^{\infty} x^i \right) \left(\sum_{j=0}^{\infty} \left(-\frac{1-a}{1+a} x \right)^j \right) \text{ (by Lemma 58)} \\
&= \frac{1}{1-a^2} \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} \left(-\frac{1-a}{1+a} \right)^j x^{i+j} \\
&= \frac{1}{1-a^2} \sum_{m=0}^{\infty} \left(\sum_{j=0}^m \left(-\frac{1-a}{1+a} \right)^j \right) x^m \text{ (combine terms of } x^{i+j} \text{ with } i+j = m) \\
&= \frac{1}{1-a^2} \sum_{m=0}^{\infty} \frac{1 - \left(-\frac{1-a}{1+a} \right)^{m+1}}{1 + \frac{1-a}{1+a}} x^m. \tag{99}
\end{aligned}$$

By comparing the coefficients in Eq. (98) and Eq. (99), the result of this lemma thus follows. \square

M.1.5 Proof of Lemma 57

We first prove a useful lemma.

Lemma 59. For any $a > b > c > 0$, we have

$$\frac{a}{b} < \frac{a-c}{b-c}.$$

Proof. Because $a > b > c > 0$, we have

$$b < a \implies bc < ac \implies ab - ac < ab - bc \implies a(b-c) < b(a-c) \implies \frac{a}{b} < \frac{a-c}{b-c}.$$

\square

Now we are ready to prove Lemma 57.

Proof of Lemma 57. Recall that l is defined in Eq. (96). For any $i \geq l$, we have

$$\frac{k+i}{i} = \frac{k}{i} + 1 \leq \frac{k}{l} + 1 \leq \frac{1+a}{2a} \quad (\text{because } l \geq \frac{2a}{1-a}k \text{ by Eq. (96)}). \quad (100)$$

Thus, by Eq. (94), we have

$$\begin{aligned} \frac{d_{a,2k,i+1}}{d_{a,2k,i}} &= \frac{\binom{2k+2i+2}{2k}}{\binom{2k+2i}{2k}} a^2 = \frac{(2k+2i+1)(2k+2i+2)}{(2i+1)(2i+2)} a^2 \\ &\leq \frac{(2k+2i)(2k+2i)}{(2i) \cdot (2i)} a^2 \quad (\text{by Lemma 59}) \\ &\leq \left(\frac{1+a}{2}\right)^2 \quad (\text{by Eq. (100)}). \end{aligned}$$

Similarly, by Eq. (95), we have

$$\begin{aligned} \frac{d_{a,2k-1,i+1}}{d_{a,2k-1,i}} &= \frac{\binom{2k+2i+2}{2k-1}}{\binom{2k+2i}{2k-1}} a^2 = \frac{(2k+2i+2)(2k+2i+1)}{(2i+2)(2i+3)} a^2 \\ &\leq \frac{(2k+2i+2)(2k+2i+3)}{(2i+2)(2i+3)} a^2 \\ &\leq \frac{(2k+2i)(2k+2i)}{(2i) \cdot (2i)} a^2 \quad (\text{by Lemma 59}) \\ &\leq \left(\frac{1+a}{2}\right)^2 \quad (\text{by Eq. (100)}). \end{aligned}$$

Iterating the above inequalities, we have

$$\frac{d_{a,2k,l+j}}{d_{a,2k,l}} \leq \left(\frac{1+a}{2}\right)^{2j}, \quad \text{and} \quad \frac{d_{a,2k-1,l+j}}{d_{a,2k-1,l}} \leq \left(\frac{1+a}{2}\right)^{2j}.$$

Thus, we have

$$\frac{\sum_{i=l+1}^{\infty} d_{a,2k,i}}{\sum_{i=0}^l d_{a,2k,i}} \leq \frac{\sum_{i=l+1}^{\infty} d_{a,2k,i}}{d_{a,2k,l}} = \sum_{j=1}^{\infty} \frac{d_{a,2k,l+j}}{d_{a,2k,l}} \leq \sum_{j=1}^{\infty} \left(\frac{1+a}{2}\right)^{2j} = \frac{\left(\frac{1+a}{2}\right)^2}{1 - \left(\frac{1+a}{2}\right)^2}.$$

We then have

$$\frac{\sum_{i=0}^{\infty} d_{a,2k,i}}{\sum_{i=0}^l d_{a,2k,i}} = \frac{\sum_{i=0}^l d_{a,2k,i} + \sum_{i=l+1}^{\infty} d_{a,2k,i}}{\sum_{i=0}^l d_{a,2k,i}} \leq 1 + \frac{\left(\frac{1+a}{2}\right)^2}{1 - \left(\frac{1+a}{2}\right)^2}.$$

Therefore, we conclude that

$$\frac{\sum_{i=0}^l d_{a,2k,i}}{\sum_{i=0}^{\infty} d_{a,2k,i}} \in \left[\frac{1}{1 + \frac{\left(\frac{1+a}{2}\right)^2}{1 - \left(\frac{1+a}{2}\right)^2}}, 1 \right].$$

Similarly, we have

$$\frac{\sum_{i=0}^l d_{a,2k-1,i}}{\sum_{i=0}^{\infty} d_{a,2k-1,i}} \in \left[\frac{1}{1 + \frac{\left(\frac{1+a}{2}\right)^2}{1 - \left(\frac{1+a}{2}\right)^2}}, 1 \right].$$

□

M.2 Expansion for a finite-degree polynomial

We plan to show the third statement of Lemma 51, which was presented in Appendix L. This is more difficult because $K^{\text{RF}}(x)$ is an infinite-degree polynomial (in contrast, Appendix M.1 deals with $K(x) = (1-a)x + a$, which is much simpler). To make progress, we first consider a finite-degree polynomial, and study the expansion. Then, in Appendix M.3, we will extend to $K^{\text{RF}}(x)$ which has infinite degree. Note that Appendix M.1 is a special case of Appendix M.2. However, since Appendix M.1 is much simpler and easy to understand, we retain the proof there, and use the result in Appendix M.2 only as a preparation for Appendix M.3.

Recall the definition of $T(x)$ in Eq. (91). We denote $K(x)$ as a polynomial, i.e.,

$$K(x) = \sum_{i=0}^{\infty} a_i x^i, \quad (101)$$

where a_i denote the coefficient of x^i in $K(x)$.

Define $u_m(\cdot)$ as a function that projects a polynomial in x to a real value such that

$$\tilde{K}(x) \frac{\pi - \arccos(\tilde{K}(x))}{2\pi} = \sum_{m=0}^{\infty} u_m(\tilde{K}(x)) \cdot x^m, \quad (102)$$

where $\tilde{K}(x)$ is any polynomial of x . In other words, $u_m(\tilde{K}(x))$ is the Taylor coefficient of x^m in $T(x)$ when $K(x) = \tilde{K}(x)$.

In this subsection, we let the number of terms of $K(x)$ be finite, i.e., there exists s such that $a_i = 0$ for all $i > s$. Further, we impose the following conditions.

Condition 2. (i) All coefficients of $K(x)$ are non-negative, i.e., $a_i \geq 0$ for all $i \in \mathbb{Z}_{\geq 0}$. (ii) The sum of all coefficients equals to 1, i.e., $\sum_{i=0}^{\infty} a_i = K(1) = 1$. (iii) $a_0 > 0$ and $a_1 > 0$.

The following lemma shows that when $K(x)$ is a polynomial with finite terms, the Taylor coefficients are on the same order as that of the even-power Taylor coefficients when $K(x) = x$. Note that, according to Eq. (102), when $K(x) = x$, $u_m(x)$ recovers the Taylor coefficients of the polynomial expansion of the function $x \frac{\pi - \arccos(x)}{2\pi}$.

Lemma 60. *Under Condition 2 and when $a_i = 0$ for all $i > s$, we must have*

$$\frac{u_j(K(x))}{u_{2\lceil j/2 \rceil}(x)} \in [\underline{C}, \overline{C}], \text{ for all } j = 1, 2, \dots,$$

where $\overline{C} > \underline{C} > 0$ are constants that only depends on $K(x)$ and are independent of j .

We prove Lemma 60 in Appendix M.2.1. Note that Lemma 60 can be seen as a generalization of Lemma 53, since $K(x) = (1-a) + a$ satisfies Condition 2 when $a \in (0, 1)$.

M.2.1 Proof of Lemma 60

We introduce some extra notations. Let b_i be the coefficients of x^i of $(K(x))^2$, i.e.,

$$(K(x))^2 = \sum_{i=0}^{2s} b_i x^i, \text{ which implies that } b_i = \sum_{j+k=i} a_j a_k \text{ for all } i \in \mathbb{Z}_{\geq 0}. \quad (103)$$

As in Lemma 27, for all $j \in \mathbb{Z}_{\geq 0}$, we define

$$t(m_0, m_1, \dots, m_j) := (m_0, m_1, \dots, m_j)! \cdot a_0^{m_0} a_1^{m_1} \dots a_j^{m_j} \text{ (we let } a_i = 0 \text{ if } i > s), \quad (104)$$

$$\mathcal{T}_{i,j} := \left\{ (m_0, m_1, \dots, m_j) \left| \begin{array}{l} m_0 + m_1 + \dots + m_j = i \\ m_1 + 2m_2 + \dots + j \cdot m_j = j \\ m_0, m_1, \dots, m_j \in \mathbb{Z}_{\geq 0} \end{array} \right. \right\},$$

$$d_{i,j} := \sum_{(m_0, m_1, \dots, m_j) \in \mathcal{T}_{i,j}} t(m_0, m_1, \dots, m_j). \quad (105)$$

Notation	Description	Definition/Expression
a_i	coefficients of x^i in $K(x)$	Eq. (101)
b_i	coefficients of x^i in $(K(x))^2$	Eq. (103)
$u_i(\tilde{K}(x))$	coefficients of x^i in $\tilde{K}(x)(\pi - \arccos(\tilde{K}(x)))/(2\pi)$	Eq. (102) and Eq. (115)
$d_{i,j}$	the coefficient of x^j in $(K(x))^i$	Eq. (105)
e_i	the coefficient of x^i in $1/(1 - (K(x))^2)$	Eq. (106) and Eq. (107)

Table 2: Summary of the notations of various coefficients.

By Lemma 27, we have

$$\begin{aligned} \frac{1}{1 - (K(x))^2} &= \sum_{i=0}^{\infty} (K(x))^{2i} = \sum_{i=0}^{\infty} \left(\sum_{j=0}^{\infty} d_{2i,j} x^j \right) \\ &= \sum_{j=0}^{\infty} \left(\sum_{i=\lceil \frac{j}{2s} \rceil}^{\infty} d_{2i,j} \right) x^j \text{ (since } d_{2i,j} = 0 \text{ when } j > 2i \cdot s \text{)}. \end{aligned}$$

Define

$$e_j \text{ for all } j \in \mathbb{Z}_{\geq 0} \text{ such that } \frac{1}{1 - (K(x))^2} = \sum_{j=0}^{\infty} e_j x^j, \quad (106)$$

i.e.,

$$e_j = \sum_{i=\lceil \frac{j}{2s} \rceil}^{\infty} d_{2i,j}. \quad (107)$$

We summarize those definitions in Table 2.

Lemma 61. *Under Condition 2, we must have*

$$\begin{aligned} \sum_{i=0}^{2s} b_i &= 1, \\ b_i &\in [0, 1] \text{ for all } i \in \{0, 1, \dots, 2s\}, \quad b_0 < 1, \text{ and } b_1 > 0. \end{aligned} \quad (108)$$

Proof. By Eq. (103) and Condition 2, we have $\sum_{i=0}^{2s} b_i = (K(1))^2 = 1$. Because $a_1 > 0$ and $\sum_{i=0}^s a_i = 1$, we have $a_0 < 1$. Thus, we have $b_0 = a_0^2 < 1$ and $b_1 = 2a_0 a_1 > 0$ (as a_0 is also positive). \square

Lemma 62. *There exist $\bar{c} \geq \underline{c} > 0$ such that for all $k \in \mathbb{Z}_{\geq 0}$, we must have $e_k \in [\underline{c}, \bar{c}]$.*

Proof. Because

$$\frac{1}{1 - (K(x))^2} = 1 + (K(x))^2 \cdot \frac{1}{1 - (K(x))^2},$$

we have

$$\sum_{j=0}^{\infty} e_j x^j = 1 + \left(\sum_{i=0}^s a_i x^i \right)^2 \cdot \left(\sum_{j=0}^{\infty} e_j x^j \right) = 1 + \left(\sum_{i=0}^{2s} b_i x^i \right) \cdot \left(\sum_{j=0}^{\infty} e_j x^j \right). \quad (109)$$

Comparing the coefficient of x^{j+2s} on both sides, we have

$$e_{j+2s} = \sum_{i=0}^{2s} e_{j+i} b_{2s-i} \text{ for all } j = 0, 1, \dots.$$

This is equivalent to

$$e_{j+2s} = b_0 e_{j+2s} + \sum_{i=0}^{2s-1} e_{j+i} b_{2s-i} \text{ for all } j = 0, 1, \dots.$$

Thus, we have

$$e_{j+2s} = \sum_{i=0}^{2s-1} \frac{b_{2s-i}}{1-b_0} e_{j+i} \text{ for all } j = 0, 1, \dots.$$

It implies that

$$e_{j+2s} \in \left[\left(\min_{i \in \{0,1,\dots,2s-1\}} e_{j+i} \right) \cdot \sum_{k=1}^{2s} \frac{b_k}{1-b_0}, \left(\max_{i \in \{0,1,\dots,2s-1\}} e_{j+i} \right) \cdot \sum_{k=1}^{2s} \frac{b_k}{1-b_0} \right]$$

for all $j = 0, 1, \dots$.

By Eq. (108), we have $\sum_{k=1}^{2s} b_k = 1 - b_0$, which implies that $\sum_{k=1}^{2s} \frac{b_k}{1-b_0} = 1$. Thus, we have

$$e_{j+2s} \in \left[\min_{i \in \{0,1,\dots,2s-1\}} e_{j+i}, \max_{i \in \{0,1,\dots,2s-1\}} e_{j+i} \right], \text{ for all } j = 0, 1, \dots.$$

Iteratively applying the above bounds, we then have

$$\begin{aligned} e_{2s}, e_{2s+1}, \dots, e_{4s-1} &\in \left[\min_{i \in \{0,1,\dots,2s-1\}} e_i, \max_{i \in \{0,1,\dots,2s-1\}} e_i \right], \\ e_{4s}, e_{4s+1}, \dots, e_{6s-1} &\in \left[\min_{i \in \{2s,2s+1,\dots,4s-1\}} e_i, \max_{i \in \{2s,2s+1,\dots,4s-1\}} e_i \right] \in \left[\min_{i \in \{0,1,\dots,2s-1\}} e_i, \right. \\ &\quad \left. \max_{i \in \{0,1,\dots,2s-1\}} e_i \right], \\ &\vdots \\ e_{2ks}, e_{2ks+1}, \dots, e_{2ks+2s-1} &\in \dots \in \left[\min_{i \in \{0,1,\dots,2s-1\}} e_i, \max_{i \in \{0,1,\dots,2s-1\}} e_i \right]. \end{aligned}$$

In other words,

$$e_k \in \left[\min_{i \in \{0,1,\dots,2s-1\}} e_i, \max_{i \in \{0,1,\dots,2s-1\}} e_i \right], \text{ for all } k = 2s, 2s+1, \dots. \quad (110)$$

By Eq. (109), we have

$$\begin{aligned} e_0 &= 1 + b_0 e_0, \\ e_1 &= b_1 e_0 + b_0 e_1, \\ e_2 &= b_2 e_0 + b_1 e_1 + b_0 e_2, \\ &\vdots \\ e_{2s} &= b_{2s} e_0 + b_{2s-1} e_1 + \dots + b_0 e_{2s}. \end{aligned}$$

Thus, we have

$$\begin{aligned} e_0 &= \frac{1}{1 - b_0}, \\ e_1 &= \frac{b_1 e_0}{1 - b_0}, \\ e_2 &= \frac{b_2 e_0 + b_1 e_1}{1 - b_0}, \\ &\vdots \\ e_{2s} &= \frac{b_{2s} e_0 + b_{2s-1} e_1 + \cdots + b_1 e_{2s-1}}{1 - b_0}. \end{aligned}$$

By Lemma 61 and using induction, we thus have

$$e_i > 0, \text{ for all } i = 0, 1, \dots, 2s - 1.$$

Thus, by Eq. (110), we have

$$e_k \in \left[\min_{i \in \{0, 1, \dots, 2s-1\}} e_i, \max_{i \in \{0, 1, \dots, 2s-1\}} e_i \right], \text{ for all } k = 0, 1, \dots.$$

□

Lemma 63. *When $i \geq j$, there exists a bijection between $\mathcal{T}_{i,j}$ and $\mathcal{T}_{i+1,j}$. Specifically, this bijection is $\mathcal{T}_{i,j} \longleftrightarrow \mathcal{T}_{i+1,j}: (m_0, m_1, \dots, m_j) \longleftrightarrow (m_0 + 1, m_1, \dots, m_j)$.*

Proof. It suffices to show that for any $(m_0, m_1, \dots, m_j) \in \mathcal{T}_{i+1,j}$, we must have $m_0 \geq 1$. To that end, note that when $i \geq j$, for any $(m_0, m_1, \dots, m_s) \in \mathcal{T}_{i+1,j}$, we have

$$\sum_{k=0}^j m_k = i + 1, \quad \sum_{k=1}^j k m_k = j.$$

Thus, we have

$$m_0 = \sum_{k=0}^j m_k - \sum_{k=1}^j k m_k + \sum_{k=1}^j (k-1) m_k = (i+1-j) + \sum_{k=1}^j (k-1) m_k \geq 1 \text{ (because } i \geq j).$$

The result of this lemma thus follows. □

Lemma 64. *If $2i \geq j$, then*

$$\frac{d_{2i+2,j}}{d_{2i,j}} \leq \left(\frac{2i}{2i-j} a_0 \right)^2.$$

(Notice that when $2i = j$, the right hand side is infinite. Nonetheless, this lemma still holds.)

Proof. We have

$$\frac{d_{2i+2,j}}{d_{2i,j}} = \frac{\sum_{\mathbf{m} \in \mathcal{T}_{2i+2,j}} t(\mathbf{m})}{\sum_{\mathbf{m} \in \mathcal{T}_{2i,j}} t(\mathbf{m})}.$$

Let $(m_0^{(k)}, m_1^{(k)}, \dots, m_j^{(k)})$ denote the k -th element in $\mathcal{T}_{2i,j}$. Because $m_0 + m_1 + \cdots + m_j = 2i$ and $m_1 + 2m_2 + \cdots + j \cdot m_j = j$, we have $m_0 \geq 2i - j$. Thus, using the definition of $t(\cdots)$ in Eq. (104), we have

$$\begin{aligned} & \frac{t(m_0^{(k)} + 2, m_1^{(k)}, \dots, m_j^{(k)})}{t(m_0^{(k)}, m_1^{(k)}, \dots, m_j^{(k)})} \\ &= \frac{(2i+1)(2i+2)}{(m_0+1)(m_0+2)} a_0^2 \\ &\leq \frac{(2i+1)(2i+2)}{(2i-j+1)(2i-j+2)} a_0^2 \text{ (because } m_0 \geq 2i-j) \\ &\leq \left(\frac{2i}{2i-j} a_0 \right)^2 \text{ (because } \frac{2i+1}{2i-j+1} \leq \frac{2i}{2i-j} \text{ and } \frac{2i+2}{2i-j+2} \leq \frac{2i}{2i-j}). \end{aligned}$$

By Lemma 63 and $2i \geq j$, we thus have

$$\begin{aligned} \frac{d_{2i+2,j}}{d_{2i,j}} &= \frac{t(m_0^{(1)} + 2, m_1^{(1)}, \dots, m_j^{(1)}) + \dots + t(m_0^{(\lceil \mathcal{T}_{2i,j} \rceil)} + 2, m_1^{(\lceil \mathcal{T}_{2i,j} \rceil)}, \dots, m_j^{(\lceil \mathcal{T}_{2i,j} \rceil)})}{t(m_0^{(1)}, m_1^{(1)}, \dots, m_j^{(1)}) + \dots + t(m_0^{(\lceil \mathcal{T}_{2i,j} \rceil)}, m_1^{(\lceil \mathcal{T}_{2i,j} \rceil)}, \dots, m_j^{(\lceil \mathcal{T}_{2i,j} \rceil)})} \\ &\leq \left(\frac{2i}{2i-j} a_0 \right)^2. \end{aligned}$$

□

Lemma 65. For any $j = 1, 2, \dots$, we must have

$$\sum_{i=\lceil \frac{j}{2s} \rceil}^{i^*} d_{2i,j} \geq \frac{1 - \left(\frac{1+a_0}{2}\right)^2}{2 - \left(\frac{1+a_0}{2}\right)^2} \sum_{i=\lceil \frac{j}{2s} \rceil}^{\infty} d_{2i,j},$$

where

$$i^* := \left\lceil \frac{1+a_0}{2(1-a_0)} j \right\rceil.$$

Proof. For all $i \geq i^*$, we have $2i \geq j$ and

$$\frac{2i}{2i-j} a_0 \leq \frac{2i^*}{2i^*-j} a_0 \leq \frac{\frac{1+a_0}{1-a_0} j}{\frac{1+a_0}{1-a_0} j - j} a_0 = \frac{1+a_0}{2} \text{ (by Lemma 59 and } 2i \geq 2i^* \geq \frac{1+a_0}{1-a_0} j \text{)}.$$

By Lemma 64, we thus have

$$\frac{d_{2i+2,j}}{d_{2i,j}} \leq \left(\frac{1+a_0}{2} \right)^2 \text{ for all } i \geq i^*.$$

Because $d_{2i,j} \geq 0$ for all i and j , we have

$$\sum_{i=i^*+1}^{\infty} d_{2i,j} \leq \sum_{i=i^*}^{\infty} d_{2i,j} \leq d_{2i^*,j} \sum_{k=0}^{\infty} \left(\frac{1+a_0}{2} \right)^{2k} = \frac{d_{2i^*,j}}{1 - \left(\frac{1+a_0}{2}\right)^2} \leq \frac{1}{1 - \left(\frac{1+a_0}{2}\right)^2} \sum_{i=\lceil \frac{j}{2s} \rceil}^{i^*} d_{2i,j}.$$

Therefore, we have

$$\left(1 + \frac{1}{1 - \left(\frac{1+a_0}{2}\right)^2} \right) \sum_{i=\lceil \frac{j}{2s} \rceil}^{i^*} d_{2i,j} \geq \sum_{i=0}^{i^*} d_{2i,j} + \sum_{i=i^*+1}^{\infty} d_{2i,j} = \sum_{i=\lceil \frac{j}{2s} \rceil}^{\infty} d_{2i,j},$$

i.e.,

$$\sum_{i=\lceil \frac{j}{2s} \rceil}^{i^*} d_{2i,j} \geq \frac{1 - \left(\frac{1+a_0}{2}\right)^2}{2 - \left(\frac{1+a_0}{2}\right)^2} \sum_{i=\lceil \frac{j}{2s} \rceil}^{\infty} d_{2i,j}.$$

□

Recall that $u_m(x)$ denotes the Taylor coefficients of $x^{\frac{\pi - \arccos(x)}{2\pi}}$. The following lemma states that $u_{2k}(x)$ is monotone decreasing with respect to k . Further, it estimates the decreasing speed. We draw the curve of $u_k(x)$ with respect to k in Fig. 11.

Lemma 66. When $k \geq 1$, we have

$$u_{2i}(x) \geq u_{2k}(x) \text{ for all } i \in \{1, 2, \dots, k\},$$

and

$$\frac{u_{2(k+m)}(x)}{u_{2k}(x)} \geq \frac{1}{\left(2 + \frac{2m}{k}\right)^2} \text{ for all } m \in \mathbb{Z}_{\geq 0}.$$

Proof. By Lemma 17, we have $u_0(x) = 0$, $u_1(x) = \frac{1}{4}$, and for all $k \geq 1$, we have

$$u_{2k}(x) = \frac{1}{2\pi} \frac{(2k-2)!}{((k-1)!)^2} \frac{4}{2k-1} \frac{1}{2^{2k}}, \quad (111)$$

$$u_{2k+1}(x) = 0. \quad (112)$$

(We also plot the curve of $u_k(x)$ with respect to k in Fig. 11, so that we can observe the general trend that is consistent with the statement of this lemma. We continue with the precise proof of this lemma below.)

By Eq. (111), we have

$$\frac{u_{2k+2}(x)}{u_{2k}(x)} = \frac{2k(2k-1)(2k-1)}{k^2(2k+1)} \cdot \frac{1}{4} = \frac{(2k-1)^2}{2k \cdot (2k+1)}. \quad (113)$$

Because $\frac{(2k-1)^2}{2k \cdot (2k+1)} \leq 1$, we know that $u_{2k}(x)$ is monotone decreasing with respect to k . Therefore, we have

$$u_{2i}(x) \geq u_{2k}(x) \text{ for all } i \in \{1, 2, \dots, k\}.$$

Iterating Eq. (113), we have

$$\begin{aligned} \frac{u_{2(k+m)}(x)}{u_{2k}(x)} &= \prod_{i=0}^{m-1} \frac{(2k+2i-1)^2}{(2k+2i)(2k+2i+1)} \\ &\geq \prod_{i=0}^{m-1} \frac{(2k+2i-1)^2}{(2k+2i+1)^2} \\ &= \frac{(2k-1)^2}{(2k+2m-1)^2} \\ &\geq \frac{k^2}{(2k+2m)^2} \text{ (because } 2k-1 \geq k \text{ due to } k \geq 1) \\ &= \frac{1}{\left(2 + \frac{2m}{k}\right)^2}. \end{aligned}$$

□

Lemma 67. Under Condition 2, for any $j = 1, 2, \dots$, we must have

$$\frac{u_j(K(x))}{u_{2\lceil j/2 \rceil}(x)} \geq \left(\frac{1}{2} \cdot \frac{1-a_0}{3-a_0}\right)^2 \frac{1 - \left(\frac{1+a_0}{2}\right)^2}{2 - \left(\frac{1+a_0}{2}\right)^2} e^j.$$

Proof. Consider i^* defined in Lemma 65, i.e., $i^* = \left\lceil \frac{1+a_0}{2(1-a_0)} j \right\rceil$. Let $k = \lceil \frac{j}{2} \rceil$ and $m = i^* - k$. We have

$$\begin{aligned} \frac{2m}{k} &= \frac{2i^*}{k} - 2 = \frac{2 \left\lceil \frac{1+a_0}{2(1-a_0)} j \right\rceil}{\lceil \frac{j}{2} \rceil} - 2 \\ &\leq \frac{2 \left(\frac{1+a_0}{2(1-a_0)} j + 1 \right)}{\frac{j}{2}} - 2 \quad \text{(because } \lceil \alpha \rceil \in [\alpha, \alpha + 1]) \\ &= \frac{2(1+a_0)}{1-a_0} + \frac{4}{j} - 2 \\ &\leq \frac{2(1+a_0)}{1-a_0} + 4 - 2 \quad \text{(because } j \geq 1) \\ &= \frac{4}{1-a_0}. \end{aligned}$$

By Lemma 66, we then have

$$\frac{u_{2i^*}(x)}{u_{2\lceil j/2\rceil}(x)} = \frac{u_{2(k+m)}(x)}{u_{2k}(x)} \geq \frac{1}{\left(2 + \frac{2m}{k}\right)^2} \geq \frac{1}{\left(2 + \frac{4}{1-a_0}\right)^2} = \left(\frac{1}{2} \cdot \frac{1-a_0}{3-a_0}\right)^2.$$

By the first part of Lemma 66, $u_{2i} \geq u_{2i^*}$ for all $i = 1, 2, \dots, i^*$. We thus have

$$\frac{u_{2i}(x)}{u_{2\lceil j/2\rceil}(x)} \geq \left(\frac{1}{2} \cdot \frac{1-a_0}{3-a_0}\right)^2 \text{ for all } i = 1, 2, \dots, i^*. \quad (114)$$

Notice that

$$\begin{aligned} & K(x) \frac{\pi - \arccos(K(x))}{2\pi} \\ &= u_1(x)K(x) + \sum_{i=1}^{\infty} u_{2i}(x) (K(x))^{2i} \\ &= u_1(x)K(x) + \sum_{i=1}^{\infty} u_{2i}(x) \sum_{j=0}^{\infty} d_{2i,j} x^j \text{ (by Lemma 27)} \\ &= \frac{1}{4}K(x) + \sum_{j=0}^{\infty} \left(\sum_{i=\lceil \frac{j}{2s} \rceil}^{\infty} u_{2i}(x) \cdot d_{2i,j} \right) x^j \text{ (since } d_{2i,j} = 0 \text{ when } j > 2i \cdot s \text{)}. \end{aligned}$$

Therefore,

$$u_j(K(x)) = \frac{a_j}{4} + \sum_{i=\lceil \frac{j}{2s} \rceil}^{\infty} u_{2i}(x) \cdot d_{2i,j} \text{ for all } j \in \mathbb{Z}_{\geq 0}. \quad (115)$$

By Eq. (115), we thus have

$$\begin{aligned} \frac{u_j(K(x))}{u_{2\lceil j/2\rceil}(x)} &\geq \frac{1}{u_{2\lceil j/2\rceil}(x)} \sum_{i=\lceil \frac{j}{2s} \rceil}^{\infty} u_{2i}(x) d_{2i,j} \text{ (notice that } a_i \geq 0 \text{ for all } i \in \mathbb{Z}_{\geq 0} \text{ by Condition 2)} \\ &\geq \frac{1}{u_{2\lceil j/2\rceil}(x)} \sum_{i=\lceil \frac{j}{2s} \rceil}^{i^*} u_{2i}(x) d_{2i,j} \\ &\geq \left(\frac{1}{2} \cdot \frac{1-a_0}{3-a_0}\right)^2 \sum_{i=\lceil \frac{j}{2s} \rceil}^{i^*} d_{2i,j} \text{ (by Eq. (114))} \\ &\geq \left(\frac{1}{2} \cdot \frac{1-a_0}{3-a_0}\right)^2 \frac{1 - \left(\frac{1+a_0}{2}\right)^2}{2 - \left(\frac{1+a_0}{2}\right)^2} \sum_{i=\lceil \frac{j}{2s} \rceil}^{\infty} d_{2i,j} \text{ (by Lemma 65)} \\ &= \left(\frac{1}{2} \cdot \frac{1-a_0}{3-a_0}\right)^2 \frac{1 - \left(\frac{1+a_0}{2}\right)^2}{2 - \left(\frac{1+a_0}{2}\right)^2} e_j \text{ (by Eq. (107)).} \end{aligned}$$

□

By Lemma 67 and Lemma 62, we can conclude the lower bound in Lemma 60. Next we will prove the upper bound in Lemma 60.

Lemma 68. *Under Condition 2, for any $j = 1, 2, \dots$, we must have*

$$\frac{u_j(K(x)) - \frac{a_j}{4}}{u_{2\lceil j/2\rceil}(x)} \leq (2 + 2s)^2 e_j.$$




Figure 11: The curve of $u_k(x)$ with respect to k . Notice that $u_k(x) = 0$ when $k = 0, 3, 5, 7, \dots$.

Proof. Let $k = \lceil \frac{j}{2s} \rceil$ and $m = \lceil \frac{j}{2} \rceil - \lceil \frac{j}{2s} \rceil$. Thus, we have

$$\frac{m}{k} = \frac{\lceil \frac{j}{2} \rceil}{\lceil \frac{j}{2s} \rceil} - 1 \leq \frac{\frac{j}{2} + 1}{\lceil \frac{j}{2s} \rceil} - 1 = \frac{\frac{j}{2}}{\lceil \frac{j}{2s} \rceil} + \left(\frac{1}{\lceil \frac{j}{2s} \rceil} - 1 \right) \leq s$$

(noting that $\lceil \frac{j}{2s} \rceil \geq 1$ since $j \geq 1$).

By Lemma 66, we then have

$$\frac{u_{2\lceil j/2 \rceil}(x)}{u_{2\lceil j/(2s) \rceil}(x)} = \frac{u_{2(k+m)}(x)}{u_{2k}(x)} \geq \frac{1}{\left(2 + \frac{2m}{k}\right)^2} \text{ (by Lemma 66)}$$

$$\geq \frac{1}{(2 + 2s)^2}.$$

By the first part of Lemma 66, we further have

$$u_{2i}(x) \leq (2 + 2s)^2 u_{2\lceil j/2 \rceil}(x) \text{ for all } 2i \geq 2 \lceil \frac{j}{2s} \rceil.$$

Thus, we have

$$u_j(K(x)) - \frac{a_j}{4} = \sum_{i=\lceil \frac{j}{2s} \rceil}^{\infty} u_{2i}(x) \cdot d_{2i,j} \text{ (by Eq. (115))}$$

$$\leq (2 + 2s)^2 u_{2\lceil j/2 \rceil}(x) \sum_{i=\lceil \frac{j}{2s} \rceil}^{\infty} d_{2i,j}$$

$$= (2 + 2s)^2 u_{2\lceil j/2 \rceil}(x) e_j \text{ (by Eq. (107)).}$$

The result of the lemma thus follows. □

Combining Lemma 67, Lemma 68, and Lemma 62, the result of Lemma 60 thus follows.

M.3 Expansion for an infinite-degree polynomial

We now return to the proof of the third statement of Lemma 51, where the polynomial $K(x)$ has infinite terms. We inherit notations a_i, b_i, u_i, d_i, e_i for the finite polynomial case in Appendix M.2. Further, we introduce some additional notations as follows.

Define

$$\tilde{b}_i := \frac{b_i}{1 - b_0}, \text{ for all } i \in \mathbb{Z}_{\geq 0}, \quad (116)$$

and

$$T_k := \sum_{i=k+1}^{\infty} \tilde{b}_i, \quad k \in \mathbb{Z}_{\geq 0}. \quad (117)$$

By Condition 2, Eq. (103), and Eq. (116), we have

$$T_k \geq 0 \text{ for all } k \in \mathbb{Z}_{\geq 0}. \quad (118)$$

Define for any $k \in \{1, 2, \dots\}$,

$$L_k := \min \{e_j \mid j = 0, 1, \dots, k-1\} \cup \{1\} - \sum_{i=k}^{\infty} T_i. \quad (119)$$

Lemma 69. *For any $i \in \mathbb{Z}_{\geq 0}$ and any $k \in \{1, 2, \dots\}$, we must have $e_i \geq L_k$. Notice that the indices i and k are not required to be equal.*

We prove Lemma 69 in Appendix M.3.1.

In order to prove the third statement of Lemma 51, by Lemma 52, we only need to lower bound $\frac{u_j(K(x))}{u_{2\lceil j/2 \rceil}(x)}$. Notice that Lemma 67 still holds when $s \rightarrow \infty$ (the proof of it will be exactly the same after replacing $\lceil j/2s \rceil$ by zero). Therefore, we only need to prove that e_i is lower bounded by a positive constant. By Lemma 69, if we can find $L_k > 0$ for some k , then we are done. By Eq. (119), In order to calculate the exact value of L_k , we need to find a way to calculate the exact value of $\sum_{k=0}^{\infty} T_k$, which is provided by the following lemma.

Lemma 70. $\sum_{k=0}^{\infty} T_k = \frac{2}{1-a_0^2} \frac{\partial K(x)}{\partial x} \Big|_{x=1}$.

Proof. We have

$$\begin{aligned} \sum_{k=0}^{\infty} T_k &= \sum_{k=0}^{\infty} \sum_{i=k+1}^{\infty} \tilde{b}_i \text{ (by Eq. (117))} \\ &= \sum_{i=1}^{\infty} i \cdot \tilde{b}_i \\ &= \frac{1}{1-b_0} \sum_{i=1}^{\infty} i \cdot b_i \text{ (by Eq. (116))} \\ &= \frac{1}{1-a_0^2} \sum_{i=1}^{\infty} i \cdot b_i \text{ (notice that } b_0 = a_0^2 \text{ by Eq. (103))} \\ &= \frac{1}{1-a_0^2} \frac{\partial (\sum_{i=0}^{\infty} b_i x^i)}{\partial x} \Big|_{x=1} \\ &= \frac{1}{1-a_0^2} \frac{\partial (K(x))^2}{\partial x} \Big|_{x=1} \text{ (by Eq. (103))} \\ &= \frac{2K(1)}{1-a_0^2} \frac{\partial K(x)}{\partial x} \Big|_{x=1} \\ &= \frac{2}{1-a_0^2} \frac{\partial K(x)}{\partial x} \Big|_{x=1} \text{ (by Condition 2).} \end{aligned}$$

□

Now we consider the case of 3-layer without bias, i.e., the case when the polynomial $K(x)$ is $2d \cdot K^{\text{RF}}(x) = \frac{\sqrt{1-x^2} + (\pi - \arccos(x))x}{\pi}$. After calculation (details in Appendix M.3.2), we have $L_3 \approx 0.069 > 0$, which completes the proof of Proposition 47.

M.3.1 Proof of Lemma 69

We first prove some useful lemmas.

Lemma 71. (i) $\sum_{i=1}^{\infty} \tilde{b}_i = 1$. (ii) $e_k = \sum_{i=0}^{k-1} \tilde{b}_{k-i} e_i$ for all $k \in \{1, 2, \dots\}$. (iii) $e_k \leq e_0 = \frac{1}{1-b_0}$ for all $k \in \mathbb{Z}_{\geq 0}$.

Proof. Note that

$$\begin{aligned} \sum_{i=1}^{\infty} \tilde{b}_i &= \frac{\sum_{i=1}^{\infty} b_i}{1-b_0} \text{ (by Eq. (116))} \\ &= 1 \quad \left(\text{since } \sum_{i=1}^{\infty} b_i = 1 - b_0 \text{ because } b_0 + \sum_{i=1}^{\infty} b_i = (K(1))^2 = 1 \right). \end{aligned}$$

Because

$$\frac{1}{1-(K(x))^2} = 1 + (K(x))^2 \cdot \frac{1}{1-(K(x))^2},$$

we have

$$\sum_{j=0}^{\infty} e_j x^j = 1 + \left(\sum_{i=0}^{\infty} a_i x^i \right)^2 \cdot \left(\sum_{j=0}^{\infty} e_j x^j \right) = 1 + \left(\sum_{i=0}^{\infty} b_i x^i \right) \cdot \left(\sum_{j=0}^{\infty} e_j x^j \right).$$

Comparing the coefficient of x^j on both sides, we have

$$\begin{aligned} e_0 &= 1 + b_0 e_0, \\ e_1 &= b_1 e_0 + b_0 e_1, \\ e_2 &= b_2 e_0 + b_1 e_1 + b_0 e_2, \\ &\vdots \end{aligned}$$

We thus have

$$\left\{ \begin{array}{l} e_0 = \frac{1}{1-b_0}, \\ e_1 = \tilde{b}_1 e_0, \\ e_2 = \tilde{b}_2 e_0 + \tilde{b}_1 e_1 = (\tilde{b}_2 + \tilde{b}_1^2) e_0, \\ e_3 = \tilde{b}_3 e_0 + \tilde{b}_2 e_1 + \tilde{b}_1 e_2 = (\tilde{b}_3 + 2\tilde{b}_2 \tilde{b}_1 + \tilde{b}_1^3) e_0, \\ e_4 = \tilde{b}_4 e_0 + \tilde{b}_3 e_1 + \tilde{b}_2 e_2 + \tilde{b}_1 e_3 = (\tilde{b}_4 + 2\tilde{b}_1 \tilde{b}_3 + \tilde{b}_2^2 + 3\tilde{b}_1^2 \tilde{b}_2 + \tilde{b}_1^4) e_0, \\ \vdots \\ e_k = \sum_{i=0}^{k-1} \tilde{b}_{k-i} e_i, \\ \vdots \end{array} \right. \quad (120)$$

We now prove that $e_k \leq e_0$ for all $k \in \mathbb{Z}_{\geq 0}$ by mathematical induction. Suppose that for all $i \leq k$, we already have $e_i \leq e_0$ (which is obviously true when $k = 0$). Thus, we have

$$e_{k+1} = \sum_{i=0}^k \tilde{b}_{k+1-i} e_i \leq e_0 \sum_{i=0}^k \tilde{b}_{k+1-i} \leq e_0 \sum_{i=0}^{\infty} \tilde{b}_i = e_0.$$

By mathematical induction, we thus have $e_i \leq e_0$ for all $i \in \mathbb{Z}_{\geq 0}$. \square

For any given real number sequence $\Delta := (\Delta_1, \Delta_2, \dots)$, we define a sequence $(e_0^\Delta, e_1^\Delta, e_2^\Delta, \dots)$ by

$$e_0^\Delta := e_0, \quad e_k^\Delta := \Delta_k + \sum_{i=0}^{k-1} \tilde{b}_{k-i} e_i^\Delta, \quad k \in \{1, 2, \dots\}. \quad (121)$$

Lemma 72. For any $k \in \{1, 2, \dots\}$ and any Δ , we must have

$$e_k^\Delta - e_k = \sum_{i=1}^k \Delta_i \cdot \frac{e_{k-i}}{e_0}. \quad (122)$$

Proof. We prove Eq. (122) by mathematical induction. When $k = 1$, we have $e_1^\Delta = \Delta_1 + \tilde{b}_1 e_0$ (by Eq. (121)) and $e_1 = \tilde{b}_1 e_0$ (by Lemma 71). Thus, Eq. (113) holds when $k = 1$. Suppose that for all $k \in \{1, 2, \dots, l\}$, Eq. (122) holds. We thus have

$$\begin{aligned} e_{l+1}^\Delta - e_{l+1} &= \Delta_{l+1} + \sum_{i=0}^l \tilde{b}_{l+1-i} (e_i^\Delta - e_i) \text{ (by Eq. (121) and Lemma 71)} \\ &= \Delta_{l+1} + \sum_{i=1}^l \tilde{b}_{l+1-i} (e_i^\Delta - e_i) \text{ (notice that } e_0^\Delta = e_0) \\ &= \Delta_{l+1} + \sum_{i=1}^l \tilde{b}_{l+1-i} \sum_{j=1}^i \Delta_j \frac{e_{i-j}}{e_0} \text{ (applying Eq. (122) by induction hypothesis).} \end{aligned} \quad (123)$$

Notice that

$$\begin{aligned} \sum_{i=1}^l \sum_{j=1}^i \tilde{b}_{l+1-i} \cdot \Delta_j \cdot e_{i-j} &= \sum_{j=1}^l \Delta_j \sum_{i=j}^l \tilde{b}_{l+1-i} e_{i-j} \text{ (by re-organizing terms)} \\ &= \sum_{j=1}^l \Delta_j \sum_{i=0}^{l-j} \tilde{b}_{l+1-i-j} e_i \text{ (replacing } i-j \text{ by } i). \end{aligned}$$

Plugging the above equation into Eq. (123), we then have

$$\begin{aligned} e_{l+1}^\Delta - e_{l+1} &= \Delta_{l+1} + \frac{1}{e_0} \sum_{j=1}^l \Delta_j \sum_{i=0}^{l-j} \tilde{b}_{l+1-i-j} e_i \\ &= \Delta_{l+1} + \frac{1}{e_0} \sum_{j=1}^l \Delta_j \cdot e_{l+1-j} \text{ (by Lemma 71)} \\ &= \frac{1}{e_0} \sum_{j=1}^{l+1} \Delta_j \cdot e_{l+1-j}, \end{aligned}$$

i.e., Eq. (122) also holds for $k = l + 1$. Thus, the mathematical induction is completed and the result of this lemma thus follows. \square

Now we are ready to prove Lemma 69.

Proof of Lemma 69. Let

$$\Delta_i = \begin{cases} 0, & \text{if } i < k, \\ T_i, & \text{if } i \geq k. \end{cases}$$

We first prove by mathematical induction that

$$e_i^\Delta \geq \min\{e_j \mid j = 0, 1, \dots, k-1\} \cup \{1\} \text{ for all } i \in \mathbb{Z}_{\geq 0}. \quad (124)$$

Towards this end, note that because $\Delta_i = 0$ for all $i < k$, we know from Lemma 72 that $e_i^\Delta = e_i$. Hence, Eq. (124) trivially holds for all $i \in \{0, 1, \dots, k-1\}$. Suppose that Eq. (124) holds for all $i \leq l \in \mathbb{Z}_{\geq 0}$, where $l \geq k-1$ denotes the index of the induction hypothesis. In order to finish the

mathematical induction, we only need to prove that Eq. (124) holds for $i = l + 1$. To this end, we have

$$\begin{aligned}
e_{l+1}^\Delta &= T_{l+1} + \sum_{i=0}^l \tilde{b}_{l+1-i} e_i^\Delta \text{ (by Eq. (121))} \\
&= \sum_{i=l+2}^{\infty} \tilde{b}_i + \sum_{i=1}^{l+1} \tilde{b}_i e_{l+1-i}^\Delta \text{ (by Eq. (117))} \\
&\geq (\min\{e_j^\Delta \mid j = 0, 1, \dots, l\} \cup \{1\}) \cdot \sum_{i=1}^{\infty} \tilde{b}_i \\
&= \min\{e_j^\Delta \mid j = 0, 1, \dots, l\} \cup \{1\} \text{ (by Lemma 71)} \\
&\geq \min\{e_j \mid j = 0, 1, \dots, k-1\} \cup \{1\} \text{ (by induction hypothesis)}.
\end{aligned}$$

Thus, Eq. (124) holds by mathematical induction. We thus have

$$\begin{aligned}
e_i &= e_i^\Delta - \sum_{j=1}^i \Delta_j \cdot \frac{e_{i-j}}{e_0} \text{ (by Lemma 72)} \\
&\geq e_i^\Delta - \sum_{j=1}^i \Delta_j \text{ (since } e_{i-j} \leq e_0 \text{ by Lemma 71)} \\
&= e_i^\Delta - \sum_{j=k}^i T_j \\
&\geq \min\{e_j \mid j = 0, 1, \dots, k-1\} \cup \{1\} - \sum_{j=k}^{\infty} T_j \text{ (by Eq. (124) and Eq. (118))} \\
&= L_k.
\end{aligned}$$

The result of this lemma thus follows. \square

M.3.2 Calculate L_3 for 3-layer without bias

Coefficients of Taylor expansion of $K(x) = 2d \cdot K^{\text{RF}}(x) = \frac{\sqrt{1-x^2} + (\pi - \arccos(x))x}{\pi}$ can be derived from Lemma 17, i.e.,

$$K(x) = \frac{1}{\pi} \left(1 + \frac{\pi}{2}x + \sum_{k=0}^{\infty} \frac{2(2k)!}{(k+1)(2k+1)(k!)^2} \left(\frac{x}{2}\right)^{2k+2} \right). \quad (125)$$

By Eq. (125), We can calculate values of a_i , b_i , and \tilde{b}_i for $i = 0, 1, 2$ by their definitions.

$$\begin{aligned}
a_0 &= \frac{1}{\pi}, \quad a_1 = \frac{1}{2}, \quad a_2 = \frac{1}{2\pi}, \\
b_0 &= a_0^2 = \frac{1}{\pi^2}, \quad b_1 = 2a_0a_1 = \frac{1}{\pi}, \quad b_2 = 2a_2a_0 + a_1^2 = \frac{1}{\pi^2} + \frac{1}{4}, \\
\tilde{b}_0 &= \frac{1}{\pi^2 - 1}, \quad \tilde{b}_1 = \frac{\pi}{\pi^2 - 1}, \quad \tilde{b}_2 = \frac{1}{\pi^2 - 1} + \frac{\pi^2}{4(\pi^2 - 1)}.
\end{aligned}$$

Then, we calculate the values of e_i by Eq. (120).

$$e_0 = \frac{\pi^2}{\pi^2 - 1} \approx 1.11, \quad e_1 = \frac{\pi^3}{(\pi^2 - 1)^2} \approx 0.39, \quad e_2 = \frac{(2\pi^2 - 1)\pi^2}{(\pi^2 - 1)^3} + \frac{\pi^4}{4(\pi^2 - 1)^2} \approx 0.57.$$

Next, we calculate $\sum_{i=0}^{\infty} T_i$ by Lemma 70.

$$\begin{aligned}
\sum_{i=0}^{\infty} T_i &= \frac{2}{1-a_0^2} \frac{\partial \frac{\sqrt{1-x^2} + (\pi - \arccos(x))x}{\pi}}{\partial x} \Big|_{x=1} \\
&= \frac{2}{1-a_0^2} \frac{1}{\pi} \left(-\frac{x}{\sqrt{1-x^2}} + (\pi - \arccos(x)) + \frac{x}{\sqrt{1-x^2}} \right) \\
&\quad \text{(notice that } \frac{\partial \sqrt{1-x^2}}{\partial x} = -\frac{x}{\sqrt{1-x^2}} \text{ and } \frac{\partial}{\partial x} \arccos(x) = -\frac{1}{\sqrt{1-x^2}} \text{)} \\
&= \frac{2}{1-a_0^2} \frac{\pi - \arccos(x)}{\pi} \Big|_{x=1} = \frac{2\pi^2}{\pi^2 - 1}.
\end{aligned}$$

By Eq. (117) and Lemma 71(i), we thus have $T_j = \sum_{i=1}^{\infty} \tilde{b}_i - \sum_{i=1}^j \tilde{b}_i = 1 - \sum_{i=1}^j \tilde{b}_i$. Therefore, we have

$$T_0 = 1, T_1 = 1 - \frac{\pi}{\pi^2 - 1}, T_2 = 1 - \frac{\pi}{\pi^2 - 1} - \left(\frac{1}{\pi^2 - 1} + \frac{\pi^2}{4(\pi^2 - 1)} \right).$$

Now we are ready to calculate L_3 by Eq. (119).

$$\begin{aligned}
L_3 &= e_1 + T_0 + T_1 + T_2 - \sum_{i=0}^{\infty} T_i \\
&= \frac{\pi^3}{(\pi^2 - 1)^2} + 3 - \frac{2\pi}{\pi^2 - 1} - \left(\frac{1}{\pi^2 - 1} + \frac{\pi^2}{4(\pi^2 - 1)} \right) - \frac{2\pi^2}{\pi^2 - 1} \\
&\approx 0.069.
\end{aligned}$$